



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : C12N 15/12, C07K 14/47, A61K 38/17, C07K 16/18, G01N 33/53, C12Q 1/68, C12N 15/62, 15/11		A2	(11) International Publication Number: <b>WO 00/53753</b>  (43) International Publication Date: 14 September 2000 (14.09.00)
(21) International Application Number: PCT/US00/00219 (22) International Filing Date: 5 January 2000 (05.01.00)		(71) Applicant (for all designated States except US): GENENTECH, INC. [US/US]; 1 DNA Way, South San Francisco, CA 94080-4990 (US).	
(30) Priority Data:		(72) Inventors; and (75) Inventors/Applicants (for US only): ASHKENAZI, Avi, J. [US/US]; 1456 Terrytown Street, San Mateo, CA 94402 (US). BAKER, Kevin, P. [GB/US]; 14006 Indian Run Drive, Damestown, MD 20878 (US). FERRARA, Napoleone [US/US]; 2090 Pacific Avenue, #704, San Francisco, CA 94109 (US). GERBER, Hanspeter [CH/US]; 1121 Tennessee Street, #5, San Francisco, CA 94107 (US). GODDARD, Audrey [CA/US]; 110 Congo Street, San Francisco, CA 94131 (US). GODOWSKI, Paul, J. [US/US]; 2627 Easton Drive, Burlingame, CA 94010 (US). GURNEY, Austin, L. [US/US]; 1 Debbie Lane, Belmont, CA 94002 (US). HILLAN, Kenneth, J. [GB/US]; 64 Seward Street, San Francisco, CA 94114 (US). KUO, Sophia, S. [US/US]; 59 Surrey Street, #3, San Francisco, CA 94131 (US). MARK, Melanie, R. [US/US]; 958 B Chula Vista Avenue, Burlingame, CA 94010 (US). MARSTERS, Scot, A. [US/US]; 990 Cherry Street, San Carlos, CA 94070 (US). PAONI, Nicholas, F. [US/US]; 1756 Terrace Drive, Belmont, CA 94002 (US). PITT, Robert, M. [US/US]; 1110 Liberty Street, El Cerrito, CA 94530 (US). WATANABE, Colin, K. [US/US]; 128 Corliss Drive, Moraga, CA 94556 (US). WILLIAMS, P., Mickey [US/US]; 509 Alto Avenue, Half Moon Bay, CA 94019 (US). WOOD, William, I. [US/US]; 35 Southdown Court, Hillsborough, CA 94010 (US).	
PCT/US99/05028 8 March 1999 (08.03.99) US 60/123,957 12 March 1999 (12.03.99) US 60/134,287 14 May 1999 (14.05.99) US PCT/US99/12252 2 June 1999 (02.06.99) US 60/141,037 23 June 1999 (23.06.99) US 60/144,758 20 July 1999 (20.07.99) US 60/145,698 26 July 1999 (26.07.99) US PCT/US99/20111 1 September 1999 (01.09.99) US PCT/US99/20594 8 September 1999 (08.09.99) US PCT/US99/21090 15 September 1999 (15.09.99) US PCT/US99/21547 15 September 1999 (15.09.99) US PCT/US99/23089 5 October 1999 (05.10.99) US PCT/US99/28313 30 November 1999 (30.11.99) US PCT/US99/28409 30 November 1999 (30.11.99) US PCT/US99/28564 2 December 1999 (02.12.99) US PCT/US99/28565 2 December 1999 (02.12.99) US Not furnished 20 December 1999 (20.12.99) US		(74) Agents: BARNES, Elizabeth, M. et al.; Genentech, Inc., 1 DNA Way, South San Francisco, CA 94080-4990 (US).	
		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
		Published Without international search report and to be republished upon receipt of that report.	
(54) Title: PROMOTION OR INHIBITION OF ANGIOGENESIS AND CARDIOVASCULARIZATION			
(57) Abstract			
<p>Compositions and methods are disclosed for stimulating or inhibiting angiogenesis and/or cardiovascularization in mammals, including humans. Pharmaceutical compositions are based on polypeptides or antagonists thereto that have been identified for one or more of these uses. Disorders that can be diagnosed, prevented, or treated by the compositions herein include trauma such as wounds, various cancers, and disorders of the vessels including atherosclerosis and cardiac hypertrophy. In addition, the present invention is directed to novel polypeptides and to nucleic acid molecules encoding those polypeptides. Also provided herein are vectors and host cells comprising those nucleic acid sequences, chimeric polypeptide molecules comprising the polypeptides of the present invention fused to heterologous polypeptide sequences, antibodies which bind to the polypeptides of the present invention and to methods for producing the polypeptides of the present invention.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## PROMOTION OR INHIBITION OF ANGIOGENESIS AND CARDIOVASCULARIZATION

### Background of the Invention

#### 5     Field of the Invention

The present invention relates to compositions and methods useful for promoting or inhibiting angiogenesis and/or cardiovascularization in mammals in need of such biological effect. This includes the diagnosis and treatment of cardiovascular disorders as well as oncological disorders.

#### 10    Description of Background

##### A.     Cardiac Disorders and Factors

Heart failure affects approximately five million Americans, and new cases of heart failure number about 400,000 each year. It is the single most frequent cause of hospitalization for people age 65 and older in the United States. Recent advances in the management of acute cardiac diseases, including acute myocardial infarction, are  
15    resulting in an expanding patient population that will eventually develop chronic heart failure. From 1979 to 1995, hospitalizations for congestive heart failure (CHF) rose from 377,000 to 872,000 (a 130 percent increase) and CHF deaths increased 116 percent.

CHF is a syndrome characterized by left ventricular dysfunction, reduced exercise tolerance, impaired quality of life, and markedly shortened life expectancy. The sine qua non of heart failure is an inability of the heart  
20    to pump blood at a rate sufficient to meet the metabolic needs of the body's tissues (in other words, there is insufficient cardiac output).

At least four major compensatory mechanisms are activated in the setting of heart failure to boost cardiac output, including peripheral vasoconstriction, increased heart rate, increased cardiac contractility, and increased plasma volume. These effects are mediated primarily by the sympathetic nervous system and the renin-angiotensin system. See, Eichhorn, American Journal of Medicine, 104: 163-169 (1998). Increased output from the  
25    sympathetic nervous system increases vascular tone, heart rate, and contractility. Angiotensin II elevates blood pressure by 1) directly stimulating vascular smooth muscle contraction, 2) promoting plasma volume expansion by stimulating aldosterone and antidiuretic hormone secretion, 3) stimulating sympathetic-mediated vascular tone, and 4) catalyzing the degradation of bradykinin, which has vasodilatory and natriuretic activity. See, review by Brown and Vaughan, Circulation, 97: 1411-1420 (1998). As noted below, angiotensin II may also have directly deleterious effects on the heart by promoting myocyte necrosis (impairing systolic function) and intracardiac fibrosis (impairing diastolic and in some cases systolic function). See, Weber, Circulation, 96: 4065-4082 (1998).  
30    

A consistent feature of congestive heart failure (CHF) is cardiac hypertrophy, an enlargement of the heart that is activated by both mechanical and hormonal stimuli and enables the heart to adapt to demands for increased  
35    cardiac output. Morgan and Baker, Circulation, 83: 13-25 (1991). This hypertrophic response is frequently

associated with a variety of distinct pathological conditions such as hypertension, aortic stenosis, myocardial infarction, cardiomyopathy, valvular regurgitation, and intracardiac shunt, all of which result in chronic hemodynamic overload.

Hypertrophy is generally defined as an increase in size of an organ or structure independent of natural growth that does not involve tumor formation. Hypertrophy of the heart is due either to an increase in the mass of the individual cells (myocytes), or to an increase in the number of cells making up the tissue (hyperplasia), or both. While the enlargement of an embryonic heart is largely dependent on an increase in myocyte number (which continues until shortly after birth), post-natal cardiac myocytes lose their proliferative capacity. Further growth occurs through hypertrophy of the individual cells.

Adult myocyte hypertrophy is initially beneficial as a short term response to impaired cardiac function by permitting a decrease in the load on individual muscle fibers. With severe, long-standing overload, however, the hypertrophied cells begin to deteriorate and die. Katz, "Heart Failure", in: Katz A.M. ed., Physiology of the Heart (New York: Raven Press, 1992) pp. 638-668. Cardiac hypertrophy is a significant risk factor for both mortality and morbidity in the clinical course of heart failure. Katz, Trends Cardiovasc. Med., 5: 37-44 (1995). For further details of the causes and pathology of cardiac hypertrophy see, e.g., Heart Disease, A Textbook of Cardiovascular Medicine, Braunwald, E. ed. (W.B. Saunders Co., 1988), Chapter 14, "Pathophysiology of Heart Failure."

On a cellular level, the heart is composed of myocytes and surrounding support cells, generically called non-myocytes. While non-myocytes are primarily fibroblast/mesenchymal cells, they also include endothelial and smooth muscle cells. Indeed, although myocytes make up most of the adult myocardial mass, they represent only about 30% of the total cell numbers present in heart. In response to hormonal, physiological, hemodynamic, and pathological stimuli, adult ventricular muscle cells can adapt to increased workloads through the activation of a hypertrophic process. This response is characterized by an increase in myocyte cell size and contractile protein content of individual cardiac muscle cells, without concomitant cell division and activation of embryonic genes, including the gene for atrial natriuretic peptide (ANP). Chien *et al.*, FASEB J., 5: 3037-3046 (1991); Chien *et al.*, Annu. Rev. Physiol., 55: 77-95 (1993). An increment in myocardial mass as a result of an increase in myocyte size that is associated with an accumulation of interstitial collagen within the extracellular matrix and around intramyocardial coronary arteries has been described in left ventricular hypertrophy secondary to pressure overload in humans. Caspari *et al.*, Cardiovasc. Res., 11: 554-558 (1977); Schwarz *et al.*, Am. J. Cardiol., 42: 895-903 (1978); Hess *et al.*, Circulation, 63: 360-371 (1981); Pearlman *et al.*, Lab. Invest., 46: 158-164 (1982).

It has also been suggested that paracrine factors produced by non-myocyte supporting cells may additionally be involved in the development of cardiac hypertrophy, and various non-myocyte derived hypertrophic factors, such as, leukocyte inhibitory factor (LIF) and endothelin, have been identified. Metcalf, Growth Factors, 7: 169-173 (1992); Kurzrock *et al.*, Endocrine Reviews, 12: 208-217 (1991); Inoue *et al.*, Proc. Natl. Acad. Sci. USA, 86: 2863-2867 (1989); Yanagisawa and Masaki, Trends Pharm. Sci., 10: 374-378 (1989); U.S. Patent No. 5,573,762 (issued November 12, 1996). Further exemplary factors that have been identified as potential mediators of cardiac hypertrophy include cardiotrophin-1 (CT-1) (Pennica *et al.*, Proc. Nat. Acad. Sci. USA, 92: 1142-1146 (1995)), catecholamines, adrenocorticosteroids, angiotensin, and prostaglandins.

At present, the treatment of cardiac hypertrophy varies depending on the underlying cardiac disease.



Catecholamines, adrenocorticosteroids, angiotensin, prostaglandins, LIF, endothelin (including endothelin-1, -2, and -3 and big endothelin), and CT-1 are among the factors identified as potential mediators of hypertrophy. For example, beta-adrenergic receptor blocking drugs (beta-blockers, *e.g.*, propranolol, timolol, tertalolol, carteolol, nadolol, betaxolol, penbutolol, acetobutolol, atenolol, metoprolol, carvedilol, etc.) and verapamil have been used extensively in the treatment of hypertrophic cardiomyopathy. The beneficial effects of beta-blockers on symptoms (*e.g.*, chest pain) and exercise tolerance are largely due to a decrease in the heart rate with a consequent prolongation of diastole and increased passive ventricular filling. Thompson *et al.*, Br. Heart J., **44**: 488-98 (1980); Harrison *et al.*, Circulation, **29**: 84-98 (1964). Verapamil has been described to improve ventricular filling and probably reducing myocardial ischemia. Bonow *et al.*, Circulation, **72**: 853-64 (1985).

Nifedipine and diltiazem have also been used occasionally in the treatment of hypertrophic cardiomyopathy. Lorell *et al.*, Circulation, **65**: 499-507 (1982); Betocchi *et al.*, Am. J. Cardiol., **78**: 451-457 (1996). However, because of its potent vasodilating properties, nifedipine may be harmful, especially in patients with outflow obstruction. Disopyramide has been used to relieve symptoms by virtue of its negative inotropic properties. Pollick, N. Engl. J. Med., **307**: 997-999 (1982). In many patients, however, the initial benefits decrease with time. Wigle *et al.*, Circulation, **92**: 1680-1692 (1995). Antihypertensive drug therapy has been reported to have beneficial effects on cardiac hypertrophy associated with elevated blood pressure. Examples of drugs used in antihypertensive therapy, alone or in combination, are calcium antagonists, *e.g.*, nitrendipine; adrenergic receptor blocking agents, *e.g.*, those listed above; angiotensin converting enzyme (ACE) inhibitors such as quinapril, captopril, enalapril, ramipril, benazepril, fosinopril, and lisinopril; diuretics, *e.g.*, chlorothiazide, hydrochlorothiazide, hydroflumethazide, methylchlorothiazide, benzthiazide, dichlorphenamide, acetazolamide, and indapamide; and calcium channel blockers, *e.g.*, diltiazem, nifedipine, verapamil, and nicardipine.

For example, treatment of hypertension with diltiazem and captopril showed a decrease in left ventricular muscle mass, but the Doppler indices of diastolic function did not normalize. Szlachcic *et al.*, Am. J. Cardiol., **63**: 198-201 (1989); Shahi *et al.*, Lancet, **336**: 458-461 (1990). These findings were interpreted to indicate that excessive amounts of interstitial collagen may remain after regression of left ventricular hypertrophy. Rossi *et al.*, Am. Heart J., **124**: 700-709 (1992). Rossi *et al.*, *supra*, investigated the effect of captopril on the prevention and regression of myocardial cell hypertrophy and interstitial fibrosis in pressure overload cardiac hypertrophy, in experimental rats.

Agents that increase cardiac contractility directly (ionotropic agents) were initially thought to benefit patients with heart failure because they improved cardiac output in the short term. However, all positive inotropic agents except digoxigenin have been found to result in increased long-term mortality, in spite of short-term improvements in cardiac performance. Massie, Curr. Op. in Cardiology, **12**: 209-217 (1997); Reddy *et al.*, Curr. Opin. Cardiol., **12**: 233-241 (1997). Beta-adrenergic receptor blockers have recently been advocated for use in heart failure. Evidence from clinical trials suggests that improvements in cardiac function can be achieved without increased mortality, though documented improvements patient survival have not yet been demonstrated. *See also*, U.S. Pat. Nos. 5,935,924, 5,624,806; 5,661,122; and 5,610,134 and WO 95/28173 regarding the use of cardiotropin-1 or antagonists thereof, or growth hormone and/or insulin-like growth factor-I in the treatment of CHF. Another treatment modality is heart transplantation, but this is limited by the availability of donor hearts.

Endothelin is a vasoconstricting peptide comprising 21 amino acids, isolated from swine arterial endothelial culture supernatant and structurally determined. Yanagisawa *et al.*, Nature, 332: 411-415 (1988). Endothelin was later found to exhibit various actions, and endothelin antibodies as endothelin antagonists have proven effective in the treatment of myocardial infarction, renal failure, and other diseases. Since endothelin is present in live bodies and exhibits vasoconstricting action, it is expected to be an endogenous factor involved in the regulation of the circulatory system, and may be associated with hypertension, cardiovascular diseases such as myocardial infarction, and renal diseases such as acute renal failure. Endothelin antagonists are described, for example, in U.S. Pat. No. 5,773,414; JP Pat. Publ. 3130299/1991, EP 457,195; EP 460,679; and EP 552,489. A new endothelin B receptor for identifying endothelin receptor antagonists is described in U.S. Pat. No. 5,773,223.

Current therapy for heart failure is primarily directed to using angiotensin-converting enzyme (ACE) inhibitors, such as captopril, and diuretics. These drugs improve hemodynamic profile and exercise tolerance and reduce the incidence of morbidity and mortality in patients with CHF. Kramer *et al.*, Circulation, 67(4): 807-816 (1983); Captopril Multicenter Research Group, J.A.C.C., 2(4): 755-763 (1983); The CONSENSUS Trial Study Group, N. Engl. J. Med., 316(23): 1429-1435 (1987); The SOLVD Investigators, N. Engl. J. Med., 325(5): 293-302 (1991). Further, they are useful in treating hypertension, left ventricular dysfunction, atherosclerotic vascular disease, and diabetic nephropathy. Brown and Vaughan, *supra*. However, despite proven efficacy, response to ACE inhibitors has been limited. For example, while prolonging survival in the setting of heart failure, ACE inhibitors appear to slow the progression towards end-stage heart failure, and substantial numbers of patients on ACE inhibitors have functional class III heart failure.

Moreover, improvement of functional capacity and exercise time is only small and mortality, although reduced, continues to be high. The CONSENSUS Trial Study Group, N. Engl. J. Med., 316(23): 1429-1453 (1987); The SOLVD Investigators, N. Engl. J. Med., 325(5): 293-302 (1991); Cohn *et al.*, N. Engl. J. Med., 325(5): 303-310 (1991); The Captopril-Digoxin Multicenter Research Group, JAMA, 259(4): 539-544 (1988). Hence, ACE inhibitors consistently appear unable to relieve symptoms in more than 60% of heart failure patients and reduce mortality of heart failure only by approximately 15-20%. For further adverse effects, *see* Brown and Vaughan, *supra*.

An alternative to ACE inhibitors is represented by specific AT<sub>I</sub> receptor antagonists. Clinical studies are planned to compare the efficacy of these two modalities in the treatment of cardiovascular and renal disease. However, animal model data suggests that the ACE/Ang II pathway, while clearly involved in cardiac hypertrophy, is not the only, or even the primary pathway active in this role. Mouse genetic "knockout" models have been made to test individual components of the pathway. In one such model, the primary cardiac receptor for Ang II, AT sub 1A, has been genetically deleted; these mice do not develop hypertrophy when Ang II is given experimentally (confirming the basic success of the model in eliminating hypertrophy secondary to Ang II). However, when the aorta is constricted in these animals (a model of hypertensive cardiac stress), the hearts still become hypertrophic. This suggests that alternative signaling pathways, not depending on this receptor (AT sub 1A), are activated in hypertension. ACE inhibitors would presumably not be able to inhibit these pathways. *See*, Harada *et al.*, Circulation, 97: 1952-1959 (1998). *See also*, Homcy, Circulation, 97: 1890-1892 (1998) regarding the enigma associated with the process and mechanism of cardiac hypertrophy.

About 750,000 patients suffer from acute myocardial infarction (AMI) annually, and approximately one-fourth of all deaths in the United States are due to AMI. In recent years, thrombolytic agents, *e.g.*, streptokinase, urokinase, and in particular tissue plasminogen activator (t-PA) have significantly increased the survival of patients who suffered myocardial infarction. When administered as a continuous intravenous infusion over 1.5 to 4 hours, t-PA produces coronary patency at 90 minutes in 69% to 90% of the treated patients. Topol *et al.*, Am. J. Cardiol., **61**, 723-728 (1988); Neuhaus *et al.*, J. Am. Coll. Cardiol., **12**: 581-587 (1988); Neuhaus *et al.*, J. Am. Coll. Cardiol., **14**: 1566-1569 (1989). The highest patency rates have been reported with high dose or accelerated dosing regimens. Topol, J. Am. Coll. Cardiol., **15**: 922-924 (1990). t-PA may also be administered as a single bolus, although due to its relatively short half-life, it is better suited for infusion therapy. Tebbe *et al.*, Am. J. Cardiol., **64**: 448-453 (1989). A t-PA variant, specifically designed to have longer half-life and very high fibrin specificity, TNK t-PA (a T103N, N117Q, KHRR(296-299)AAAA t-PA variant, Keyt *et al.*, Proc. Natl. Acad. Sci. USA, **91**: 3670-3674 (1994)) is particularly suitable for bolus administration. However, despite all these advances, the long-term prognosis of patient survival depends greatly on the post-infarction monitoring and treatment of the patients, which should include monitoring and treatment of cardiac hypertrophy.

#### B. Growth Factors

Various naturally occurring polypeptides reportedly induce the proliferation of endothelial cells. Among those polypeptides are the basic and acidic fibroblast growth factors (FGF) (Burgess and Maciag, Annual Rev. Biochem., **58**: 575 (1989)), platelet-derived endothelial cell growth factor (PD-ECGF) (Ishikawa *et al.*, Nature, **338**: 557 (1989)), and vascular endothelial growth factor (VEGF). Leung *et al.*, Science, **246**: 1306 (1989); Ferrara and Henzel, Biochem. Biophys. Res. Commun., **161**: 851 (1989); Tischer *et al.*, Biochem. Biophys. Res. Commun., **165**: 1198 (1989); EP 471,754B granted July 31, 1996.

Media conditioned by cells transfected with the human VEGF (hVEGF) cDNA promoted the proliferation of capillary endothelial cells, whereas control cells did not. Leung *et al.*, Science, **246**: 1306 (1989). Several additional cDNAs were identified in human cDNA libraries that encode 121-, 189-, and 206-amino acid isoforms of hVEGF (also collectively referred to as hVEGF-related proteins). The 121-amino acid protein differs from hVEGF by virtue of the deletion of the 44 amino acids between residues 116 and 159 in hVEGF. The 189-amino acid protein differs from hVEGF by virtue of the insertion of 24 amino acids at residue 116 in hVEGF, and apparently is identical to human vascular permeability factor (hVPF). The 206-amino acid protein differs from hVEGF by virtue of an insertion of 41 amino acids at residue 116 in hVEGF. Houck *et al.*, Mol. Endocrin., **5**: 1806 (1991); Ferrara *et al.*, J. Cell. Biochem., **47**: 211 (1991); Ferrara *et al.*, Endocrine Reviews, **13**: 18 (1992); Keck *et al.*, Science, **246**: 1309 (1989); Connolly *et al.*, J. Biol. Chem., **264**: 20017 (1989); EP 370,989 published May 30, 1990.

It is now well established that angiogenesis, which involves the formation of new blood vessels from preexisting endothelium, is implicated in the pathogenesis of a variety of disorders. These include solid tumors and metastasis, atherosclerosis, retrolental fibroplasia, hemangiomas, chronic inflammation, intraocular neovascular syndromes such as proliferative retinopathies, *e.g.*, diabetic retinopathy, age-related macular degeneration (AMD), neovascular glaucoma, immune rejection of transplanted corneal tissue and other tissues, rheumatoid arthritis, and

psoriasis. Folkman *et al.*, J. Biol. Chem., **267**: 10931-10934 (1992); Klagsbrun *et al.*, Annu. Rev. Physiol., **53**: 217-239 (1991); and Garner A., "Vascular diseases", In: Pathobiology of Ocular Disease. A Dynamic Approach, Garner A., Klintworth GK, eds., 2nd Edition (Marcel Dekker, NY, 1994), pp 1625-1710.

In the case of tumor growth, angiogenesis appears to be crucial for the transition from hyperplasia to neoplasia, and for providing nourishment to the growing solid tumor. Folkman *et al.*, Nature, **339**: 58 (1989). The neovascularization allows the tumor cells to acquire a growth advantage and proliferative autonomy compared to the normal cells. Accordingly, a correlation has been observed between density of microvessels in tumor sections and patient survival in breast cancer as well as in several other tumors. Weidner *et al.*, N. Engl. J. Med., **324**: 1-6 (1991); Horak *et al.*, Lancet, **340**: 1120-1124 (1992); Macchiarini *et al.*, Lancet, **340**: 145-146 (1992).

The search for positive regulators of angiogenesis has yielded many candidates, including aFGF, bFGF, TGF- $\alpha$ , TGF- $\beta$ , HGF, TNF- $\alpha$ , angiogenin, IL-8, etc. Folkman *et al.*, J.B.C., *supra*, and Klagsbrun *et al.*, *supra*. The negative regulators so far identified include thrombospondin (Good *et al.*, Proc. Natl. Acad. Sci. USA., **87**: 6624-6628 (1990)), the 16-kilodalton N-terminal fragment of prolactin (Clapp *et al.*, Endocrinology, **133**: 1292-1299 (1993)), angiostatin (O'Reilly *et al.*, Cell, **79**: 315-328 (1994)), and endostatin. O'Reilly *et al.*, Cell, **88**: 277-285 (1996).

Work done over the last several years has established the key role of VEGF, not only in stimulating vascular endothelial cell proliferation, but also in inducing vascular permeability and angiogenesis. Ferrara *et al.*, Endocr. Rev., **18**: 4-25 (1997). The finding that the loss of even a single VEGF allele results in embryonic lethality points to an irreplaceable role played by this factor in the development and differentiation of the vascular system. Furthermore, VEGF has been shown to be a key mediator of neovascularization associated with tumors and intraocular disorders. Ferrara *et al.*, Endocr. Rev., *supra*. The VEGF mRNA is overexpressed by the majority of human tumors examined. Berkman *et al.*, J. Clin. Invest., **91**: 153-159 (1993); Brown *et al.*, Human Pathol., **26**: 86-91 (1995); Brown *et al.*, Cancer Res., **53**: 4727-4735 (1993); Mattern *et al.*, Brit. J. Cancer, **73**: 931-934 (1996); Dvorak *et al.*, Am. J. Pathol., **146**: 1029-1039 (1995).

Also, the concentration levels of VEGF in eye fluids are highly correlated to the presence of active proliferation of blood vessels in patients with diabetic and other ischemia-related retinopathies. Aiello *et al.*, N. Engl. J. Med., **331**: 1480-1487 (1994). Furthermore, recent studies have demonstrated the localization of VEGF in choroidal neovascular membranes in patients affected by AMD. Lopez *et al.*, Invest. Ophthalmol. Vis. Sci., **37**: 855-868 (1996).

Anti-VEGF neutralizing antibodies suppress the growth of a variety of human tumor cell lines in nude mice (Kim *et al.*, Nature, **362**: 841-844 (1993); Warren *et al.*, J. Clin. Invest., **95**: 1789-1797 (1995); Borgström *et al.*, Cancer Res., **56**: 4032-4039 (1996); Melnyk *et al.*, Cancer Res., **56**: 921-924 (1996)) and also inhibit intraocular angiogenesis in models of ischemic retinal disorders. Adamis *et al.*, Arch. Ophthalmol., **114**: 66-71 (1996). Therefore, anti-VEGF monoclonal antibodies or other inhibitors of VEGF action are promising candidates for the treatment of solid tumors and various intraocular neovascular disorders. Such antibodies are described, for example, in EP 817,648 published January 14, 1998 and in PCT/US 98/06724 filed April 3, 1998.

There exist several other growth factors and mitogens, including transforming oncogenes, that are capable of rapidly inducing a complex set of genes to be expressed by certain cells. Lau and Nathans, Molecular Aspects of

Cellular Regulation, 6: 165-202 (1991). These genes, which have been named immediate-early- or early-response genes, are transcriptionally activated within minutes after contact with a growth factor or mitogen, independent of *de novo* protein synthesis. A group of these intermediate-early genes encodes secreted, extracellular proteins that are needed for coordination of complex biological processes such as differentiation and proliferation, regeneration, and wound healing. Ryseck *et al.*, Cell Growth Differ., 2: 235-233 (1991).

Highly-related proteins that belong to this group include *cef 10* (Simmons *et al.*, Proc. Natl. Acad. Sci. USA, 86: 1178-1182 (1989)), *cyr 61*, which is rapidly activated by serum- or platelet-derived growth factor (PDGF) (O'Brien *et al.*, Mol. Cell Biol., 10: 3569-3577 (1990), human connective tissue growth factor (CTGF) (Bradham *et al.*, J. Cell. Biol., 114: 1285-1294 (1991)), which is secreted by human vascular endothelial cells in high levels after activation with transforming growth factor beta (TGF- $\beta$ ), exhibits PDGF-like biological and immunological activities, and competes with PDGF for a particular cell surface receptor, *fisp-12* (Ryseck *et al.*, Cell Growth Differ., 2: 235-233 (1991)), human vascular IBP-like growth factor (VIGF) (WO 96/17931), and *nov*, normally arrested in adult kidney cells, which was found to be overexpressed in myeloblastosis-associated-virus-type-1-induced nephroblastomas. Jolait *et al.*, Mol. Cell. Biol., 12: 10-21 (1992).

The expression of these immediate-early genes acts as "third messengers" in the cascade of events triggered by growth factors. It is also thought that they are needed to integrate and coordinate complex biological processes, such as differentiation and wound healing in which cell proliferation is a common event.

As additional mitogens, insulin-like growth factor binding proteins (IGFBPs) have been shown, in complex with insulin-like growth factor (IGF), to stimulate increased binding of IGF to fibroblast and smooth muscle cell surface receptors. Clemmons *et al.*, J. Clin. Invest., 77: 1548 (1986). Inhibitory effects of IGFBP on various IGF actions *in vitro* include stimulation of glucose transport by adipocytes, sulfate incorporation by chondrocytes, and thymidine incorporation in fibroblast. Zapf *et al.*, J. Clin. Invest., 63: 1077 (1979). In addition, inhibitory effects of IGFBPs on growth factor-mediated mitogen activity in normal cells have been shown.

#### C. Need for Further Treatments

In view of the role of vascular endothelial cell growth and angiogenesis in many diseases and disorders, it is desirable to have a means of reducing or inhibiting one or more of the biological effects causing these processes. It is also desirable to have a means of assaying for the presence of pathogenic polypeptides in normal and diseased conditions, and especially cancer. Further, in a specific aspect, as there is no generally applicable therapy for the treatment of cardiac hypertrophy, the identification of factors that can prevent or reduce cardiac myocyte hypertrophy is of primary importance in the development of new therapeutic strategies to inhibit pathophysiological cardiac growth. While there are several treatment modalities for various cardiovascular and oncologic disorders, there is still a need for additional therapeutic approaches.

### Summary of the Invention

#### A. Embodiments

Accordingly, the present invention concerns compositions and methods for promoting or inhibiting angiogenesis and/or cardiovascularization in mammals. The present invention is based on the identification of

proteins that test positive in various cardiovascular assays that test promotion or inhibition of certain biological activities. Accordingly, the proteins are believed to be useful drugs for the diagnosis and/or treatment (including prevention) of disorders where such effects are desired, such as the promotion or inhibition of angiogenesis, inhibition or stimulation of vascular endothelial cell growth, stimulation of growth or proliferation of vascular endothelial cells, inhibition of tumor growth, inhibition of angiogenesis-dependent tissue growth, stimulation of angiogenesis-dependent tissue growth, inhibition of cardiac hypertrophy and stimulation of cardiac hypertrophy, *e.g.*, for the treatment of congestive heart failure.

In one embodiment, the present invention provides a composition comprising a PRO polypeptide in admixture with a pharmaceutically acceptable carrier. In one aspect, the composition comprises a therapeutically effective amount of the polypeptide. In another aspect, the composition comprises a further active ingredient, namely, a cardiovascular, endothelial or angiogenic agent or an angiostatic agent, preferably an angiogenic or angiostatic agent. Preferably, the composition is sterile. The PRO polypeptide may be administered in the form of a liquid pharmaceutical formulation, which may be preserved to achieve extended storage stability. Preserved liquid pharmaceutical formulations might contain multiple doses of PRO polypeptide, and might, therefore, be suitable for repeated use.

In a further embodiment, the present invention provides a method for preparing such a composition useful for the treatment of a cardiovascular, endothelial or angiogenic disorder comprising admixing a therapeutically effective amount of a PRO polypeptide with a pharmaceutically acceptable carrier.

In another embodiment, the present invention provides a composition comprising an agonist or antagonist of a PRO polypeptide in admixture with a pharmaceutically acceptable carrier. In one aspect, the composition comprises a therapeutically effective amount of the agonist or antagonist. In another aspect, the composition comprises a further active ingredient, namely, a cardiovascular, endothelial or angiogenic agent or an angiostatic agent, preferably an angiogenic or angiostatic agent. Preferably, the composition is sterile. The PRO polypeptide agonist or antagonist may be administered in the form of a liquid pharmaceutical formulation, which may be preserved to achieve extended storage stability. Preserved liquid pharmaceutical formulations might contain multiple doses of a PRO polypeptide agonist or antagonist, and might, therefore, be suitable for repeated use.

In a further embodiment, the present invention provides a method for preparing such a composition useful for the treatment of a cardiovascular, endothelial or angiogenic disorder comprising admixing a therapeutically effective amount of a PRO polypeptide agonist or antagonist with a pharmaceutically acceptable carrier.

In yet another embodiment, the present invention concerns a composition comprising an anti-PRO antibody in admixture with a pharmaceutically acceptable carrier. In one aspect, the composition comprises a therapeutically effective amount of the antibody. In another aspect, the composition comprises a further active ingredient, namely, a cardiovascular, endothelial or angiogenic agent or an angiostatic agent, preferably an angiogenic or angiostatic agent. Preferably, the composition is sterile. The composition may be administered in the form of a liquid pharmaceutical formulation, which may be preserved to achieve extended storage stability. Preserved liquid pharmaceutical formulations might contain multiple doses of the anti-PRO antibody, and might, therefore, be suitable for repeated use. In preferred embodiments, the antibody is a monoclonal antibody, an antibody fragment, a humanized antibody, or a single-chain antibody.

In a further embodiment, the present invention provides a method for preparing such a composition useful for the treatment of a cardiovascular, endothelial or angiogenic disorder comprising admixing a therapeutically effective amount of an anti-PRO antibody with a pharmaceutically acceptable carrier.

In a still further aspect, the present invention provides an article of manufacture comprising:

- 5 (a) a composition of matter comprising a PRO polypeptide or agonist or antagonist thereof;
- (b) a container containing said composition; and
- (c) a label affixed to said container, or a package insert included in said container referring to the use of said PRO polypeptide or agonist or antagonist thereof in the treatment of a cardiovascular, endothelial or angiogenic disorder, wherein the agonist or antagonist may be an antibody which binds to the PRO polypeptide. The composition may comprise a therapeutically effective amount of the PRO polypeptide or the agonist or antagonist thereof.
- 10

In another embodiment, the present invention provides a method for identifying an agonist of a PRO polypeptide comprising:

- 15 (a) contacting cells and a test compound to be screened under conditions suitable for the induction of a cellular response normally induced by a PRO polypeptide; and
- (b) determining the induction of said cellular response to determine if the test compound is an effective agonist, wherein the induction of said cellular response is indicative of said test compound being an effective agonist.

In another embodiment, the present invention provides a method for identifying an agonist of a PRO polypeptide comprising:

20

- (a) contacting cells and a test compound to be screened under conditions suitable for the stimulation of cell proliferation by a PRO polypeptide; and
- (b) measuring the proliferation of said cells to determine if the test compound is an effective agonist, wherein the stimulation of cell proliferation is indicative of said test compound being an effective agonist.

25 In another embodiment, the invention provides a method for identifying a compound that inhibits the activity of a PRO polypeptide comprising contacting a test compound with a PRO polypeptide under conditions and for a time sufficient to allow the test compound and polypeptide to interact and determining whether the activity of the PRO polypeptide is inhibited. In a specific preferred aspect, either the test compound or the PRO polypeptide is immobilized on a solid support. In another preferred aspect, the non-immobilized component carries a detectable label. In a preferred aspect, this method comprises the steps of:

30

- (a) contacting cells and a test compound to be screened in the presence of a PRO polypeptide under conditions suitable for the induction of a cellular response normally induced by a PRO polypeptide; and
- (b) determining the induction of said cellular response to determine if the test compound is an effective antagonist.

35 In another preferred aspect, this process comprises the steps of:

- (a) contacting cells and a test compound to be screened in the presence of a PRO polypeptide under conditions suitable for the stimulation of cell proliferation by a PRO polypeptide; and
- (b) measuring the proliferation of the cells to determine if the test compound is an effective antagonist.

In another embodiment, the invention provides a method for identifying a compound that inhibits the expression of a PRO polypeptide in cells that normally expresses the polypeptide, wherein the method comprises contacting the cells with a test compound and determining whether the expression of the PRO polypeptide is inhibited. In a preferred aspect, this method comprises the steps of:

- (a) contacting cells and a test compound to be screened under conditions suitable for allowing expression of the PRO polypeptide; and
- (b) determining the inhibition of expression of said polypeptide.

In a still further embodiment, the invention provides a compound that inhibits the expression of a PRO polypeptide, such as a compound that is identified by the methods set forth above.

Another aspect of the present invention is directed to an agonist or an antagonist of a PRO polypeptide which may optionally be identified by the methods described above.

One type of antagonist of a PRO polypeptide that inhibits one or more of the functions or activities of the PRO polypeptide is an antibody. Hence, in another aspect, the invention provides an isolated antibody that binds a PRO polypeptide. In a preferred aspect, the antibody is a monoclonal antibody, which preferably has non-human complementarity-determining-region (CDR) residues and human framework-region (FR) residues. The antibody may be labeled and may be immobilized on a solid support. In a further aspect, the antibody is an antibody fragment, a single-chain antibody, or a humanized antibody. Preferably, the antibody specifically binds to the polypeptide.

In a still further aspect, the present invention provides a method for diagnosing a disease or susceptibility to a disease which is related to a mutation in a PRO polypeptide-encoding nucleic acid sequence comprising determining the presence or absence of said mutation in the PRO polypeptide nucleic acid sequence, wherein the presence or absence of said mutation is indicative of the presence of said disease or susceptibility to said disease.

In a still further aspect, the invention provides a method of diagnosing a cardiovascular, endothelial or angiogenic disorder in a mammal which comprises analyzing the level of expression of a gene encoding a PRO polypeptide (a) in a test sample of tissue cells obtained from said mammal, and (b) in a control sample of known normal tissue cells of the same cell type, wherein a higher or lower expression level in the test sample as compared to the control sample is indicative of the presence of a cardiovascular, endothelial or angiogenic disorder in said mammal. The expression of a gene encoding a PRO polypeptide may optionally be accomplished by measuring the level of mRNA or the polypeptide in the test sample as compared to the control sample.

In a still further aspect, the present invention provides a method of diagnosing a cardiovascular, endothelial or angiogenic disorder in a mammal which comprises detecting the presence or absence of a PRO polypeptide in a test sample of tissue cells obtained from said mammal, wherein the presence or absence of said PRO polypeptide in said test sample is indicative of the presence of a cardiovascular, endothelial or angiogenic disorder in said mammal.

In a still further embodiment, the invention provides a method of diagnosing a cardiovascular, endothelial or angiogenic disorder in a mammal comprising (a) contacting an anti-PRO antibody with a test sample of tissue cells obtained from the mammal, and (b) detecting the formation of a complex between the antibody and the PRO polypeptide in the test sample, wherein the formation of said complex is indicative of the presence of a



cardiovascular, endothelial or angiogenic disorder in the mammal. The detection may be qualitative or quantitative, and may be performed in comparison with monitoring the complex formation in a control sample of known normal tissue cells of the same cell type. A larger or smaller quantity of complexes formed in the test sample indicates the presence of a cardiovascular, endothelial or angiogenic dysfunction in the mammal from which the test tissue cells were obtained. The antibody preferably carries a detectable label. Complex formation can be monitored, for example, by light microscopy, flow cytometry, fluorimetry, or other techniques known in the art. The test sample is usually obtained from an individual suspected to have a cardiovascular, endothelial or angiogenic disorder.

In another embodiment, the invention provides a method for determining the presence of a PRO polypeptide in a sample comprising exposing a sample suspected of containing the PRO polypeptide to an anti-PRO antibody and determining binding of said antibody to a component of said sample. In a specific aspect, the sample comprises a cell suspected of containing the PRO polypeptide and the antibody binds to the cell. The antibody is preferably detectably labeled and/or bound to a solid support.

In further aspects, the invention provides a cardiovascular, endothelial or angiogenic disorder diagnostic kit comprising an anti-PRO antibody and a carrier in suitable packaging. Preferably, such kit further comprises instructions for using said antibody to detect the presence of the PRO polypeptide. Preferably, the carrier is a buffer, for example. Preferably, the cardiovascular, endothelial or angiogenic disorder is cancer.

In yet another embodiment, the present invention provides a method for treating a cardiovascular, endothelial or angiogenic disorder in a mammal comprising administering to the mammal an effective amount of a PRO polypeptide. Preferably, the disorder is cardiac hypertrophy, trauma such as wounds or burns, or a type of cancer. In a further aspect, the mammal is further exposed to angioplasty or a drug that treats cardiovascular, endothelial or angiogenic disorders such as ACE inhibitors or chemotherapeutic agents if the cardiovascular, endothelial or angiogenic disorder is a type of cancer. Preferably, the mammal is human, preferably one who is at risk of developing cardiac hypertrophy and more preferably has suffered myocardial infarction.

In another preferred aspect, the cardiac hypertrophy is characterized by the presence of an elevated level of  $\text{PGF}_{2\alpha}$ . Alternatively, the cardiac hypertrophy may be induced by myocardial infarction, wherein preferably the administration of the PRO polypeptide is initiated within 48 hours, more preferably within 24 hours, following myocardial infarction.

In another preferred embodiment, the cardiovascular, endothelial or angiogenic disorder is cardiac hypertrophy and said PRO polypeptide is administered together with a cardiovascular, endothelial or angiogenic agent. The preferred cardiovascular, endothelial or angiogenic agent for this purpose is selected from the group consisting of an antihypertensive drug, an ACE inhibitor, an endothelin receptor antagonist and a thrombolytic agent. If a thrombolytic agent is administered, preferably the PRO polypeptide is administered following administration of such agent. More preferably, the thrombolytic agent is recombinant human tissue plasminogen activator.

In another preferred aspect, the cardiovascular, endothelial or angiogenic disorder is cardiac hypertrophy and the PRO polypeptide is administered following primary angioplasty for the treatment of acute myocardial infarction, preferably wherein the mammal is further exposed to angioplasty or a cardiovascular, endothelial, or angiogenic agent.

In another preferred embodiment, the cardiovascular, endothelial or angiogenic disorder is a cancer and the

PRO polypeptide is administered in combination with a chemotherapeutic agent, a growth inhibitory agent or a cytotoxic agent.

In a further embodiment, the invention concerns a method for treating a cardiovascular, endothelial or angiogenic disorder in a mammal comprising administering to the mammal an effective amount of an agonist of a PRO polypeptide. Preferably, the cardiovascular, endothelial or angiogenic disorder is cardiac hypertrophy, trauma, a cancer, or age-related macular degeneration. Also preferred is where the mammal is human, and where an effective amount of an angiogenic or angiostatic agent is administered in conjunction with the agonist.

In a further embodiment, the invention concerns a method for treating a cardiovascular, endothelial or angiogenic disorder in a mammal comprising administering to the mammal an effective amount of an antagonist of a PRO polypeptide. Preferably, the cardiovascular, endothelial or angiogenic disorder is cardiac hypertrophy, trauma, a cancer, or age-related macular degeneration. Also preferred is where the mammal is human, and where an effective amount of an angiogenic or angiostatic agent is administered in conjunction with the antagonist.

In a further embodiment, the invention concerns a method for treating a cardiovascular, endothelial or angiogenic disorder in a mammal comprising administering to the mammal an effective amount of an anti-PRO antibody. Preferably, the cardiovascular, endothelial or angiogenic disorder is cardiac hypertrophy, trauma, a cancer, or age-related macular degeneration. Also preferred is where the mammal is human, and where an effective amount of an angiogenic or angiostatic agent is administered in conjunction with the antibody.

In still further embodiments, the invention provides a method for treating a cardiovascular, endothelial or angiogenic disorder in a mammal that suffers therefrom comprising administering to the mammal a nucleic acid molecule that codes for either (a) a PRO polypeptide, (b) an agonist of a PRO polypeptide or (c) an antagonist of a PRO polypeptide, wherein said agonist or antagonist may be an anti-PRO antibody. In a preferred embodiment, the mammal is human. In another preferred embodiment, the gene is administered via *ex vivo* gene therapy. In a further preferred embodiment, the gene is comprised within a vector, more preferably an adenoviral, adeno-associated viral, lentiviral, or retroviral vector.

In yet another aspect, the invention provides a recombinant retroviral particle comprising a retroviral vector consisting essentially of a promoter, nucleic acid encoding (a) a PRO polypeptide, (b) an agonist polypeptide of a PRO polypeptide, or (c) an antagonist polypeptide of a PRO polypeptide, and a signal sequence for cellular secretion of the polypeptide, wherein the retroviral vector is in association with retroviral structural proteins. Preferably, the signal sequence is from a mammal, such as from a native PRO polypeptide.

In a still further embodiment, the invention supplies an *ex vivo* producer cell comprising a nucleic acid construct that expresses retroviral structural proteins and also comprises a retroviral vector consisting essentially of a promoter, nucleic acid encoding (a) a PRO polypeptide, (b) an agonist polypeptide of a PRO polypeptide or (c) an antagonist polypeptide of a PRO polypeptide, and a signal sequence for cellular secretion of the polypeptide, wherein said producer cell packages the retroviral vector in association with the structural proteins to produce recombinant retroviral particles.

In yet another embodiment, the invention provides a method for inhibiting endothelial cell growth in a mammal comprising administering to the mammal (a) a PRO polypeptide, (b) an agonist of a PRO polypeptide, or (c) an antagonist of a PRO polypeptide, wherein endothelial cell growth in said mammal is inhibited, and wherein said

agonist or antagonist may be an anti-PRO antibody. Preferably, the mammal is human and the endothelial cell growth is associated with a tumor or a retinal disorder.

In yet another embodiment, the invention provides a method for stimulating endothelial cell growth in a mammal comprising administering to the mammal (a) a PRO polypeptide, (b) an agonist of a PRO polypeptide, or (c) an antagonist of a PRO polypeptide, wherein endothelial cell growth in said mammal is stimulated, and wherein said agonist or antagonist may be an anti-PRO antibody. Preferably, the mammal is human.

In yet another embodiment, the invention provides a method for inhibiting cardiac hypertrophy in a mammal comprising administering to the mammal (a) a PRO polypeptide, (b) an agonist of a PRO polypeptide, or (c) an antagonist of a PRO polypeptide, wherein cardiac hypertrophy in said mammal is inhibited, and wherein said agonist or antagonist may be an anti-PRO antibody. Preferably, the mammal is human and the cardiac hypertrophy has been induced by myocardial infarction.

In yet another embodiment, the invention provides a method for stimulating cardiac hypertrophy in a mammal comprising administering to the mammal (a) a PRO polypeptide, (b) an agonist of a PRO polypeptide, or (c) an antagonist of a PRO polypeptide, wherein cardiac hypertrophy in said mammal is stimulated, and wherein said agonist or antagonist may be an anti-PRO antibody. Preferably, the mammal is human who suffers from congestive heart failure.

In yet another embodiment, the invention provides a method for inhibiting angiogenesis induced by a PRO polypeptide in a mammal comprising administering a therapeutically effective amount of an anti-PRO antibody to the mammal. Preferably, the mammal is a human, and more preferably the mammal has a tumor or a retinal disorder.

In yet another embodiment, the invention provides a method for stimulating angiogenesis induced by a PRO polypeptide in a mammal comprising administering a therapeutically effective amount of a PRO polypeptide to the mammal. Preferably, the mammal is a human, and more preferably angiogenesis would promote tissue regeneration or wound healing.

#### B. Additional Embodiments

In other embodiments of the present invention, the invention provides an isolated nucleic acid molecule comprising a nucleotide sequence that encodes a PRO polypeptide.

In one aspect, the isolated nucleic acid molecule comprises a nucleotide sequence having at least about 80% sequence identity, preferably at least about 81% sequence identity, more preferably at least about 82% sequence identity, yet more preferably at least about 83% sequence identity, yet more preferably at least about 84% sequence identity, yet more preferably at least about 85% sequence identity, yet more preferably at least about 86% sequence identity, yet more preferably at least about 87% sequence identity, yet more preferably at least about 88% sequence identity, yet more preferably at least about 89% sequence identity, yet more preferably at least about 90% sequence identity, yet more preferably at least about 91% sequence identity, yet more preferably at least about 92% sequence identity, yet more preferably at least about 93% sequence identity, yet more preferably at least about 94% sequence identity, yet more preferably at least about 95% sequence identity, yet more preferably at least about 96% sequence identity, yet more preferably at least about 97% sequence identity, yet more preferably at least about 98% sequence

identity and yet more preferably at least about 99% sequence identity to (a) a DNA molecule encoding a PRO polypeptide having a full-length amino acid sequence as disclosed herein, an amino acid sequence lacking the signal peptide as disclosed herein, an extracellular domain of a transmembrane protein, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of the full-length amino acid sequence as disclosed herein, or (b) the complement of the DNA molecule of (a).

In other aspects, the isolated nucleic acid molecule comprises a nucleotide sequence having at least about 80% sequence identity, preferably at least about 81% sequence identity, more preferably at least about 82% sequence identity, yet more preferably at least about 83% sequence identity, yet more preferably at least about 84% sequence identity, yet more preferably at least about 85% sequence identity, yet more preferably at least about 86% sequence identity, yet more preferably at least about 87% sequence identity, yet more preferably at least about 88% sequence identity, yet more preferably at least about 89% sequence identity, yet more preferably at least about 90% sequence identity, yet more preferably at least about 91% sequence identity, yet more preferably at least about 92% sequence identity, yet more preferably at least about 93% sequence identity, yet more preferably at least about 94% sequence identity, yet more preferably at least about 95% sequence identity, yet more preferably at least about 96% sequence identity, yet more preferably at least about 97% sequence identity, yet more preferably at least about 98% sequence identity and yet more preferably at least about 99% sequence identity to (a) a DNA molecule comprising the coding sequence of a full-length PRO polypeptide cDNA as disclosed herein, the coding sequence of a PRO polypeptide lacking the signal peptide as disclosed herein, the coding sequence of an extracellular domain of a transmembrane PRO polypeptide, with or without the signal peptide, as disclosed herein or the coding sequence of any other specifically defined fragment of the full-length amino acid sequence as disclosed herein, or (b) the complement of the DNA molecule of (a).

In a further aspect, the invention concerns an isolated nucleic acid molecule comprising a nucleotide sequence having at least about 80% sequence identity, preferably at least about 81% sequence identity, more preferably at least about 82% sequence identity, yet more preferably at least about 83% sequence identity, yet more preferably at least about 84% sequence identity, yet more preferably at least about 85% sequence identity, yet more preferably at least about 86% sequence identity, yet more preferably at least about 87% sequence identity, yet more preferably at least about 88% sequence identity, yet more preferably at least about 89% sequence identity, yet more preferably at least about 90% sequence identity, yet more preferably at least about 91% sequence identity, yet more preferably at least about 92% sequence identity, yet more preferably at least about 93% sequence identity, yet more preferably at least about 94% sequence identity, yet more preferably at least about 95% sequence identity, yet more preferably at least about 96% sequence identity, yet more preferably at least about 97% sequence identity, yet more preferably at least about 98% sequence identity and yet more preferably at least about 99% sequence identity to (a) a DNA molecule that encodes the same mature polypeptide encoded by any of the human protein cDNAs deposited with the ATCC as disclosed herein, or (b) the complement of the DNA molecule of (a).

Another aspect of the invention provides an isolated nucleic acid molecule comprising a nucleotide sequence encoding a PRO polypeptide which is either transmembrane domain-deleted or transmembrane domain-inactivated, or is complementary to such encoding nucleotide sequence, wherein the transmembrane domain(s) of such polypeptide are disclosed herein. Therefore, soluble extracellular domains of the herein described PRO

polypeptides are contemplated.

Another embodiment is directed to fragments of a PRO polypeptide coding sequence, or the complement thereof, that may find use as, for example, hybridization probes, for encoding fragments of a PRO polypeptide that may optionally encode a polypeptide comprising a binding site for an anti-PRO antibody or as antisense oligonucleotide probes. Such nucleic acid fragments are usually at least about 20 nucleotides in length, preferably at least about 30 nucleotides in length, more preferably at least about 40 nucleotides in length, yet more preferably at least about 50 nucleotides in length, yet more preferably at least about 60 nucleotides in length, yet more preferably at least about 70 nucleotides in length, yet more preferably at least about 80 nucleotides in length, yet more preferably at least about 90 nucleotides in length, yet more preferably at least about 100 nucleotides in length, yet more preferably at least about 110 nucleotides in length, yet more preferably at least about 120 nucleotides in length, yet more preferably at least about 130 nucleotides in length, yet more preferably at least about 140 nucleotides in length, yet more preferably at least about 150 nucleotides in length, yet more preferably at least about 160 nucleotides in length, yet more preferably at least about 170 nucleotides in length, yet more preferably at least about 180 nucleotides in length, yet more preferably at least about 190 nucleotides in length, yet more preferably at least about 200 nucleotides in length, yet more preferably at least about 250 nucleotides in length, yet more preferably at least about 300 nucleotides in length, yet more preferably at least about 350 nucleotides in length, yet more preferably at least about 400 nucleotides in length, yet more preferably at least about 450 nucleotides in length, yet more preferably at least about 500 nucleotides in length, yet more preferably at least about 600 nucleotides in length, yet more preferably at least about 700 nucleotides in length, yet more preferably at least about 800 nucleotides in length, yet more preferably at least about 900 nucleotides in length and yet more preferably at least about 1000 nucleotides in length, wherein in this context the term "about" means the referenced nucleotide sequence length plus or minus 10% of that referenced length. It is noted that novel fragments of a PRO polypeptide-encoding nucleotide sequence may be determined in a routine manner by aligning the PRO polypeptide-encoding nucleotide sequence with other known nucleotide sequences using any of a number of well known sequence alignment programs and determining which PRO polypeptide-encoding nucleotide sequence fragment(s) are novel. All of such PRO polypeptide-encoding nucleotide sequences are contemplated herein. Also contemplated are the PRO polypeptide fragments encoded by these nucleotide molecule fragments, preferably those PRO polypeptide fragments that comprise a binding site for an anti-PRO antibody.

In another embodiment, the invention provides isolated PRO polypeptide encoded by any of the isolated nucleic acid sequences hereinabove identified.

In a certain aspect, the invention concerns an isolated PRO polypeptide, comprising an amino acid sequence having at least about 80% sequence identity, preferably at least about 81% sequence identity, more preferably at least about 82% sequence identity, yet more preferably at least about 83% sequence identity, yet more preferably at least about 84% sequence identity, yet more preferably at least about 85% sequence identity, yet more preferably at least about 86% sequence identity, yet more preferably at least about 87% sequence identity, yet more preferably at least about 88% sequence identity, yet more preferably at least about 89% sequence identity, yet more preferably at least about 90% sequence identity, yet more preferably at least about 91% sequence identity, yet more preferably at least about 92% sequence identity, yet more preferably at least about 93% sequence identity, yet more preferably

at least about 94% sequence identity, yet more preferably at least about 95% sequence identity, yet more preferably at least about 96% sequence identity, yet more preferably at least about 97% sequence identity, yet more preferably at least about 98% sequence identity and yet more preferably at least about 99% sequence identity to a PRO polypeptide having a full-length amino acid sequence as disclosed herein, an amino acid sequence lacking the signal peptide as disclosed herein, an extracellular domain of a transmembrane protein, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of the full-length amino acid sequence as disclosed herein.

In a further aspect, the invention concerns an isolated PRO polypeptide comprising an amino acid sequence having at least about 80% sequence identity, preferably at least about 81% sequence identity, more preferably at least about 82% sequence identity, yet more preferably at least about 83% sequence identity, yet more preferably at least about 84% sequence identity, yet more preferably at least about 85% sequence identity, yet more preferably at least about 86% sequence identity, yet more preferably at least about 87% sequence identity, yet more preferably at least about 88% sequence identity, yet more preferably at least about 89% sequence identity, yet more preferably at least about 90% sequence identity, yet more preferably at least about 91% sequence identity, yet more preferably at least about 92% sequence identity, yet more preferably at least about 93% sequence identity, yet more preferably at least about 94% sequence identity, yet more preferably at least about 95% sequence identity, yet more preferably at least about 96% sequence identity, yet more preferably at least about 97% sequence identity, yet more preferably at least about 98% sequence identity and yet more preferably at least about 99% sequence identity to an amino acid sequence encoded by any of the human protein cDNAs deposited with the ATCC as disclosed herein.

In a further aspect, the invention concerns an isolated PRO polypeptide comprising an amino acid sequence scoring at least about 80% positives, preferably at least about 81% positives, more preferably at least about 82% positives, yet more preferably at least about 83% positives, yet more preferably at least about 84% positives, yet more preferably at least about 85% positives, yet more preferably at least about 86% positives, yet more preferably at least about 87% positives, yet more preferably at least about 88% positives, yet more preferably at least about 89% positives, yet more preferably at least about 90% positives, yet more preferably at least about 91% positives, yet more preferably at least about 92% positives, yet more preferably at least about 93% positives, yet more preferably at least about 94% positives, yet more preferably at least about 95% positives, yet more preferably at least about 96% positives, yet more preferably at least about 97% positives, yet more preferably at least about 98% positives and yet more preferably at least about 99% positives when compared with the amino acid sequence of a PRO polypeptide having a full-length amino acid sequence as disclosed herein, an amino acid sequence lacking the signal peptide as disclosed herein, an extracellular domain of a transmembrane protein, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of the full-length amino acid sequence as disclosed herein.

In a specific aspect, the invention provides an isolated PRO polypeptide without the N-terminal signal sequence and/or the initiating methionine and is encoded by a nucleotide sequence that encodes such an amino acid sequence as hereinbefore described. Processes for producing the same are also herein described, wherein those processes comprise culturing a host cell comprising a vector which comprises the appropriate encoding nucleic acid molecule under conditions suitable for expression of the PRO polypeptide and recovering the PRO polypeptide from

the cell culture.

Another aspect of the invention provides an isolated PRO polypeptide which is either transmembrane domain-deleted or transmembrane domain-inactivated. Processes for producing the same are also herein described, wherein those processes comprise culturing a host cell comprising a vector which comprises the appropriate encoding nucleic acid molecule under conditions suitable for expression of the PRO polypeptide and recovering the PRO polypeptide from the cell culture.

In yet another embodiment, the invention concerns agonists and antagonists of a native PRO polypeptide as defined herein. In a particular embodiment, the agonist or antagonist is an anti-PRO antibody or a small molecule.

In a further embodiment, the invention concerns a method of identifying agonists or antagonists to a PRO polypeptide which comprise contacting the PRO polypeptide with a candidate molecule and monitoring a biological activity mediated by said PRO polypeptide. Preferably, the PRO polypeptide is a native PRO polypeptide.

In a still further embodiment, the invention concerns a composition of matter comprising a PRO polypeptide, or an agonist or antagonist of a PRO polypeptide as herein described, or an anti-PRO antibody, in combination with a carrier. Optionally, the carrier is a pharmaceutically acceptable carrier.

Another embodiment of the present invention is directed to the use of a PRO polypeptide, or an agonist or antagonist thereof as hereinbefore described, or an anti-PRO antibody, for the preparation of a medicament useful in the treatment of a condition which is responsive to the PRO polypeptide, an agonist or antagonist thereof or an anti-PRO antibody.

In additional embodiments of the present invention, the invention provides vectors comprising DNA encoding any of the herein described polypeptides. Host cell comprising any such vector are also provided. By way of example, the host cells may be CHO cells, *E. coli*, yeast, or Baculovirus-infected insect cells. A process for producing any of the herein described polypeptides is further provided and comprises culturing host cells under conditions suitable for expression of the desired polypeptide and recovering the desired polypeptide from the cell culture.

In other embodiments, the invention provides chimeric molecules comprising any of the herein described polypeptides fused to a heterologous polypeptide or amino acid sequence. Example of such chimeric molecules comprise any of the herein described polypeptides fused to an epitope tag sequence or a Fc region of an immunoglobulin.

In yet another embodiment, the invention provides an antibody which specifically binds to any of the above or below described polypeptides. Optionally, the antibody is a monoclonal antibody, humanized antibody, antibody fragment or single-chain antibody.

In yet other embodiments, the invention provides oligonucleotide probes useful for isolating genomic and cDNA nucleotide sequences or as antisense probes, wherein those probes may be derived from any of the above or below described nucleotide sequences.

#### Brief Description of the Drawings

Figures 1A through 1B show a nucleotide sequence (SEQ ID NO:1) of a native sequence PRO172 cDNA, wherein SEQ ID NO:1 is a clone designated herein as "DNA35916-1161".

Figure 2 shows the amino acid sequence (SEQ ID NO:2) derived from the coding sequence of SEQ ID NO:1 shown in Figures 1A through 1B.

Figures 3A through 3B show a nucleotide sequence (SEQ ID NO:8) of a native sequence PRO175 cDNA, wherein SEQ ID NO:8 is a clone designated herein as "DNA19355-1150".

5 Figure 4 shows the amino acid sequence (SEQ ID NO:9) derived from the coding sequence of SEQ ID NO:8 shown in Figures 3A through 3B.

Figure 5 shows a nucleotide sequence (SEQ ID NO:10) of a native sequence PRO178 cDNA, wherein SEQ ID NO:10 is a clone designated herein as "DNA23339-1130".

10 Figure 6 shows the amino acid sequence (SEQ ID NO:11) derived from the coding sequence of SEQ ID NO:10 shown in Figure 5.

Figures 7A through 7B show a nucleotide sequence (SEQ ID NO:15) of a native sequence PRO188 cDNA, wherein SEQ ID NO:15 is a clone designated herein as "DNA28497-1130".

Figure 8 shows the amino acid sequence (SEQ ID NO:16) derived from the coding sequence of SEQ ID NO:15 shown in Figures 7A through 7B.

15 Figures 9A through 9B show a nucleotide sequence (SEQ ID NO:20) of a native sequence PRO356 cDNA, wherein SEQ ID NO:20 is a clone designated herein as "DNA47470-1130-P1".

Figure 10 shows the amino acid sequence (SEQ ID NO:21) derived from the coding sequence of SEQ ID NO:20 shown in Figures 9A through 9B.

20 Figures 11A through 11B show a nucleotide sequence (SEQ ID NO:25) of a native sequence PRO179 cDNA, wherein SEQ ID NO:25 is a clone designated herein as "DNA16451-1078".

Figure 12 shows the amino acid sequence (SEQ ID NO:26) derived from the coding sequence of SEQ ID NO:25 shown in Figures 11A through 11B.

Figure 13 shows a nucleotide sequence (SEQ ID NO:30) of a native sequence PRO197 cDNA, wherein SEQ ID NO:30 is a clone designated herein as "DNA22780-1078".

25 Figure 14 shows the amino acid sequence (SEQ ID NO:31) derived from the coding sequence of SEQ ID NO:30 shown in Figure 13.

Figure 15 shows a nucleotide sequence (SEQ ID NO:35) of a native sequence PRO198 cDNA, wherein SEQ ID NO:35 is a clone designated herein as "DNA33457-1078".

30 Figure 16 shows the amino acid sequence (SEQ ID NO:36) derived from the coding sequence of SEQ ID NO:35 shown in Figure 15.

Figure 17 shows a nucleotide sequence (SEQ ID NO:40) of a native sequence PRO182 cDNA, wherein SEQ ID NO:40 is a clone designated herein as "DNA27865-1091".

Figure 18 shows the amino acid sequence (SEQ ID NO:41) derived from the coding sequence of SEQ ID NO:40 shown in Figure 17.

35 Figure 19 shows a nucleotide sequence (SEQ ID NO:45) of a native sequence PRO195 cDNA, wherein SEQ ID NO:45 is a clone designated herein as "DNA26847-1395".

Figure 20 shows the amino acid sequence (SEQ ID NO:46) derived from the coding sequence of SEQ ID NO:45 shown in Figure 19.



Figures 21A through 21B show a nucleotide sequence (SEQ ID NO:50) of a native sequence PRO200 cDNA, wherein SEQ ID NO:50 is a clone designated herein as "DNA29101-1122".

Figure 22 shows the amino acid sequence (SEQ ID NO:51) derived from the coding sequence of SEQ ID NO:50 shown in Figures 21A through 21B.

5 Figure 23 shows a nucleotide sequence (SEQ ID NO:56) of a native sequence PRO211 cDNA, wherein SEQ ID NO:56 is a clone designated herein as "DNA32292-1131".

Figure 24 shows the amino acid sequence (SEQ ID NO:57) derived from the coding sequence of SEQ ID NO:56 shown in Figure 23.

10 Figures 25A through 25B show a nucleotide sequence (SEQ ID NO:61) of a native sequence PRO217 cDNA, wherein SEQ ID NO:61 is a clone designated herein as "DNA33094-1131".

Figure 26 shows the amino acid sequence (SEQ ID NO:62) derived from the coding sequence of SEQ ID NO:61 shown in Figures 25A through 25B.

Figures 27A through 27B show a nucleotide sequence (SEQ ID NO:66) of a native sequence PRO219 cDNA, wherein SEQ ID NO:66 is a clone designated herein as "DNA32290-1164".

15 Figure 28 shows the amino acid sequence (SEQ ID NO:67) derived from the coding sequence of SEQ ID NO:66 shown in Figures 27A through 27B.

Figure 29 shows a nucleotide sequence (SEQ ID NO:71) of a native sequence PRO221 cDNA, wherein SEQ ID NO:71 is a clone designated herein as "DNA33089-1132".

20 Figure 30 shows the amino acid sequence (SEQ ID NO:72) derived from the coding sequence of SEQ ID NO:71 shown in Figure 29.

Figure 31 shows a nucleotide sequence (SEQ ID NO:76) of a native sequence PRO224 cDNA, wherein SEQ ID NO:76 is a clone designated herein as "DNA33221-1133".

Figure 32 shows the amino acid sequence (SEQ ID NO:77) derived from the coding sequence of SEQ ID NO:76 shown in Figure 31.

25 Figures 33A through 33B show a nucleotide sequence (SEQ ID NO:81) of a native sequence PRO228 cDNA, wherein SEQ ID NO:81 is a clone designated herein as "DNA33092-1202".

Figure 34 shows the amino acid sequence (SEQ ID NO:82) derived from the coding sequence of SEQ ID NO:81 shown in Figures 33A through 33B.

30 Figure 35 shows a nucleotide sequence (SEQ ID NO:90) of a native sequence PRO245 cDNA, wherein SEQ ID NO:90 is a clone designated herein as "DNA35638-1141".

Figure 36 shows the amino acid sequence (SEQ ID NO:91) derived from the coding sequence of SEQ ID NO:90 shown in Figure 35.

Figure 37 shows a nucleotide sequence (SEQ ID NO:95) of a native sequence PRO246 cDNA, wherein SEQ ID NO:95 is a clone designated herein as "DNA35639-1172".

35 Figure 38 shows the amino acid sequence (SEQ ID NO:96) derived from the coding sequence of SEQ ID NO:95 shown in Figure 37.

Figure 39 shows a nucleotide sequence (SEQ ID NO:100) of a native sequence PRO258 cDNA, wherein SEQ ID NO:100 is a clone designated herein as "DNA35918-1174".

Figure 40 shows the amino acid sequence (SEQ ID NO:101) derived from the coding sequence of SEQ ID NO:100 shown in Figure 39.

Figure 41 shows a nucleotide sequence (SEQ ID NO:107) of a native sequence PRO261 cDNA, wherein SEQ ID NO:107 is a clone designated herein as "DNA33473-1176".

5 Figure 42 shows the amino acid sequence (SEQ ID NO:108) derived from the coding sequence of SEQ ID NO:107 shown in Figure 41.

Figure 43 shows a nucleotide sequence (SEQ ID NO:112) of a native sequence PRO272 cDNA, wherein SEQ ID NO:112 is a clone designated herein as "DNA40620-1183".

10 Figure 44 shows the amino acid sequence (SEQ ID NO:113) derived from the coding sequence of SEQ ID NO:112 shown in Figure 43.

Figure 45 shows a nucleotide sequence (SEQ ID NO:118) of a native sequence PRO301 cDNA, wherein SEQ ID NO:118 is a clone designated herein as "DNA40628-1216".

Figure 46 shows the amino acid sequence (SEQ ID NO:119) derived from the coding sequence of SEQ ID NO:118 shown in Figure 45.

15 Figure 47 shows a nucleotide sequence (SEQ ID NO:126) of a native sequence PRO322 cDNA, wherein SEQ ID NO:126 is a clone designated herein as "DNA48336-1309".

Figure 48 shows the amino acid sequence (SEQ ID NO:127) derived from the coding sequence of SEQ ID NO:126 shown in Figure 47.

20 Figure 49 shows a nucleotide sequence (SEQ ID NO:131) of a native sequence PRO328 cDNA, wherein SEQ ID NO:131 is a clone designated herein as "DNA40587-1231".

Figure 50 shows the amino acid sequence (SEQ ID NO:132) derived from the coding sequence of SEQ ID NO:131 shown in Figure 49.

Figures 51A through 51B show a nucleotide sequence (SEQ ID NO:136) of a native sequence PRO331 cDNA, wherein SEQ ID NO:136 is a clone designated herein as "DNA40981-1234".

25 Figure 52 shows the amino acid sequence (SEQ ID NO:137) derived from the coding sequence of SEQ ID NO:136 shown in Figures 51A through 51B.

Figure 53 shows a nucleotide sequence (SEQ ID NO:141) of a native sequence PRO364 cDNA, wherein SEQ ID NO:141 is a clone designated herein as "DNA47365-1206".

30 Figure 54 shows the amino acid sequence (SEQ ID NO:142) derived from the coding sequence of SEQ ID NO:141 shown in Figure 53.

Figure 55 shows a nucleotide sequence (SEQ ID NO:151) of a native sequence PRO366 cDNA, wherein SEQ ID NO:151 is a clone designated herein as "DNA33085-1110".

Figure 56 shows the amino acid sequence (SEQ ID NO:152) derived from the coding sequence of SEQ ID NO:151 shown in Figure 55.

35 Figure 57 shows a nucleotide sequence (SEQ ID NO:153) of a native sequence PRO535 cDNA, wherein SEQ ID NO:153 is a clone designated herein as "DNA49143-1429".

Figure 58 shows the amino acid sequence (SEQ ID NO:154) derived from the coding sequence of SEQ ID NO:153 shown in Figure 57.

Figure 59 shows a nucleotide sequence (SEQ ID NO:155) of a native sequence PRO819 cDNA, wherein SEQ ID NO:155 is a clone designated herein as "DNA57695-1340".

Figure 60 shows the amino acid sequence (SEQ ID NO:156) derived from the coding sequence of SEQ ID NO:155 shown in Figure 59.

5 Figure 61 shows a nucleotide sequence (SEQ ID NO:157) of a native sequence PRO826 cDNA, wherein SEQ ID NO:157 is a clone designated herein as "DNA57694-1341".

Figure 62 shows the amino acid sequence (SEQ ID NO:158) derived from the coding sequence of SEQ ID NO:157 shown in Figure 61.

10 Figure 63 shows a nucleotide sequence (SEQ ID NO:159) of a native sequence PRO1160 cDNA, wherein SEQ ID NO:159 is a clone designated herein as "DNA62872-1509".

Figure 64 shows the amino acid sequence (SEQ ID NO:160) derived from the coding sequence of SEQ ID NO:159 shown in Figure 63.

Figure 65 shows a nucleotide sequence (SEQ ID NO:164) of a native sequence PRO1186 cDNA, wherein SEQ ID NO:164 is a clone designated herein as "DNA60621-1516".

15 Figure 66 shows the amino acid sequence (SEQ ID NO:165) derived from the coding sequence of SEQ ID NO:164 shown in Figure 65.

Figures 67A through 67B show a nucleotide sequence (SEQ ID NO:166) of a native sequence PRO1246 cDNA, wherein SEQ ID NO:166 is a clone designated herein as "DNA64885-1529".

20 Figure 68 shows the amino acid sequence (SEQ ID NO:167) derived from the coding sequence of SEQ ID NO:166 shown in Figures 67A through 67B.

### Detailed Description of the Invention

#### I. Definitions

25 The phrases "cardiovascular, endothelial and angiogenic disorder", "cardiovascular, endothelial and angiogenic dysfunction", "cardiovascular, endothelial or angiogenic disorder" and "cardiovascular, endothelial or angiogenic dysfunction" are used interchangeably and refer in part to systemic disorders that affect vessels, such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins, and/or lymphatics. This would include indications that stimulate angiogenesis and/or cardiovascularization, and those that inhibit angiogenesis and/or cardiovascularization. Such disorders include, for example, arterial disease, such as atherosclerosis, hypertension, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, and arterial restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, cancer such as vascular tumors, *e.g.*, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, Kaposi's sarcoma, lymphangioma, and lymphangiosarcoma, tumor angiogenesis, trauma such as wounds, burns, and other injured tissue, implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. This would also include angina, myocardial infarctions such as acute myocardial infarctions, cardiac hypertrophy, and heart failure such as CHF.

35

“Hypertrophy”, as used herein, is defined as an increase in mass of an organ or structure independent of natural growth that does not involve tumor formation. Hypertrophy of an organ or tissue is due either to an increase in the mass of the individual cells (true hypertrophy), or to an increase in the number of cells making up the tissue (hyperplasia), or both. Certain organs, such as the heart, lose the ability to divide shortly after birth. Accordingly, “cardiac hypertrophy” is defined as an increase in mass of the heart, which, in adults, is characterized by an increase in myocyte cell size and contractile protein content without concomitant cell division. The character of the stress responsible for inciting the hypertrophy, (*e.g.*, increased preload, increased afterload, loss of myocytes, as in myocardial infarction, or primary depression of contractility), appears to play a critical role in determining the nature of the response. The early stage of cardiac hypertrophy is usually characterized morphologically by increases in the size of myofibrils and mitochondria, as well as by enlargement of mitochondria and nuclei. At this stage, while muscle cells are larger than normal, cellular organization is largely preserved. At a more advanced stage of cardiac hypertrophy, there are preferential increases in the size or number of specific organelles, such as mitochondria, and new contractile elements are added in localized areas of the cells, in an irregular manner. Cells subjected to long-standing hypertrophy show more obvious disruptions in cellular organization, including markedly enlarged nuclei with highly lobulated membranes, which displace adjacent myofibrils and cause breakdown of normal Z-band registration. The phrase “cardiac hypertrophy” is used to include all stages of the progression of this condition, characterized by various degrees of structural damage of the heart muscle, regardless of the underlying cardiac disorder. Hence, the term also includes physiological conditions instrumental in the development of cardiac hypertrophy, such as elevated blood pressure, aortic stenosis, or myocardial infarction.

“Heart failure” refers to an abnormality of cardiac function where the heart does not pump blood at the rate needed for the requirements of metabolizing tissues. The heart failure can be caused by a number of factors, including ischemic, congenital, rheumatic, or idiopathic forms.

“Congestive heart failure” (CHF) is a progressive pathologic state where the heart is increasingly unable to supply adequate cardiac output (the volume of blood pumped by the heart over time) to deliver the oxygenated blood to peripheral tissues. As CHF progresses, structural and hemodynamic damages occur. While these damages have a variety of manifestations, one characteristic symptom is ventricular hypertrophy. CHF is a common end result of a number of various cardiac disorders.

“Myocardial infarction” generally results from atherosclerosis of the coronary arteries, often with superimposed coronary thrombosis. It may be divided into two major types: transmural infarcts, in which myocardial necrosis involves the full thickness of the ventricular wall, and subendocardial (nontransmural) infarcts, in which the necrosis involves the subendocardium, the intramural myocardium, or both, without extending all the way through the ventricular wall to the epicardium. Myocardial infarction is known to cause both a change in hemodynamic effects and an alteration in structure in the damaged and healthy zones of the heart. Thus, for example, myocardial infarction reduces the maximum cardiac output and the stroke volume of the heart. Also associated with myocardial infarction is a stimulation of the DNA synthesis occurring in the interstice as well as an increase in the formation of collagen in the areas of the heart not affected.

As a result of the increased stress or strain placed on the heart in prolonged hypertension due, for example, to the increased total peripheral resistance, cardiac hypertrophy has long been associated with “hypertension”. A

characteristic of the ventricle that becomes hypertrophic as a result of chronic pressure overload is an impaired diastolic performance. Fouad *et al.*, J. Am. Coll. Cardiol., 4: 1500-1506 (1984); Smith *et al.*, J. Am. Coll. Cardiol., 5: 869-874 (1985). A prolonged left ventricular relaxation has been detected in early essential hypertension, in spite of normal or supranormal systolic function. Hartford *et al.*, Hypertension, 6: 329-338 (1984). However, there is no close parallelism between blood pressure levels and cardiac hypertrophy. Although improvement in left ventricular function in response to antihypertensive therapy has been reported in humans, patients variously treated with a diuretic (hydrochlorothiazide), a  $\beta$ -blocker (propranolol), or a calcium channel blocker (diltiazem), have shown reversal of left ventricular hypertrophy, without improvement in diastolic function. Inouye *et al.*, Am. J. Cardiol., 53: 1583-7 (1984).

Another complex cardiac disease associated with cardiac hypertrophy is "hypertrophic cardiomyopathy". This condition is characterized by a great diversity of morphologic, functional, and clinical features (Maron *et al.*, N. Engl. J. Med., 316: 780-789 (1987); Spirito *et al.*, N. Engl. J. Med., 320: 749-755 (1989); Louie and Edwards, Prog. Cardiovasc. Dis., 36: 275-308 (1994); Wagle *et al.*, Circulation, 92: 1680-1692 (1995)), the heterogeneity of which is accentuated by the fact that it afflicts patients of all ages. Spirito *et al.*, N. Engl. J. Med., 336: 775-785 (1997). The causative factors of hypertrophic cardiomyopathy are also diverse and little understood. In general, mutations in genes encoding sarcomeric proteins are associated with hypertrophic cardiomyopathy. Recent data suggest that  $\beta$ -myosin heavy chain mutations may account for approximately 30 to 40 percent of cases of familial hypertrophic cardiomyopathy. Watkins *et al.*, N. Engl. J. Med., 326: 1108-1114 (1992); Schwartz *et al.*, Circulation, 91: 532-540 (1995); Marian and Roberts, Circulation, 92: 1336-1347 (1995); Thierfelder *et al.*, Cell, 77: 701-712 (1994); Watkins *et al.*, Nat. Gen., 11: 434-437 (1995). Besides  $\beta$ -myosin heavy chain, other locations of genetic mutations include cardiac troponin T, alpha topomyosin, cardiac myosin binding protein C, essential myosin light chain, and regulatory myosin light chain. See, Malik and Watkins, Curr. Opin. Cardiol., 12: 295-302 (1997).

Supravalvular "aortic stenosis" is an inherited vascular disorder characterized by narrowing of the ascending aorta, but other arteries, including the pulmonary arteries, may also be affected. Untreated aortic stenosis may lead to increased intracardiac pressure resulting in myocardial hypertrophy and eventually heart failure and death. The pathogenesis of this disorder is not fully understood, but hypertrophy and possibly hyperplasia of medial smooth muscle are prominent features of this disorder. It has been reported that molecular variants of the elastin gene are involved in the development and pathogenesis of aortic stenosis. U.S. Patent No. 5,650,282 issued July 22, 1997.

"Valvular regurgitation" occurs as a result of heart diseases resulting in disorders of the cardiac valves. Various diseases, like rheumatic fever, can cause the shrinking or pulling apart of the valve orifice, while other diseases may result in endocarditis, an inflammation of the endocardium or lining membrane of the atrioventricular orifices and operation of the heart. Defects such as the narrowing of the valve stenosis or the defective closing of the valve result in an accumulation of blood in the heart cavity or regurgitation of blood past the valve. If uncorrected, prolonged valvular stenosis or insufficiency may result in cardiac hypertrophy and associated damage to the heart muscle, which may eventually necessitate valve replacement.

The treatment of all these, and other cardiovascular, endothelial and angiogenic disorders, which may or may not be accompanied by cardiac hypertrophy, is encompassed by the present invention.

The terms "cancer", "cancerous", and "malignant" refer to or describe the physiological condition in mammals

that is typically characterized by unregulated cell growth. Examples of cancer include but are not limited to, carcinoma including adenocarcinoma, lymphoma, blastoma, melanoma, sarcoma, and leukemia. More particular examples of such cancers include squamous cell cancer, small-cell lung cancer, non-small cell lung cancer, gastrointestinal cancer, Hodgkin's and non-Hodgkin's lymphoma, pancreatic cancer, glioblastoma, cervical cancer, ovarian cancer, liver cancer such as hepatic carcinoma and hepatoma, bladder cancer, breast cancer, colon cancer, colorectal cancer, endometrial carcinoma, salivary gland carcinoma, kidney cancer such as renal cell carcinoma and Wilms' tumors, basal cell carcinoma, melanoma, prostate cancer, vulval cancer, thyroid cancer, testicular cancer, esophageal cancer, and various types of head and neck cancer. The preferred cancers for treatment herein are breast, colon, lung, melanoma, ovarian, and others involving vascular tumors as noted above.

The term "cytotoxic agent" as used herein refers to a substance that inhibits or prevents the function of cells and/or causes destruction of cells. The term is intended to include radioactive isotopes (e.g.,  $^{131}\text{I}$ ,  $^{125}\text{I}$ ,  $^{90}\text{Y}$ , and  $^{186}\text{Re}$ ), chemotherapeutic agents, and toxins such as enzymatically active toxins of bacterial, fungal, plant, or animal origin, or fragments thereof.

A "chemotherapeutic agent" is a chemical compound useful in the treatment of cancer. Examples of chemotherapeutic agents include alkylating agents, folic acid antagonists, anti-metabolites of nucleic acid metabolism, antibiotics, pyrimidine analogs, 5-fluorouracil, cisplatin, purine nucleosides, amines, amino acids, triazol nucleosides, or corticosteroids. Specific examples include Adriamycin, Doxorubicin, 5-Fluorouracil, Cytosine arabinoside ("Ara-C"), Cyclophosphamide, Thiotepe, Busulfan, Cytosine, Taxol, Toxotere, Methotrexate, Cisplatin, Melphalan, Vinblastine, Bleomycin, Etoposide, Ifosfamide, Mitomycin C, Mitoxantrone, Vincristine, Vinorelbine, Carboplatin, Teniposide, Daunomycin, Carminomycin, Aminopterin, Dactinomycin, Mitomycins, Esperamicins (see U.S. Pat. No. 4,675,187), Melphalan, and other related nitrogen mustards. Also included in this definition are hormonal agents that act to regulate or inhibit hormone action on tumors, such as tamoxifen and onapristone.

A "growth-inhibitory agent" when used herein refers to a compound or composition that inhibits growth of a cell, such as an Wnt-overexpressing cancer cell, either *in vitro* or *in vivo*. Thus, the growth-inhibitory agent is one which significantly reduces the percentage of malignant cells in S phase. Examples of growth-inhibitory agents include agents that block cell cycle progression (at a place other than S phase), such as agents that induce G1 arrest and M-phase arrest. Classical M-phase blockers include the vincas (vincristine and vinblastine), taxol, and topo II inhibitors such as doxorubicin, daunorubicin, etoposide, and bleomycin. Those agents that arrest G1 also spill over into S-phase arrest, for example, DNA alkylating agents such as tamoxifen, prednisone, dacarbazine, mechlorethamine, cisplatin, methotrexate, 5-fluorouracil, and ara-C. Further information can be found in The Molecular Basis of Cancer, Mendelsohn and Israel, eds., Chapter 1, entitled "Cell cycle regulation, oncogenes, and antineoplastic drugs" by Murakami *et al.* (WB Saunders: Philadelphia, 1995), especially p. 13. Additional examples include tumor necrosis factor (TNF), an antibody capable of inhibiting or neutralizing the angiogenic activity of acidic or basic FGF or hepatocyte growth factor (HGF), an antibody capable of inhibiting or neutralizing the coagulant activities of tissue factor, protein C, or protein S (see, WO 91/01753, published 21 February 1991), or an antibody capable of binding to HER2 receptor (WO 89/06692), such as the 4D5 antibody (and functional equivalents thereof) (e.g., WO 92/22653).

"Treatment" is an intervention performed with the intention of preventing the development or altering the pathology of a cardiovascular, endothelial, and angiogenic disorder. The concept of treatment is used in the broadest sense, and specifically includes the prevention (prophylaxis), moderation, reduction, and curing of cardiovascular, endothelial, and angiogenic disorders of any stage. Accordingly, "treatment" refers to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or slow down (lessen) a cardiovascular, endothelial, and angiogenic disorder such as hypertrophy. Those in need of treatment include those already with the disorder as well as those prone to have the disorder or those in whom the disorder is to be prevented. The disorder may result from any cause, including idiopathic, cardiopathic, or myopathic causes, or ischemia or ischemic insults, such as myocardial infarction.

"Chronic" administration refers to administration of the agent(s) in a continuous mode as opposed to an acute mode, so as to maintain the initial effect, such as an anti-hypertrophic effect, for an extended period of time.

"Mammal" for purposes of treatment refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, horses, cats, cows, sheep, pigs, etc. Preferably, the mammal is human.

Administration "in combination with" one or more further therapeutic agents includes simultaneous (concurrent) and consecutive administration in any order.

The phrase "cardiovascular, endothelial or angiogenic agents" refers generically to any drug that acts in treating cardiovascular, endothelial, and angiogenic disorders. Examples of cardiovascular agents are those that promote vascular homeostasis by modulating blood pressure, heart rate, heart contractility, and endothelial and smooth muscle biology, all of which factors have a role in cardiovascular disease. Specific examples of these include angiotensin-II receptor antagonists; endothelin receptor antagonists such as, for example, BOSENTAN<sup>TM</sup> and MOXONODIN<sup>TM</sup>; interferon-gamma (IFN- $\gamma$ ); des-aspartate-angiotensin I; thrombolytic agents, *e.g.*, streptokinase, urokinase, t-PA, and a t-PA variant specifically designed to have longer half-life and very high fibrin specificity, TNK t-PA (a T103N, N117Q, KHRR(296-299)AAAA t-PA variant, Keyt *et al.*, Proc. Natl. Acad. Sci. USA 91, 3670-3674 (1994)); inotropic or hypertensive agents such as digoxigenin and  $\beta$ -adrenergic receptor blocking agents, *e.g.*, propranolol, timolol, tertalolol, carteolol, nadolol, betaxolol, penbutolol, acetobutolol, atenolol, metoprolol, and carvedilol; angiotensin converting enzyme (ACE) inhibitors, *e.g.*, quinapril, captopril, enalapril, ramipril, benazepril, fosinopril, and lisinopril; diuretics, *e.g.*, chlorothiazide, hydrochlorothiazide, hydroflumethazide, methylchlorothiazide, benzthiazide, dichlorphenamide, acetazolamide, and indapamide; and calcium channel blockers, *e.g.*, diltiazem, nifedipine, verapamil, nicardipine. One preferred category of this type is a therapeutic agent used for the treatment of cardiac hypertrophy or of a physiological condition instrumental in the development of cardiac hypertrophy, such as elevated blood pressure, aortic stenosis, or myocardial infarction.

"Angiogenic agents" and "endothelial agents" are active agents that promote angiogenesis and/or endothelial cell growth, or, if applicable, vasculogenesis. This would include factors that accelerate wound healing, such as growth hormone, insulin-like growth factor-I (IGF-I), VEGF, VIGF, PDGF, epidermal growth factor (EGF), CTGF and members of its family, FGF, and TGF- $\alpha$  and TGF- $\beta$ .

"Angiostatic agents" are active agents that inhibit angiogenesis or vasculogenesis or otherwise inhibit or prevent growth of cancer cells. Examples include antibodies or other antagonists to angiogenic agents as defined

above, such as antibodies to VEGF. They additionally include cytotherapeutic agents such as cytotoxic agents, chemotherapeutic agents, growth-inhibitory agents, apoptotic agents, and other agents to treat cancer, such as anti-HER-2, anti-CD20, and other bioactive and organic chemical agents.

5 In a pharmacological sense, in the context of the present invention, a "therapeutically effective amount" of an active agent such as a PRO polypeptide or agonist or antagonist thereto or an anti-PRO antibody, refers to an amount effective in the treatment of a cardiovascular, endothelial or angiogenic disorder in a mammal and can be determined empirically.

10 As used herein, an "effective amount" of an active agent such as a PRO polypeptide or agonist or antagonist thereto or an anti-PRO antibody, refers to an amount effective for carrying out a stated purpose, wherein such amounts may be determined empirically for the desired effect.

15 The terms "PRO polypeptide" and "PRO" as used herein and when immediately followed by a numerical designation refer to various polypeptides, wherein the complete designation (*i.e.*, PRO/number) refers to specific polypeptide sequences as described herein. The terms "PRO/number polypeptide" and "PRO/number" wherein the term "number" is provided as an actual numerical designation as used herein encompass native sequence polypeptides and polypeptide variants (which are further defined herein). The PRO polypeptides described herein may be isolated from a variety of sources, such as from human tissue types or from another source, or prepared by recombinant or synthetic methods.

20 A "native sequence PRO polypeptide" comprises a polypeptide having the same amino acid sequence as the corresponding PRO polypeptide derived from nature. Such native sequence PRO polypeptides can be isolated from nature or can be produced by recombinant or synthetic means. The term "native sequence PRO polypeptide" specifically encompasses naturally-occurring truncated or secreted forms of the specific PRO polypeptide (*e.g.*, an extracellular domain sequence), naturally-occurring variant forms (*e.g.*, alternatively spliced forms) and naturally-occurring allelic variants of the polypeptide. In various embodiments of the invention, the native sequence PRO polypeptides disclosed herein are mature or full-length native sequence polypeptides comprising the full-length amino acids sequences shown in the accompanying figures. Start and stop codons are shown in bold font and underlined in the figures. However, while the PRO polypeptide disclosed in the accompanying figures are shown to begin with methionine residues designated herein as amino acid position 1 in the figures, it is conceivable and possible that other methionine residues located either upstream or downstream from the amino acid position 1 in the figures may be employed as the starting amino acid residue for the PRO polypeptides.

30 The PRO polypeptide "extracellular domain" or "ECD" refers to a form of the PRO polypeptide which is essentially free of the transmembrane and cytoplasmic domains. Ordinarily, a PRO polypeptide ECD will have less than 1% of such transmembrane and/or cytoplasmic domains and preferably, will have less than 0.5% of such domains. It will be understood that any transmembrane domains identified for the PRO polypeptides of the present invention are identified pursuant to criteria routinely employed in the art for identifying that type of hydrophobic domain. The exact boundaries of a transmembrane domain may vary but most likely by no more than about 5 amino acids at either end of the domain as initially identified herein. Optionally, therefore, an extracellular domain of a PRO polypeptide may contain from about 5 or fewer amino acids on either side of the transmembrane domain/extracellular domain boundary as identified in the Examples or specification and such polypeptides, with

35



or without the associated signal peptide, and nucleic acid encoding them, are contemplated by the present invention.

The approximate location of the "signal peptides" of the various PRO polypeptides disclosed herein are shown in the present specification and/or the accompanying figures. It is noted, however, that the C-terminal boundary of a signal peptide may vary, but most likely by no more than about 5 amino acids on either side of the signal peptide C-terminal boundary as initially identified herein, wherein the C-terminal boundary of the signal peptide may be identified pursuant to criteria routinely employed in the art for identifying that type of amino acid sequence element (*e.g.*, Nielsen *et al.*, Prot. Eng., 10:1-6 (1997) and von Heinje *et al.*, Nucl. Acids Res., 14:4683-4690 (1986)). Moreover, it is also recognized that, in some cases, cleavage of a signal sequence from a secreted polypeptide is not entirely uniform, resulting in more than one secreted species. These mature polypeptides, where the signal peptide is cleaved within no more than about 5 amino acids on either side of the C-terminal boundary of the signal peptide as identified herein, and the polynucleotides encoding them, are contemplated by the present invention.

"PRO polypeptide variant" means an active PRO polypeptide as defined above or below having at least about 80% amino acid sequence identity with a full-length native sequence PRO polypeptide sequence as disclosed herein, a PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Such PRO polypeptide variants include, for instance, PRO polypeptides wherein one or more amino acid residues are added, or deleted, at the N- or C-terminus of the full-length native amino acid sequence. Ordinarily, a PRO polypeptide variant will have at least about 80% amino acid sequence identity, preferably at least about 81% amino acid sequence identity, more preferably at least about 82% amino acid sequence identity, more preferably at least about 83% amino acid sequence identity, more preferably at least about 84% amino acid sequence identity, more preferably at least about 85% amino acid sequence identity, more preferably at least about 86% amino acid sequence identity, more preferably at least about 87% amino acid sequence identity, more preferably at least about 88% amino acid sequence identity, more preferably at least about 89% amino acid sequence identity, more preferably at least about 90% amino acid sequence identity, more preferably at least about 91% amino acid sequence identity, more preferably at least about 92% amino acid sequence identity, more preferably at least about 93% amino acid sequence identity, more preferably at least about 94% amino acid sequence identity, more preferably at least about 95% amino acid sequence identity, more preferably at least about 96% amino acid sequence identity, more preferably at least about 97% amino acid sequence identity, more preferably at least about 98% amino acid sequence identity and most preferably at least about 99% amino acid sequence identity with a full-length native sequence PRO polypeptide sequence as disclosed herein, a PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of a full-length PRO polypeptide sequence as disclosed herein. Ordinarily, PRO variant polypeptides are at least about 10 amino acids in length, often at least about 20 amino acids in length, more often at least about 30 amino acids in length, more often at least about 40 amino acids in length, more often at least about 50 amino acids in length, more often at least about 60 amino acids in length, more often at least about 70 amino acids in length, more often at least about 80 amino acids in length, more often at least about 90 amino acids in length, more often at least about 100 amino acids in length, more often

at least about 150 amino acids in length, more often at least about 200 amino acids in length, more often at least about 300 amino acids in length, or more.

As shown below, Table 1 provides the complete source code for the ALIGN-2 sequence comparison computer program. This source code may be routinely compiled for use on a UNIX operating system to provide the ALIGN-2 sequence comparison computer program.

In addition, Tables 2A-2D show hypothetical exemplifications for using the below described method to determine % amino acid sequence identity (Tables 2A-2B) and % nucleic acid sequence identity (Tables 2C-2D) using the ALIGN-2 sequence comparison computer program, wherein "PRO" represents the amino acid sequence of a hypothetical PRO polypeptide of interest, "Comparison Protein" represents the amino acid sequence of a polypeptide against which the "PRO" polypeptide of interest is being compared, "PRO-DNA" represents a hypothetical PRO-encoding nucleic acid sequence of interest, "Comparison DNA" represents the nucleotide sequence of a nucleic acid molecule against which the "PRO-DNA" nucleic acid molecule of interest is being compared, "X", "Y", and "Z" each represent different hypothetical amino acid residues and "N", "L" and "V" each represent different hypothetical nucleotides.

Table 1

```

/*
 *
 * C-C increased from 12 to 15
 * Z is average of EQ
 * B is average of ND
 * match with stop is _M; stop-stop = 0; J (joker) match = 0
 */
#define _M      -8      /* value of a match with a stop */

int _day[26][26] = {
/*   A B C D E F G H I J K L M N O P Q R S T U V W X Y Z */
/* A */ { 2, 0, -2, 0, 0, -4, 1, -1, -1, 0, -1, -2, -1, 0, _M, 1, 0, -2, 1, 1, 0, 0, -6, 0, -3, 0},
/* B */ { 0, 3, -4, 3, 2, -5, 0, 1, -2, 0, 0, -3, -2, 2, _M, -1, 1, 0, 0, 0, 0, -2, -5, 0, -3, 1},
/* C */ {-2, -4, 15, -5, -5, -4, -3, -3, -2, 0, -5, -6, -5, -4, _M, -3, -5, -4, 0, -2, 0, -2, -8, 0, 0, -5},
/* D */ { 0, 3, -5, 4, 3, -6, 1, 1, -2, 0, 0, -4, -3, 2, _M, -1, 2, -1, 0, 0, 0, -2, -7, 0, -4, 2},
/* E */ { 0, 2, -5, 3, 4, -5, 0, 1, -2, 0, 0, -3, -2, 1, _M, -1, 2, -1, 0, 0, 0, -2, -7, 0, -4, 3},
/* F */ {-4, -5, -4, -6, -5, 9, -5, -2, 1, 0, -5, 2, 0, -4, _M, -5, -5, -4, -3, -3, 0, -1, 0, 0, 7, -5},
/* G */ { 1, 0, -3, 1, 0, -5, 5, -2, -3, 0, -2, -4, -3, 0, _M, -1, -1, -3, 1, 0, 0, -1, -7, 0, -5, 0},
/* H */ {-1, 1, -3, 1, 1, -2, -2, 6, -2, 0, 0, -2, -2, 2, _M, 0, 3, 2, -1, -1, 0, -2, -3, 0, 0, 2},
/* I */ {-1, -2, -2, -2, -2, 1, -3, -2, 5, 0, -2, 2, 2, -2, _M, -2, -2, -2, -1, 0, 0, 4, -5, 0, -1, -2},
/* J */ { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* K */ {-1, 0, -5, 0, 0, -5, -2, 0, -2, 0, 5, -3, 0, 1, _M, -1, 1, 3, 0, 0, 0, -2, -3, 0, -4, 0},
/* L */ {-2, -3, -6, -4, -3, 2, -4, -2, 2, 0, -3, 6, 4, -3, _M, -3, -2, -3, -3, -1, 0, 2, -2, 0, -1, -2},
/* M */ {-1, -2, -5, -3, -2, 0, -3, -2, 2, 0, 0, 4, 6, -2, _M, -2, -1, 0, -2, -1, 0, 2, -4, 0, -2, -1},
/* N */ { 0, 2, -4, 2, 1, -4, 0, 2, -2, 0, 1, -3, -2, 2, _M, -1, 1, 0, 1, 0, 0, -2, -4, 0, -2, 1},
/* O */ {_M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, 0, _M, _M, _M, _M, _M, _M, _M, _M, _M},
/* P */ { 1, -1, -3, -1, -1, -5, -1, 0, -2, 0, -1, -3, -2, -1, _M, 6, 0, 0, 1, 0, 0, -1, -6, 0, -5, 0},
/* Q */ { 0, 1, -5, 2, 2, -5, -1, 3, -2, 0, 1, -2, -1, 1, _M, 0, 4, 1, -1, -1, 0, -2, -5, 0, -4, 3},
/* R */ {-2, 0, -4, -1, -1, -4, -3, 2, -2, 0, 3, -3, 0, 0, _M, 0, 1, 6, 0, -1, 0, -2, 2, 0, -4, 0},
/* S */ { 1, 0, 0, 0, 0, -3, 1, -1, -1, 0, 0, -3, -2, 1, _M, 1, -1, 0, 2, 1, 0, -1, -2, 0, -3, 0},
/* T */ { 1, 0, -2, 0, 0, -3, 0, -1, 0, 0, 0, -1, -1, 0, _M, 0, -1, -1, 1, 3, 0, 0, -5, 0, -3, 0},
/* U */ { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* V */ { 0, -2, -2, -2, -2, -1, -1, -2, 4, 0, -2, 2, 2, -2, _M, -1, -2, -2, -1, 0, 0, 4, -6, 0, -2, -2},
/* W */ {-6, -5, -8, -7, -7, 0, -7, -3, -5, 0, -3, -2, -4, -4, _M, -6, -5, 2, -2, -5, 0, -6, 17, 0, 0, -6},
/* X */ { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* Y */ {-3, -3, 0, -4, -4, 7, -5, 0, -1, 0, -4, -1, -2, -2, _M, -5, -4, -4, -3, -3, 0, -2, 0, 0, 10, -4},
/* Z */ { 0, 1, -5, 2, 3, -5, 0, 2, -2, 0, 0, -2, -1, 1, _M, 0, 3, 0, 0, 0, 0, -2, -6, 0, -4, 4}
};

```

```

/*
*/
#include <stdio.h>
#include <ctype.h>

#define MAXJMP      16      /* max jumps in a diag */
#define MAXGAP      24      /* don't continue to penalize gaps larger than this */
#define JMPS        1024    /* max jmps in an path */
#define MX          4       /* save if there's at least MX-1 bases since last jmp */

#define DMAT        3       /* value of matching bases */
#define DMIS        0       /* penalty for mismatched bases */
#define DINS0       8       /* penalty for a gap */
#define DINS1       1       /* penalty per base */
#define PINS0       8       /* penalty for a gap */
#define PINS1       4       /* penalty per residue */

struct jmp {
    short          n[MAXJMP];    /* size of jmp (neg for dely) */
    unsigned short x[MAXJMP];    /* base no. of jmp in seq x */
};
/* limits seq to 2^16 -1 */

struct diag {
    int            score;        /* score at last jmp */
    long           offset;       /* offset of prev block */
    short          jmp;         /* current jmp index */
    struct jmp     jp;          /* list of jmps */
};

struct path {
    int            spc;          /* number of leading spaces */
    short          n[JMPS];      /* size of jmp (gap) */
    int            x[JMPS];      /* loc of jmp (last elem before gap) */
};

char    *ofile;                /* output file name */
char    *namex[2];              /* seq names: getseqs() */
char    *prog;                  /* prog name for err msgs */
char    *seqx[2];               /* seqs: getseqs() */
int      dmax;                  /* best diag: nw() */
int      dmax0;                 /* final diag */
int      dna;                   /* set if dna: main() */
int      endgaps;               /* set if penalizing end gaps */
int      gapx, gapy;             /* total gaps in seqs */
int      len0, len1;             /* seq lens */
int      ngapx, ngapy;           /* total size of gaps */
int      smax;                  /* max score: nw() */
int      *xbm;                  /* bitmap for matching */
long     offset;                /* current offset in jmp file */
struct   diag    dx;             /* holds diagonals */
struct   path    pp[2];          /* holds path for seqs */

char    *calloc(), *malloc(), *index(), *strcpy();
char    *getseq(), *g_calloc();

```

```

/* Needleman-Wunsch alignment program
*
* usage: progs file1 file2
* where file1 and file2 are two dna or two protein sequences.
* The sequences can be in upper- or lower-case and may contain ambiguity
* Any lines beginning with ';', '>' or '<' are ignored
* Max file length is 65535 (limited by unsigned short x in the jmp struct)
* A sequence with 1/3 or more of its elements ACGTU is assumed to be DNA
* Output is in the file "align.out"
*
* The program may create a tmp file in /tmp to hold info about traceback.
* Original version developed under BSD 4.3 on a vax 8650
*/
#include "nw.h"
#include "day.h"

static _dbval[26] = {
    1,14,2,13,0,0,4,11,0,0,12,0,3,15,0,0,0,5,6,8,8,7,9,0,10,0
};

static _pbval[26] = {
    1, 2[(1 < ('D'-'A'))|(1 < ('N'-'A'))], 4, 8, 16, 32, 64,
    128, 256, 0xFFFFFFFF, 1 < 10, 1 < 11, 1 < 12, 1 < 13, 1 < 14,
    1 < 15, 1 < 16, 1 < 17, 1 < 18, 1 < 19, 1 < 20, 1 < 21, 1 < 22,
    1 < 23, 1 < 24, 1 < 25[(1 < ('E'-'A'))|(1 < ('Q'-'A'))]
};

main(ac, av)
int ac;
char *av[];
{
    prog = av[0];
    if (ac != 3) {
        fprintf(stderr, "usage: %s file1 file2\n", prog);
        fprintf(stderr, "where file1 and file2 are two dna or two protein sequences.\n");
        fprintf(stderr, "The sequences can be in upper- or lower-case\n");
        fprintf(stderr, "Any lines beginning with ';' or '<' are ignored\n");
        fprintf(stderr, "Output is in the file 'align.out'\n");
        exit(1);
    }
    namex[0] = av[1];
    namex[1] = av[2];
    seqx[0] = getseq(namex[0], &len0);
    seqx[1] = getseq(namex[1], &len1);
    xbm = (dna)? _dbval : _pbval;

    endgaps = 0; /* 1 to penalize endgaps */
    ofile = "align.out"; /* output file */

    nw(); /* fill in the matrix, get the possible jumps */
    readjumps(); /* get the actual jumps */
    print(); /* print stats, alignment */

    cleanup(); /* unlink any tmp files */
}

```

```

/* do the alignment, return best score: main()
 * dna: values in Fitch and Smith, PNAS, 80, 1382-1386, 1983
 * pro: PAM 250 values
 * When scores are equal, we prefer mismatches to any gap, prefer
 * a new gap to extending an ongoing gap, and prefer a gap in seqx
 * to a gap in seq y.
 */
nw()
{
    char          *px, *py;          /* seqs and ptrs */
    int           *ndely, *dely;      /* keep track of dely */
    int           ndelx, delx;        /* keep track of delx */
    int           *tmp;               /* for swapping row0, row1 */
    int           mis;                /* score for each type */
    int           ins0, ins1;         /* insertion penalties */
    register      id;                 /* diagonal index */
    register      ij;                 /* jmp index */
    register      *col0, *col1;       /* score for curr, last row */
    register      xx, yy;             /* index into seqs */

    dx = (struct diag *)g_calloc("to get diags", len0+len1+1, sizeof(struct diag));

    ndely = (int *)g_calloc("to get ndely", len1+1, sizeof(int));
    dely = (int *)g_calloc("to get dely", len1+1, sizeof(int));
    col0 = (int *)g_calloc("to get col0", len1+1, sizeof(int));
    col1 = (int *)g_calloc("to get col1", len1+1, sizeof(int));
    ins0 = (dna)? DINS0 : PINS0;
    ins1 = (dna)? DINS1 : PINS1;

    smax = -10000;
    if (endgaps) {
        for (col0[0] = dely[0] = -ins0, yy = 1; yy <= len1; yy++) {
            col0[yy] = dely[yy] = col0[yy-1] - ins1;
            ndely[yy] = yy;
        }
        col0[0] = 0;          /* Waterman Bull Math Biol 84 */
    }
    else
        for (yy = 1; yy <= len1; yy++)
            dely[yy] = -ins0;

    /* fill in match matrix
     */
    for (px = seqx[0], xx = 1; xx <= len0; px++, xx++) {
        /* initialize first entry in col
         */
        if (endgaps) {
            if (xx == 1)
                col1[0] = delx = -(ins0+ins1);
            else
                col1[0] = delx = col0[0] - ins1;
            ndelx = xx;
        }
        else {
            col1[0] = 0;
            delx = -ins0;
            ndelx = 0;
        }
    }
}

```

...nw

```

for (py = seqx[1], yy = 1; yy <= len1; py++, yy++) {
    mis = col0[yy-1];
    if (dna)
        mis += (xbm[*px-'A']&xbm[*py-'A'])? DMAT : DMIS;
    else
        mis += _day[*px-'A'][*py-'A'];

    /* update penalty for del in x seq;
     * favor new del over ongoing del
     * ignore MAXGAP if weighting endgaps
     */
    if (endgaps || ndely[yy] < MAXGAP) {
        if (col0[yy] - ins0 >= dely[yy]) {
            dely[yy] = col0[yy] - (ins0+ins1);
            ndely[yy] = 1;
        } else {
            dely[yy] -= ins1;
            ndely[yy]++;
        }
    } else {
        if (col0[yy] - (ins0+ins1) >= dely[yy]) {
            dely[yy] = col0[yy] - (ins0+ins1);
            ndely[yy] = 1;
        } else
            ndely[yy]++;
    }

    /* update penalty for del in y seq;
     * favor new del over ongoing del
     */
    if (endgaps || ndelx < MAXGAP) {
        if (col1[yy-1] - ins0 >= delx) {
            delx = col1[yy-1] - (ins0+ins1);
            ndelx = 1;
        } else {
            delx -= ins1;
            ndelx++;
        }
    } else {
        if (col1[yy-1] - (ins0+ins1) >= delx) {
            delx = col1[yy-1] - (ins0+ins1);
            ndelx = 1;
        } else
            ndelx++;
    }

    /* pick the maximum score; we're favoring
     * mis over any del and delx over dely
     */

```

...nw

```

        id = xx - yy + len1 - 1;
        if (mis >= delx && mis >= dely[yy])
            coll[yy] = mis;
        else if (delx >= dely[yy]) {
            coll[yy] = delx;
            ij = dx[id].ijmp;
            if (dx[id].jp.n[0] && (!dna || (ndelx >= MAXJMP
            && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
                dx[id].ijmp++;
                if (++ij >= MAXJMP) {
                    writejumps(id);
                    ij = dx[id].ijmp = 0;
                    dx[id].offset = offset;
                    offset += sizeof(struct jmp) + sizeof(offset);
                }
            }
            dx[id].jp.n[ij] = ndelx;
            dx[id].jp.x[ij] = xx;
            dx[id].score = delx;
        }
        else {
            coll[yy] = dely[yy];
            ij = dx[id].ijmp;

        if (dx[id].jp.n[0] && (!dna || (ndely[yy] >= MAXJMP
            && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
                dx[id].ijmp++;
                if (++ij >= MAXJMP) {
                    writejumps(id);
                    ij = dx[id].ijmp = 0;
                    dx[id].offset = offset;
                    offset += sizeof(struct jmp) + sizeof(offset);
                }
            }
            dx[id].jp.n[ij] = -ndely[yy];
            dx[id].jp.x[ij] = xx;
            dx[id].score = dely[yy];
        }
        if (xx == len0 && yy < len1) {
            /* last col
            */
            if (endgaps)
                coll[yy] -= ins0+ins1*(len1-yy);
            if (coll[yy] > smax) {
                smax = coll[yy];
                dmax = id;
            }
        }
    }
    if (endgaps && xx < len0)
        coll[yy-1] -= ins0+ins1*(len0-xx);
    if (coll[yy-1] > smax) {
        smax = coll[yy-1];
        dmax = id;
    }
    tmp = col0; col0 = coll; coll = tmp;
}
(void) free((char *)ndely);
(void) free((char *)dely);
(void) free((char *)col0);
(void) free((char *)coll);
}

```



```

/*
 *
 * print() -- only routine visible outside this module
 *
 * static:
 * getmat() -- trace back best path, count matches: print()
 * pr_align() -- print alignment of described in array p[]: print()
 * dumpblock() -- dump a block of lines with numbers, stars: pr_align()
 * nums() -- put out a number line: dumpblock()
 * putline() -- put out a line (name, [num], seq, [num]): dumpblock()
 * stars() -- put a line of stars: dumpblock()
 * stripname() -- strip any path and prefix from a seqname
 */

#include "nw.h"

#define SPC 3
#define P_LINE 256 /* maximum output line */
#define P_SPC 3 /* space between name or num and seq */

extern _day[26][26];
int olen; /* set output line length */
FILE *fx; /* output file */

print()
{
    int lx, ly, firstgap, lastgap; /* overlap */

    if ((fx = fopen(ofile, "w")) == 0) {
        fprintf(stderr, "%s: can't write %s\n", prog, ofile);
        cleanup(1);
    }
    fprintf(fx, "< first sequence: %s (length = %d)\n", namex[0], len0);
    fprintf(fx, "< second sequence: %s (length = %d)\n", namex[1], len1);
    olen = 60;
    lx = len0;
    ly = len1;
    firstgap = lastgap = 0;
    if (dmax < len1 - 1) { /* leading gap in x */
        pp[0].spc = firstgap = len1 - dmax - 1;
        ly -= pp[0].spc;
    }
    else if (dmax > len1 - 1) { /* leading gap in y */
        pp[1].spc = firstgap = dmax - (len1 - 1);
        lx -= pp[1].spc;
    }
    if (dmax0 < len0 - 1) { /* trailing gap in x */
        lastgap = len0 - dmax0 - 1;
        lx -= lastgap;
    }
    else if (dmax0 > len0 - 1) { /* trailing gap in y */
        lastgap = dmax0 - (len0 - 1);
        ly -= lastgap;
    }
    getmat(lx, ly, firstgap, lastgap);
    pr_align();
}

```

print

```

/*
 * trace back the best path, count matches
 */
static
getmat(lx, ly, firstgap, lastgap)                                getmat
{
    int lx, ly;                                /* "core" (minus endgaps) */
    int firstgap, lastgap;                      /* leading trailing overlap */

    int nm, i0, i1, siz0, siz1;
    char outx[32];
    double pct;
    register n0, n1;
    register char *p0, *p1;

    /* get total matches, score
    */
    i0 = i1 = siz0 = siz1 = 0;
    p0 = seqx[0] + pp[1].spc;
    p1 = seqx[1] + pp[0].spc;
    n0 = pp[1].spc + 1;
    n1 = pp[0].spc + 1;

    nm = 0;
    while ( *p0 && *p1 ) {
        if (siz0) {
            p1++;
            n1++;
            siz0--;
        }
        else if (siz1) {
            p0++;
            n0++;
            siz1--;
        }
        else {
            if (xbm[*p0-'A']&xbm[*p1-'A'])
                nm++;
            if (n0++ == pp[0].x[i0])
                siz0 = pp[0].n[i0++];
            if (n1++ == pp[1].x[i1])
                siz1 = pp[1].n[i1++];
            p0++;
            p1++;
        }
    }

    /* pct homology:
    * if penalizing endgaps, base is the shorter seq
    * else, knock off overhangs and take shorter core
    */
    if (endgaps)
        lx = (len0 < len1)? len0 : len1;
    else
        lx = (lx < ly)? lx : ly;
    pct = 100.*((double)nm)/((double)lx);
    fprintf(fx, "\n");
    fprintf(fx, "< %d match%s in an overlap of %d: %.2f percent similarity\n",
        nm, (nm == 1)? "" : "es", lx, pct);
}

```

```

fprintf(fx, "< gaps in first sequence: %d", gapx);
if (gapx) {
    (void) sprintf(outx, " (%d %s%s)",
        ngapx, (dna)? "base": "residue", (ngapx == 1)? "" : "s");
    fprintf(fx, "%s", outx);

    fprintf(fx, ", gaps in second sequence: %d", gapy);
    if (gapy) {
        (void) sprintf(outx, " (%d %s%s)",
            ngapy, (dna)? "base": "residue", (ngapy == 1)? "" : "s");
        fprintf(fx, "%s", outx);
    }
    if (dna)
        fprintf(fx,
            "\n< score: %d (match = %d, mismatch = %d, gap penalty = %d + %d per base)\n",
            smax, DMAT, DMIS, DINS0, DINS1);
    else
        fprintf(fx,
            "\n< score: %d (Dayhoff PAM 250 matrix, gap penalty = %d + %d per residue)\n",
            smax, PINS0, PINS1);
    if (endgaps)
        fprintf(fx,
            "< endgaps penalized. left endgap: %d %s%s, right endgap: %d %s%s\n",
            firstgap, (dna)? "base" : "residue", (firstgap == 1)? "" : "s",
            lastgap, (dna)? "base" : "residue", (lastgap == 1)? "" : "s");
    else
        fprintf(fx, "< endgaps not penalized\n");
}

static      nm;          /* matches in core -- for checking */
static      lmax;        /* lengths of stripped file names */
static      ij[2];       /* jmp index for a path */
static      nc[2];       /* number at start of current line */
static      ni[2];       /* current elem number -- for gapping */
static      siz[2];
static char *ps[2];      /* ptr to current element */
static char *po[2];      /* ptr to next output char slot */
static char out[2][P_LINE]; /* output line */
static char star[P_LINE]; /* set by stars() */

/*
 * print alignment of described in struct path pp[]
 */
static
pr_align()
{
    int      nn;          /* char count */
    int      more;
    register i;

    for (i = 0, lmax = 0; i < 2; i++) {
        nn = stripname(namecx[i]);
        if (nn > lmax)
            lmax = nn;

        nc[i] = 1;
        ni[i] = 1;
        siz[i] = ij[i] = 0;
        ps[i] = seqx[i];
        po[i] = out[i];
    }
}

```

...getmat

pr\_align

```

for (nn = nm = 0, more = 1; more; ) {
    for (i = more = 0; i < 2; i++) {
        /*
         * do we have more of this sequence?
         */
        if (!*ps[i])
            continue;

        more++;

        if (pp[i].spc) { /* leading space */
            *po[i]++ = ' ';
            pp[i].spc--;
        }
        else if (siz[i]) { /* in a gap */
            *po[i]++ = '-';
            siz[i]--;
        }
        else { /* we're putting a seq element
                */
            *po[i] = *ps[i];
            if (islower(*ps[i]))
                *ps[i] = toupper(*ps[i]);
            po[i]++;
            ps[i]++;

            /*
             * are we at next gap for this seq?
             */
            if (ni[i] == pp[i].x[ij[i]]) {
                /*
                 * we need to merge all gaps
                 * at this location
                 */
                siz[i] = pp[i].n[ij[i] + +];
                while (ni[i] == pp[i].x[ij[i]])
                    siz[i] += pp[i].n[ij[i] + +];
            }
            ni[i]++;
        }
    }
    if (++nn == olen || !more && nn) {
        dumpblock();
        for (i = 0; i < 2; i++)
            po[i] = out[i];
        nn = 0;
    }
}

/*
 * dump a block of lines, including numbers, stars: pr_align()
 */
static
dumpblock()
{
    register    i;

    for (i = 0; i < 2; i++)
        *po[i]-- = '\0';
}

```

...pr\_align

dumpblock

...dumpblock

```

(void) putc('\n', fx);
for (i = 0; i < 2; i++) {
    if (*out[i] && (*out[i] != ' ' || *(po[i]) != ' ')) {
        if (i == 0)
            nums(i);
        if (i == 0 && *out[1])
            stars();
        putline(i);
        if (i == 0 && *out[1])
            fprintf(fx, star);
        if (i == 1)
            nums(i);
    }
}
}

/*
 * put out a number line: dumpblock()
 */
static
nums(ix)
    int ix;        /* index in out[] holding seq line */
{
    char          nline[P_LINE];
    register      i, j;
    register char *pn, *px, *py;

    for (pn = nline, i = 0; i < lmax + P_SPC; i++, pn++)
        *pn = ' ';
    for (i = nc[ix], py = out[ix]; *py; py++, pn++) {
        if (*py == ' ' || *py == '-')
            *pn = ' ';
        else {
            if (i % 10 == 0 || (i == 1 && nc[ix] != 1)) {
                j = (i < 0)? -i : i;
                for (px = pn; j /= 10, px--)
                    *px = j % 10 + '0';
                if (i < 0)
                    *px = '-';
            }
            else
                *pn = ' ';
            i++;
        }
    }
    *pn = '\0';
    nc[ix] = i;
    for (pn = nline; *pn; pn++)
        (void) putc(*pn, fx);
    (void) putc('\n', fx);
}

/*
 * put out a line (name, [num], seq, [num]): dumpblock()
 */
static
putline(ix)
    int ix;
{

```

nums

putline

...putline

```

int      i;
register char      *px;

for (px = namex[ix], i = 0; *px && *px != ':'; px++, i++)
    (void) putc(*px, fx);
for (; i < lmax+P_SPC; i++)
    (void) putc(' ', fx);

/* these count from 1:
 * ni[] is current element (from 1)
 * nc[] is number at start of current line
 */
for (px = out[ix]; *px; px++)
    (void) putc(*px&0x7F, fx);
(void) putc('\n', fx);
}

```

```

/*
 * put a line of stars (seqs always in out[0], out[1]): dumpblock()
 */

```

static

stars()

stars

```

{
    int      i;
    register char      *p0, *p1, cx, *px;

    if (!*out[0] || (*out[0] == ' ' && *(po[0]) == ' ') ||
        !*out[1] || (*out[1] == ' ' && *(po[1]) == ' '))
        return;
    px = star;
    for (i = lmax+P_SPC; i--;)
        *px++ = ' ';

    for (p0 = out[0], p1 = out[1]; *p0 && *p1; p0++, p1++) {
        if (isalpha(*p0) && isalpha(*p1)) {
            if (xbm[*p0-'A']&xbm[*p1-'A']) {
                cx = '*';
                nm++;
            }
            else if (!dna && _day[*p0-'A'][*p1-'A'] > 0)
                cx = '.';
            else
                cx = ' ';
        }
        else
            cx = ' ';
        *px++ = cx;
    }
    *px++ = '\n';
    *px = '\0';
}

```

```
/*
 * strip path or prefix from pn, return len: pr_align()
 */
static
stripname(pn)
    char      *pn;      /* file name (may be path) */
{
    register char      *px, *py;

    py = 0;
    for (px = pn; *px; px++)
        if (*px == '/')
            py = px + 1;
    if (py)
        (void) strcpy(pn, py);
    return(strlen(pn));
}
```

**stripname**

```

/*
 * cleanup() -- cleanup any tmp file
 * getseq() -- read in seq, set dna, len, maxlen
 * g_calloc() -- calloc() with error checkin
 * readjumps() -- get the good jumps, from tmp file if necessary
 * writejumps() -- write a filled array of jumps to a tmp file: nw()
 */
#include "nw.h"
#include <sys/file.h>

char *jname = "/tmp/homgXXXXXX";          /* tmp file for jumps */
FILE *fj;

int cleanup();                             /* cleanup tmp file */
long lseek();

/*
 * remove any tmp file if we blow
 */
cleanup(i)                                cleanup
{
    int i;
    {
        if (fj)
            (void) unlink(jname);
        exit(i);
    }
}

/*
 * read, return ptr to seq, set dna, len, maxlen
 * skip lines starting with ';', '<', or '>'
 * seq in upper or lower case
 */
char *
getseq(file, lcn)                          getseq
{
    char *file;          /* file name */
    int *len;            /* seq len */
    {
        char line[1024], *pseq;
        register char *px, *py;
        int natgc, tlen;
        FILE *fp;

        if ((fp = fopen(file, "r")) == 0) {
            fprintf(stderr, "%s: can't read %s\n", prog, file);
            exit(1);
        }
        tlen = natgc = 0;
        while (fgets(line, 1024, fp)) {
            if (*line == ';' || *line == '<' || *line == '>')
                continue;
            for (px = line; *px != '\n'; px++)
                if (isupper(*px) || islower(*px))
                    tlen++;
        }
        if ((pseq = malloc((unsigned)(tlen+6))) == 0) {
            fprintf(stderr, "%s: malloc() failed to get %d bytes for %s\n", prog, tlen+6, file);
            exit(1);
        }
        pseq[0] = pseq[1] = pseq[2] = pseq[3] = '\0';
    }
}

```



...getseq

```

py = pseq + 4;
*len = tlen;
rewind(fp);

while (fgets(line, 1024, fp)) {
    if (*line == ';' || *line == '<' || *line == '>')
        continue;
    for (px = line; *px != '\n'; px++) {
        if (isupper(*px))
            *py++ = *px;
        else if (islower(*px))
            *py++ = toupper(*px);
        if (index("ATGCU", *(py-1)))
            natgc++;
    }
}
*py++ = '\0';
*py = '\0';
(void) fclose(fp);
dna = natgc > (tlen/3);
return(pseq+4);
}

```

char \*

g\_calloc(msg, nx, sz)

g\_calloc

```

char *msg; /* program, calling routine */
int nx, sz; /* number and size of elements */
{
    char *px, *calloc();

    if ((px = calloc((unsigned)nx, (unsigned)sz)) == 0) {
        if (*msg) {
            fprintf(stderr, "%s: g_calloc() failed %s (n=%d, sz=%d)\n", prog, msg, nx, sz);
            exit(1);
        }
    }
    return(px);
}

```

/\*

\* get final jmps from dx[] or tmp file, set pp[], reset dmax: main()

\*/

readjmps()

readjmps

```

{
    int fd = -1;
    int siz, i0, i1;
    register i, j, xx;

    if (fj) {
        (void) fclose(fj);
        if ((fd = open(jname, O_RDONLY, 0)) < 0) {
            fprintf(stderr, "%s: can't open() %s\n", prog, jname);
            cleanup(1);
        }
    }
    for (i = i0 = i1 = 0, dmax0 = dmax, xx = len0; i++) {
        while (1) {
            for (j = dx[dmax].ijmp; j >= 0 && dx[dmax].jp.x[j] >= xx; j--)
                ;
        }
    }
}

```

Page 2 of nwsubr.c

## ...readjumps

```

        if (j < 0 && dx[dmax].offset && fj) {
            (void) lseek(fd, dx[dmax].offset, 0);
            (void) read(fd, (char *)&dx[dmax].jp, sizeof(struct jmp));
            (void) read(fd, (char *)&dx[dmax].offset, sizeof(dx[dmax].offset));
            dx[dmax].ijmp = MAXJMP-1;
        }
        else
            break;
    }
    if (i >= JMPS) {
        fprintf(stderr, "%s: too many gaps in alignment\n", prog);
        cleanup(1);
    }
    if (j >= 0) {
        siz = dx[dmax].jp.n[j];
        xx = dx[dmax].jp.x[j];
        dmax += siz;
        if (siz < 0) { /* gap in second seq */
            pp[1].n[i1] = -siz;
            xx += siz;

            /* id = xx - yy + len1 - 1
             */
            pp[1].x[i1] = xx - dmax + len1 - 1;
            gapy++;
            ngapy -= siz;
        }
        /* ignore MAXGAP when doing endgaps */
        siz = (-siz < MAXGAP || endgaps)? -siz : MAXGAP;
        i1++;
    }
    else if (siz > 0) { /* gap in first seq */
        pp[0].n[i0] = siz;
        pp[0].x[i0] = xx;
        gapx++;
        ngapx += siz;
        /* ignore MAXGAP when doing endgaps */
        siz = (siz < MAXGAP || endgaps)? siz : MAXGAP;
        i0++;
    }
    }
    else
        break;
}

/* reverse the order of jumps
 */
for (j = 0, i0--; j < i0; j++, i0--) {
    i = pp[0].n[j]; pp[0].n[j] = pp[0].n[i0]; pp[0].n[i0] = i;
    i = pp[0].x[j]; pp[0].x[j] = pp[0].x[i0]; pp[0].x[i0] = i;
}
for (j = 0, i1--; j < i1; j++, i1--) {
    i = pp[1].n[j]; pp[1].n[j] = pp[1].n[i1]; pp[1].n[i1] = i;
    i = pp[1].x[j]; pp[1].x[j] = pp[1].x[i1]; pp[1].x[i1] = i;
}
if (fd >= 0)
    (void) close(fd);
if (fj) {
    (void) unlink(jname);
    fj = 0; offset = 0; }

```

```
/*
 * write a filled jmp struct offset of the prev one (if any): nw()
 */
writejumps(ix)
    int ix;
{
    char      *mktemp();

    if (!fj) {
        if (mktemp(jname) < 0) {
            fprintf(stderr, "%s: can't mktemp() %s\n", prog, jname);
            cleanup(1);
        }
        if ((fj = fopen(jname, "w")) == 0) {
            fprintf(stderr, "%s: can't write %s\n", prog, jname);
            exit(1);
        }
    }
    (void) fwrite((char *)&dx[ix].jp, sizeof(struct jmp), 1, fj);
    (void) fwrite((char *)&dx[ix].offset, sizeof(dx[ix].offset), 1, fj);
}
```

**writejumps**

Table 2A

PRO	XXXXXXXXXXXXXXXXXX	(Length = 15 amino acids)
Comparison Protein	XXXXXXXXYYYYYYY	(Length = 12 amino acids)

% amino acid sequence identity =

(the number of identically matching amino acid residues between the two polypeptide sequences as determined by ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide) =

5 divided by 15 = 33.3%

Table 2B

PRO	XXXXXXXXXX	(Length = 10 amino acids)
Comparison Protein	XXXXXXXXYYYZZYZ	(Length = 15 amino acids)

% amino acid sequence identity =

(the number of identically matching amino acid residues between the two polypeptide sequences as determined by ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide) =

5 divided by 10 = 50%

Table 2C

PRO-DNA	NNNNNNNNNNNNNN	(Length = 14 nucleotides)
Comparison DNA	NNNNNNLLLLLLLLLL	(Length = 16 nucleotides)

% nucleic acid sequence identity =

(the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence) =

6 divided by 14 = 42.9%

Table 2D

PRO-DNA	NNNNNNNNNNNN	(Length = 12 nucleotides)
Comparison DNA	NNNNLLL	(Length = 9 nucleotides)

% nucleic acid sequence identity =

(the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence) =

4 divided by 12 = 33.3%

"Percent (%) amino acid sequence identity" with respect to the PRO polypeptide sequences identified herein is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in a PRO sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN, ALIGN-2 or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full-length of the sequences being compared. For purposes herein, however, % amino acid sequence identity values are obtained as described below by using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Table 1. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc., and the source code shown in Table 1 has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, California or may be compiled from the source code provided in Table 1. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

For purposes herein, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

$$100 \text{ times the fraction } X/Y$$

where X is the number of amino acid residues scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A. As examples of % amino acid sequence identity calculations, Tables 2A-2B demonstrate how to calculate the % amino acid sequence identity of the amino acid sequence designated "Comparison Protein" to the amino acid sequence designated "PRO".

Unless specifically stated otherwise, all % amino acid sequence identity values used herein are obtained as described above using the ALIGN-2 sequence comparison computer program. However, % amino acid sequence identity may also be determined using the sequence comparison program NCBI-BLAST2 (Altschul *et al.*, Nucleic Acids Res., 25:3389-3402 (1997)). The NCBI-BLAST2 sequence comparison program may be downloaded from <http://www.ncbi.nlm.nih.gov>. NCBI-BLAST2 uses several search parameters, wherein all of those search parameters are set to default values including, for example, unmask = yes, strand = all, expected occurrences = 10,



minimum low complexity length = 15/5, multi-pass e-value = 0.01, constant for multi-pass = 25, dropoff for final gapped alignment = 25 and scoring matrix = BLOSUM62.

In situations where NCBI-BLAST2 is employed for amino acid sequence comparisons, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

$$100 \text{ times the fraction } X/Y$$

where X is the number of amino acid residues scored as identical matches by the sequence alignment program NCBI-BLAST2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A.

In addition, % amino acid sequence identity may also be determined using the WU-BLAST-2 computer program (Altschul *et al.*, Methods in Enzymology, 266:460-480 (1996)). Most of the WU-BLAST-2 search parameters are set to the default values. Those not set to default values, *i.e.*, the adjustable parameters, are set with the following values: overlap span = 1, overlap fraction = 0.125, word threshold (T) = 11, and scoring matrix = BLOSUM62. For purposes herein, a % amino acid sequence identity value is determined by dividing (a) the number of matching identical amino acids residues between the amino acid sequence of the PRO polypeptide of interest having a sequence derived from the native PRO polypeptide and the comparison amino acid sequence of interest (*i.e.*, the sequence against which the PRO polypeptide of interest is being compared which may be a PRO variant polypeptide) as determined by WU-BLAST-2 by (b) the total number of amino acid residues of the PRO polypeptide of interest. For example, in the statement "a polypeptide comprising an amino acid sequence A which has or having at least 80% amino acid sequence identity to the amino acid sequence B", the amino acid sequence A is the comparison amino acid sequence of interest and the amino acid sequence B is the amino acid sequence of the PRO polypeptide of interest.

"PRO variant polynucleotide" or "PRO variant nucleic acid sequence" means a nucleic acid molecule which encodes an active PRO polypeptide as defined below and which has at least about 80% nucleic acid sequence identity with a nucleic acid sequence encoding a full-length native sequence PRO polypeptide sequence as disclosed herein, a full-length native sequence PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Ordinarily, a PRO variant polynucleotide will have at least about 80% nucleic acid sequence identity, more preferably at least about 81% nucleic acid sequence identity, more preferably at least about 82% nucleic acid sequence identity, more preferably at least about 83% nucleic acid sequence identity, more preferably at least about 84% nucleic acid sequence identity, more preferably at least about 85% nucleic acid sequence identity, more preferably at least about 86% nucleic acid

sequence identity, more preferably at least about 87% nucleic acid sequence identity, more preferably at least about 88% nucleic acid sequence identity, more preferably at least about 89% nucleic acid sequence identity, more preferably at least about 90% nucleic acid sequence identity, more preferably at least about 91% nucleic acid sequence identity, more preferably at least about 92% nucleic acid sequence identity, more preferably at least about 93% nucleic acid sequence identity, more preferably at least about 94% nucleic acid sequence identity, more preferably at least about 95% nucleic acid sequence identity, more preferably at least about 96% nucleic acid sequence identity, more preferably at least about 97% nucleic acid sequence identity, more preferably at least about 98% nucleic acid sequence identity and yet more preferably at least about 99% nucleic acid sequence identity with a nucleic acid sequence encoding a full-length native sequence PRO polypeptide sequence as disclosed herein, a full-length native sequence PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal sequence, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Variants do not encompass the native nucleotide sequence.

Ordinarily, PRO variant polynucleotides are at least about 30 nucleotides in length, often at least about 60 nucleotides in length, more often at least about 90 nucleotides in length, more often at least about 120 nucleotides in length, more often at least about 150 nucleotides in length, more often at least about 180 nucleotides in length, more often at least about 210 nucleotides in length, more often at least about 240 nucleotides in length, more often at least about 270 nucleotides in length, more often at least about 300 nucleotides in length, more often at least about 450 nucleotides in length, more often at least about 600 nucleotides in length, more often at least about 900 nucleotides in length, or more.

"Percent (%) nucleic acid sequence identity" with respect to the PRO polypeptide-encoding nucleic acid sequences identified herein is defined as the percentage of nucleotides in a candidate sequence that are identical with the nucleotides in a PRO polypeptide-encoding nucleic acid sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity. Alignment for purposes of determining percent nucleic acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN, ALIGN-2 or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full-length of the sequences being compared. For purposes herein, however, % nucleic acid sequence identity values are obtained as described below by using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Table 1. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc., and the source code shown in Table 1 has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, California or may be compiled from the source code provided in Table 1. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

For purposes herein, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence D (which can alternatively be phrased as a given nucleic acid sequence C that has or comprises a certain % nucleic acid sequence identity to, with, or against a given nucleic acid sequence D) is calculated as follows:

5

100 times the fraction  $W/Z$

where W is the number of nucleotides scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C is not equal to the length of nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C. As examples of %  
 10 nucleic acid sequence identity calculations, Tables 2C-2D demonstrate how to calculate the % nucleic acid sequence identity of the nucleic acid sequence designated "Comparison DNA" to the nucleic acid sequence designated "PRO-DNA".

Unless specifically stated otherwise, all % nucleic acid sequence identity values used herein are obtained as described above using the ALIGN-2 sequence comparison computer program. However, % nucleic acid sequence  
 15 identity may also be determined using the sequence comparison program NCBI-BLAST2 (Altschul *et al.*, Nucleic Acids Res., 25:3389-3402 (1997)). The NCBI-BLAST2 sequence comparison program may be downloaded from <http://www.ncbi.nlm.nih.gov>. NCBI-BLAST2 uses several search parameters, wherein all of those search parameters are set to default values including, for example, unmask = yes, strand = all, expected occurrences = 10, minimum low complexity length = 15/5, multi-pass e-value = 0.01, constant for multi-pass = 25, dropoff for final  
 20 gapped alignment = 25 and scoring matrix = BLOSUM62.

In situations where NCBI-BLAST2 is employed for sequence comparisons, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence D (which can alternatively be phrased as a given nucleic acid sequence C that has or comprises a certain % nucleic acid sequence identity to, with, or against a given nucleic acid sequence D) is calculated as follows:

25

100 times the fraction  $W/Z$

where W is the number of nucleotides scored as identical matches by the sequence alignment program NCBI-BLAST2 in that program's alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C is not equal to the length of nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C.

30

In addition, % nucleic acid sequence identity values may also be generated using the WU-BLAST-2 computer program (Altschul *et al.*, Methods in Enzymology, 266:460-480 (1996)). Most of the WU-BLAST-2 search parameters are set to the default values. Those not set to default values, *i.e.*, the adjustable parameters, are set with the following values: overlap span = 1, overlap fraction = 0.125, word threshold (T) = 11, and scoring

matrix = BLOSUM62. For purposes herein, a % nucleic acid sequence identity value is determined by dividing (a) the number of matching identical nucleotides between the nucleic acid sequence of the PRO polypeptide-encoding nucleic acid molecule of interest having a sequence derived from the native sequence PRO polypeptide-encoding nucleic acid and the comparison nucleic acid molecule of interest (*i.e.*, the sequence against which the PRO polypeptide-encoding nucleic acid molecule of interest is being compared which may be a variant PRO polynucleotide) as determined by WU-BLAST-2 by (b) the total number of nucleotides of the PRO polypeptide-encoding nucleic acid molecule of interest. For example, in the statement "an isolated nucleic acid molecule comprising a nucleic acid sequence A which has or having at least 80% nucleic acid sequence identity to the nucleic acid sequence B", the nucleic acid sequence A is the comparison nucleic acid molecule of interest and the nucleic acid sequence B is the nucleic acid sequence of the PRO polypeptide-encoding nucleic acid molecule of interest.

In other embodiments, PRO variant polynucleotides are nucleic acid molecules that encode an active PRO polypeptide and which are capable of hybridizing, preferably under stringent hybridization and wash conditions, to nucleotide sequences encoding the full-length PRO polypeptide shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:9), Figure 6 (SEQ ID NO:11), Figure 8 (SEQ ID NO:16), Figure 10 (SEQ ID NO:21), Figure 12 (SEQ ID NO:26), Figure 14 (SEQ ID NO:31), Figure 16 (SEQ ID NO:36), Figure 18 (SEQ ID NO:41), Figure 20 (SEQ ID NO:46), Figure 22 (SEQ ID NO:51), Figure 24 (SEQ ID NO:57), Figure 26 (SEQ ID NO:62), Figure 28 (SEQ ID NO:67), Figure 30 (SEQ ID NO:72), Figure 32 (SEQ ID NO:77), Figure 34 (SEQ ID NO:82), Figure 36 (SEQ ID NO:91), Figure 38 (SEQ ID NO:96), Figure 40 (SEQ ID NO:101), Figure 42 (SEQ ID NO:108), Figure 44 (SEQ ID NO:113), Figure 46 (SEQ ID NO:119), Figure 48 (SEQ ID NO:127), Figure 50 (SEQ ID NO:132), Figure 52 (SEQ ID NO:137), Figure 54 (SEQ ID NO:142), Figure 56 (SEQ ID NO:152), Figure 58 (SEQ ID NO:154), Figure 60 (SEQ ID NO:156), Figure 62 (SEQ ID NO:158), Figure 64 (SEQ ID NO:160), Figure 66 (SEQ ID NO:165), and Figure 68 (SEQ ID NO:167), respectively. PRO variant polypeptides may be those that are encoded by a PRO variant polynucleotide.

The term "positives", in the context of the amino acid sequence identity comparisons performed as described above, includes amino acid residues in the sequences compared that are not only identical, but also those that have similar properties. Amino acid residues that score a positive value to an amino acid residue of interest are those that are either identical to the amino acid residue of interest or are a preferred substitution (as defined in Table 3 below) of the amino acid residue of interest.

For purposes herein, the % value of positives of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % positives to, with, or against a given amino acid sequence B) is calculated as follows:

$$100 \text{ times the fraction } X/Y$$

where X is the number of amino acid residues scoring a positive value by the sequence alignment program ALIGN-2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B,

the % positives of A to B will not equal the % positives of B to A.

- “Isolated”, when used to describe the various polypeptides disclosed herein, means a polypeptide that has been identified and separated and/or recovered from a component of its natural environment. Preferably, the isolated polypeptide is free of association with all components with which it is naturally associated. Contaminant components of its natural environment are materials that would typically interfere with diagnostic or therapeutic uses for the polypeptide, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In preferred embodiments, the polypeptide will be purified (1) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (2) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue or, preferably, silver stain.
- 10 Isolated polypeptide includes polypeptide *in situ* within recombinant cells, since at least one component of the PRO natural environment will not be present. Ordinarily, however, isolated polypeptide will be prepared by at least one purification step.

- An “isolated” nucleic acid molecule encoding a PRO polypeptide or an “isolated” nucleic acid molecule encoding an anti-PRO antibody is a nucleic acid molecule that is identified and separated from at least one contaminant nucleic acid molecule with which it is ordinarily associated in the natural source of the PRO-encoding nucleic acid or the natural source of the anti-PRO-encoding nucleic acid. Preferably, the isolated nucleic acid is free of association with all components with which it is naturally associated. An isolated PRO-encoding nucleic acid molecule or an isolated anti-PRO-encoding nucleic acid molecule is other than in the form or setting in which it is found in nature. Isolated nucleic acid molecules therefore are distinguished from the PRO-encoding nucleic acid molecule or from the anti-PRO-encoding nucleic acid molecule as it exists in natural cells. However, an isolated nucleic acid molecule encoding a PRO polypeptide or an isolated nucleic acid molecule encoding an anti-PRO antibody includes PRO-nucleic acid molecules or anti-PRO-nucleic acid molecules contained in cells that ordinarily express PRO polypeptides or anti-PRO antibodies where, for example, the nucleic acid molecule is in a chromosomal location different from that of natural cells.

- 25 The term “control sequences” refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

- Nucleic acid is “operably linked” when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a PRO polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, “operably linked” means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

"Stringency" of hybridization reactions is readily determinable by one of ordinary skill in the art, and generally is an empirical calculation dependent upon probe length, washing temperature, and salt concentration. In general, longer probes require higher temperatures for proper annealing, while shorter probes need lower temperatures. Hybridization generally depends on the ability of denatured DNA to reanneal when complementary strands are present in an environment below their melting temperature. The higher the degree of desired homology between the probe and hybridizable sequence, the higher the relative temperature that can be used. As a result, it follows that higher relative temperatures would tend to make the reaction conditions more stringent, while lower temperatures less so. For additional details and explanation of stringency of hybridization reactions, *see*, Ausubel *et al.*, Current Protocols in Molecular Biology (Wiley Interscience Publishers, 1995).

"Stringent conditions" or "high-stringency conditions", as defined herein, may be identified by those that: (1) employ low ionic strength and high temperature for washing, for example, 0.015 M sodium chloride/0.0015 M sodium citrate/0.1% sodium dodecyl sulfate at 50°C; (2) employ during hybridization a denaturing agent, such as formamide, for example, 50% (v/v) formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50mM sodium phosphate buffer at pH 6.5 with 750 mM sodium chloride, 75 mM sodium citrate at 42°C; or (3) employ 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42°C, with washes at 42°C in 0.2 x SSC (sodium chloride/sodium citrate) and 50% formamide at 55°C, followed by a high-stringency wash consisting of 0.1 x SSC containing EDTA at 55°C.

"Moderately-stringent conditions" may be identified as described by Sambrook *et al.*, Molecular Cloning: A Laboratory Manual (New York: Cold Spring Harbor Press, 1989), and include the use of washing solution and hybridization conditions (*e.g.*, temperature, ionic strength, and % SDS) less stringent than those described above. An example of moderately stringent conditions is overnight incubation at 37°C in a solution comprising: 20% formamide, 5 x SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5 x Denhardt's solution, 10% dextran sulfate, and 20 mg/ml denatured sheared salmon sperm DNA, followed by washing the filters in 1 x SSC at about 37-50°C. The skilled artisan will recognize how to adjust the temperature, ionic strength, etc. as necessary to accommodate factors such as probe length and the like.

The modifier "epitope-tagged" when used herein refers to a chimeric polypeptide comprising a PRO polypeptide fused to a "tag polypeptide". The tag polypeptide has enough residues to provide an epitope against which an antibody can be made, yet is short enough such that it does not interfere with activity of the polypeptide to which it is fused. The tag polypeptide preferably also is fairly unique so that the antibody does not substantially cross-react with other epitopes. Suitable tag polypeptides generally have at least six amino acid residues and usually between about 8 and 50 amino acid residues (preferably, between about 10 and 20 amino acid residues).

"Active" or "activity" in the context of PRO variants refers to form(s) of PRO proteins that retain the biologic and/or immunologic activities of a native or naturally-occurring PRO polypeptide.

"Biological activity" in the context of a molecule that antagonizes a PRO polypeptide that can be identified by the screening assays disclosed herein (*e.g.*, an organic or inorganic small molecule, peptide, etc.) is used to refer

to the ability of such molecules to bind or complex with the PRO polypeptide identified herein, or otherwise interfere with the interaction of the PRO polypeptides with other cellular proteins or otherwise inhibits the transcription or translation of the PRO polypeptide. Particularly preferred biological activity includes cardiac hypertrophy, activity that acts on systemic disorders that affect vessels, such as diabetes mellitus, as well as diseases of the arteries, capillaries, veins, and/or lymphatics, and cancer.

The term "antagonist" is used in the broadest sense, and includes any molecule that partially or fully blocks, inhibits, or neutralizes one or more of the biological activities of a native PRO polypeptide disclosed herein, for example, if applicable, its mitogenic or angiogenic activity. Antagonists of a PRO polypeptide may act by interfering with the binding of a PRO polypeptide to a cellular receptor, by incapacitating or killing cells that have been activated by a PRO polypeptide, or by interfering with vascular endothelial cell activation after binding of a PRO polypeptide to a cellular receptor. All such points of intervention by a PRO polypeptide antagonist shall be considered equivalent for purposes of this invention. The antagonists inhibit the mitogenic, angiogenic, or other biological activity of PRO polypeptides, and thus are useful for the treatment of diseases or disorders characterized by undesirable excessive neovascularization, including by way of example tumors, and especially solid malignant tumors, rheumatoid arthritis, psoriasis, atherosclerosis, diabetic and other retinopathies, retrolental fibroplasia, age-related macular degeneration, neovascular glaucoma, hemangiomas, thyroid hyperplasias (including Grave's disease), corneal and other tissue transplantation, and chronic inflammation. The antagonists also are useful for the treatment of diseases or disorders characterized by undesirable excessive vascular permeability, such as edema associated with brain tumors, ascites associated with malignancies, Meigs' syndrome, lung inflammation, nephrotic syndrome, pericardial effusion (such as that associated with pericarditis), and pleural effusion. In a similar manner, the term "agonist" is used in the broadest sense and includes any molecule that mimics a biological activity of a native PRO polypeptide disclosed herein. Suitable agonist or antagonist molecules specifically include agonist or antagonist antibodies or antibody fragments, fragments, or amino acid sequence variants of native PRO polypeptides, peptides, small organic molecules, etc.

A "small molecule" is defined herein to have a molecular weight below about 500 daltons.

The term "PRO polypeptide receptor" as used herein refers to a cellular receptor for a PRO polypeptide, ordinarily a cell-surface receptor found on vascular endothelial cells, as well as variants thereof that retain the ability to bind a PRO polypeptide.

"Antibodies" (Abs) and "immunoglobulins" (Igs) are glycoproteins having the same structural characteristics. While antibodies exhibit binding specificity to a specific antigen, immunoglobulins include both antibodies and other antibody-like molecules that lack antigen specificity. Polypeptides of the latter kind are, for example, produced at low levels by the lymph system and at increased levels by myelomas. The term "antibody" is used in the broadest sense and specifically covers, without limitation, intact monoclonal antibodies, polyclonal antibodies, multispecific antibodies (e.g., bispecific antibodies) formed from at least two intact antibodies, and antibody fragments, so long as they exhibit the desired biological activity.

"Native antibodies" and "native immunoglobulins" are usually heterotetrameric glycoproteins of about 150,000 daltons, composed of two identical light (L) chains and two identical heavy (H) chains. Each light chain is linked

to a heavy chain by one covalent disulfide bond, while the number of disulfide linkages varies among the heavy chains of different immunoglobulin isotypes. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at one end a variable domain ( $V_H$ ) followed by a number of constant domains. Each light chain has a variable domain at one end ( $V_L$ ) and a constant domain at its other end; the constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light-chain variable domain is aligned with the variable domain of the heavy chain. Particular amino acid residues are believed to form an interface between the light- and heavy-chain variable domains.

The term "variable" refers to the fact that certain portions of the variable domains differ extensively in sequence among antibodies and are used in the binding and specificity of each particular antibody to and for its particular antigen. However, the variability is not evenly distributed throughout the variable domains of antibodies. It is concentrated in three segments called complementarity-determining regions (CDRs) or hypervariable regions both in the light-chain and the heavy-chain variable domains. The more highly conserved portions of variable domains are called the framework regions (FR). The variable domains of native heavy and light chains each comprise four FR regions, largely adopting a  $\beta$ -sheet configuration, connected by three CDRs, which form loops connecting, and in some cases forming part of, the  $\beta$ -sheet structure. The CDRs in each chain are held together in close proximity by the FR regions and, with the CDRs from the other chain, contribute to the formation of the antigen-binding site of antibodies. See, Kabat *et al.*, NIH Publ. No.91-3242, Vol. I, pages 647-669 (1991). The constant domains are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody-dependent cellular toxicity.

"Antibody fragments" comprise a portion of an intact antibody, preferably the antigen-binding or variable region of the intact antibody. Examples of antibody fragments include Fab, Fab', F(ab')<sub>2</sub>, and Fv fragments; diabodies; linear antibodies (Zapata *et al.*, Protein Eng., 8(10): 1057-1062 (1995)); single-chain antibody molecules; and multispecific antibodies formed from antibody fragments.

Papain digestion of antibodies produces two identical antigen-binding fragments, called "Fab" fragments, each with a single antigen-binding site, and a residual "Fc" fragment, whose name reflects its ability to crystallize readily. Pepsin treatment yields an F(ab')<sub>2</sub> fragment that has two antigen-combining sites and is still capable of cross-linking antigen.

"Fv" is the minimum antibody fragment that contains a complete antigen-recognition and -binding site. This region consists of a dimer of one heavy- and one light-chain variable domain in tight, non-covalent association. It is in this configuration that the three CDRs of each variable domain interact to define an antigen-binding site on the surface of the  $V_H$ - $V_L$  dimer. Collectively, the six CDRs confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

The Fab fragment also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. Fab' fragments differ from Fab fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free thiol group.



F(ab')<sub>2</sub> antibody fragments originally were produced as pairs of Fab' fragments that have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

The "light chains" of antibodies (immunoglobulins) from any vertebrate species can be assigned to one of two clearly distinct types, called kappa (κ) and lambda (λ), based on the amino acid sequences of their constant domains.

5 Depending on the amino acid sequence of the constant domain of their heavy chains, immunoglobulins can be assigned to different classes. There are five major classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM; and several of these may be further divided into subclasses (isotypes), *e.g.*, IgG1, IgG2, IgG3, IgG4, IgA, and IgA2. The heavy-chain constant domains that correspond to the different classes of immunoglobulins are called α, δ, ε, γ, and μ, respectively. The subunit structures and three-dimensional configurations of different classes of  
10 immunoglobulins are well known.

The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally-occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to conventional (polyclonal)  
15 antibody preparations that typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. In addition to their specificity, the monoclonal antibodies are advantageous in that they are synthesized by the hybridoma culture, uncontaminated by other immunoglobulins. The modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of  
20 the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler *et al.*, Nature, 256: 495 (1975), or may be made by recombinant DNA methods (*see, e.g.*, U.S. Patent No. 4,816,567). The "monoclonal antibodies" may also be isolated from phage antibody libraries using the techniques described in Clackson *et al.*, Nature, 352: 624-628 (1991) and Marks *et al.*, J. Mol. Biol., 222: 581-597 (1991), for example.

25 The monoclonal antibodies herein specifically include "chimeric" antibodies (immunoglobulins) in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the  
30 desired biological activity. U.S. Patent No. 4,816,567; Morrison *et al.*, Proc. Natl. Acad. Sci. USA, 81: 6851-6855 (1984).

"Humanized" forms of non-human (*e.g.*, murine) antibodies are chimeric immunoglobulins, immunoglobulin chains, or fragments thereof (such as Fv, Fab, Fab', F(ab')<sub>2</sub>, or other antigen-binding subsequences of antibodies) that contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies  
35 are human immunoglobulins (recipient antibody) in which residues from a CDR of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity, and capacity. In some instances, Fv FR residues of the human immunoglobulin are replaced

by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues that are found neither in the recipient antibody nor in the imported CDR or framework sequences. These modifications are made to further refine and maximize antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin sequence. The humanized antibody preferably also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, *see* Jones *et al.*, Nature, **321**: 522-525 (1986); Reichmann *et al.*, Nature, **332**: 323-329 (1988); and Presta, Curr. Op. Struct. Biol., **2**: 593-596 (1992). The humanized antibody includes a PRIMATIZED™ antibody wherein the antigen-binding region of the antibody is derived from an antibody produced by immunizing macaque monkeys with the antigen of interest.

"Single-chain Fv" or "sFv" antibody fragments comprise the V<sub>H</sub> and V<sub>L</sub> domains of an antibody, wherein these domains are present in a single polypeptide chain. Preferably, the Fv polypeptide further comprises a polypeptide linker between the V<sub>H</sub> and V<sub>L</sub> domains that enables the sFv to form the desired structure for antigen binding. For a review of sFv *see*, Pluckthun in The Pharmacology of Monoclonal Antibodies, Vol. 113, Rosenberg and Moore, eds. (Springer-Verlag: New York, 1994), pp. 269-315.

The term "diabodies" refers to small antibody fragments with two antigen-binding sites, which fragments comprise a heavy-chain variable domain (V<sub>H</sub>) connected to a light-chain variable domain (V<sub>L</sub>) in the same polypeptide chain (V<sub>H</sub> - V<sub>L</sub>). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies are described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger *et al.*, Proc. Natl. Acad. Sci. USA, **90**: 6444-6448 (1993).

An "isolated" antibody is one that has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials that would interfere with diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In preferred embodiments, the antibody will be purified (1) to greater than 95% by weight of antibody as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions using Coomassie blue or, preferably, silver stain. Isolated antibody includes the antibody *in situ* within recombinant cells, since at least one component of the antibody's natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

The word "label" when used herein refers to a detectable compound or other composition that is conjugated directly or indirectly to the antibody so as to generate a "labeled" antibody. The label may be detectable by itself (*e.g.*, radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, may catalyze chemical alteration of a substrate compound or composition that is detectable. Radionuclides that can serve as detectable labels include, for example, I-131, I-123, I-125, Y-90, Re-188, At-211, Cu-67, Bi-212, and Pd-109. The label may also be a non-

detectable entity such as a toxin.

By "solid phase" is meant a non-aqueous matrix to which an antibody of the present invention can adhere. Examples of solid phases encompassed herein include those formed partially or entirely of glass (*e.g.*, controlled pore glass), polysaccharides (*e.g.*, agarose), polyacrylamides, polystyrene, polyvinyl alcohol and silicones. In certain embodiments, depending on the context, the solid phase can comprise the well of an assay plate; in others it is a purification column (*e.g.*, an affinity chromatography column). This term also includes a discontinuous solid phase of discrete particles, such as those described in U.S. Patent No. 4,275,149.

A "liposome" is a small vesicle composed of various types of lipids, phospholipids and/or surfactant that is useful for delivery of a drug (such as the PRO polypeptide or antibodies thereto disclosed herein) to a mammal.

The components of the liposome are commonly arranged in a bilayer formation, similar to the lipid arrangement of biological membranes.

As used herein, the term "immunoadhesin" designates antibody-like molecules that combine the binding specificity of a heterologous protein (an "adhesin") with the effector functions of immunoglobulin constant domains. Structurally, the immunoadhesins comprise a fusion of an amino acid sequence with the desired binding specificity that is other than the antigen recognition and binding site of an antibody (*i.e.*, is "heterologous"), and an immunoglobulin constant domain sequence. The adhesin part of an immunoadhesin molecule typically is a contiguous amino acid sequence comprising at least the binding site of a receptor or a ligand. The immunoglobulin constant domain sequence in the immunoadhesin may be obtained from any immunoglobulin, such as IgG-1, IgG-2, IgG-3, or IgG-4 subtypes, IgA (including IgA-1 and IgA-2), IgE, IgD, or IgM.

## II. Compositions and Methods of the Invention

A. PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 and PRO1246 Variants

In addition to the full-length native sequence PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 and PRO1246 polypeptides described herein, it is contemplated that PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 and PRO1246 variants can be prepared. PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 and PRO1246 variants can be prepared by introducing appropriate nucleotide changes into the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195,

PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 DNA, and/or by synthesis of the desired PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide. Those skilled in the art will appreciate that amino acid changes may alter post-translational processes of the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246, such as changing the number or position of glycosylation sites or altering the membrane anchoring characteristics.

Variations in the native full-length sequence PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 or in various domains of the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 described herein, can be made, for example, using any of the techniques and guidelines for conservative and non-conservative mutations set forth, for instance, in U.S. Patent No. 5,364,934. Variations may be a substitution, deletion or insertion of one or more codons encoding the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 that results in a change in the amino acid sequence of the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 as compared with the native sequence PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246. Optionally the variation is by substitution of at least one amino acid with any other amino acid in one or more of the domains of the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246. Guidance in determining which amino acid residue may be inserted, substituted or deleted without adversely affecting the desired activity may be found by comparing the sequence of the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179,

PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 with that of homologous known protein molecules and minimizing the number of amino acid sequence changes made in regions of high homology. Amino acid  
5 substitutions can be the result of replacing one amino acid with another amino acid having similar structural and/or chemical properties, such as the replacement of a leucine with a serine, *i.e.*, conservative amino acid replacements. Insertions or deletions may optionally be in the range of about 1 to 5 amino acids. The variation allowed may be determined by systematically making insertions, deletions or substitutions of amino acids in the sequence and testing the resulting variants for activity exhibited by the full-length or mature native sequence.

- 10 In particular embodiments, conservative substitutions of interest are shown in Table 3 under the heading of preferred substitutions. If such substitutions result in a change in biological activity, then more substantial changes, denominated exemplary substitutions in Table 3, or as further described below in reference to amino acid classes, are introduced and the products screened.

Table 3

	Original	Exemplary	Preferred
	<u>Residue</u>	<u>Substitutions</u>	<u>Substitutions</u>
	Ala (A)	val; leu; ile	val
5	Arg (R)	lys; gln; asn	lys
	Asn (N)	gln; his; lys; arg	gln
	Asp (D)	glu	glu
	Cys (C)	ser	ser
	Gln (Q)	asn	asn
10	Glu (E)	asp	asp
	Gly (G)	pro; ala	ala
	His (H)	asn; gln; lys; arg	arg
	Ile (I)	leu; val; met; ala; phe; norleucine	leu
15	Leu (L)	norleucine; ile; val; met; ala; phe	ile
	Lys (K)	arg; gln; asn	arg
	Met (M)	leu; phe; ile	leu
	Phe (F)	leu; val; ile; ala; tyr	leu
20	Pro (P)	ala	ala
	Ser (S)	thr	thr
	Thr (T)	ser	ser
	Trp (W)	tyr; phe	tyr
	Tyr (Y)	trp; phe; thr; ser	phe
25	Val (V)	ile; leu; met; phe; ala; norleucine	leu

- Substantial modifications in function or immunological identity of the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side-chain properties:
- 30 (1) hydrophobic: norleucine, met, ala, val, leu, ile;  
 (2) neutral hydrophilic: cys, ser, thr;  
 (3) acidic: asp, glu;  
 (4) basic: asn, gln, his, lys, arg;  
 (5) residues that influence chain orientation: gly, pro; and  
 40 (6) aromatic: trp, tyr, phe.

Non-conservative substitutions will entail exchanging a member of one of these classes for another class. Such substituted residues also may be introduced into the conservative substitution sites or, more preferably, into the remaining (non-conserved) sites.

The variations can be made using methods known in the art such as oligonucleotide-mediated (site-directed)

mutagenesis, alanine scanning, and PCR mutagenesis. Site-directed mutagenesis [Carter *et al.*, Nucl. Acids Res., 13:4331 (1986); Zoller *et al.*, Nucl. Acids Res., 10:6487 (1987)], cassette mutagenesis [Wells *et al.*, Gene, 34:315 (1985)], restriction selection mutagenesis [Wells *et al.*, Philos. Trans. R. Soc. London SerA, 317:415 (1986)] or other known techniques can be performed on the cloned DNA to produce the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 variant DNA.

Scanning amino acid analysis can also be employed to identify one or more amino acids along a contiguous sequence. Among the preferred scanning amino acids are relatively small, neutral amino acids. Such amino acids include alanine, glycine, serine, and cysteine. Alanine is typically a preferred scanning amino acid among this group because it eliminates the side-chain beyond the beta-carbon and is less likely to alter the main-chain conformation of the variant [Cunningham and Wells, Science, 244: 1081-1085 (1989)]. Alanine is also typically preferred because it is the most common amino acid. Further, it is frequently found in both buried and exposed positions [Creighton, The Proteins, (W.H. Freeman & Co., N.Y.); Chothia, J. Mol. Biol., 150:1 (1976)]. If alanine substitution does not yield adequate amounts of variant, an isoteric amino acid can be used.

B. Modifications of PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 and PRO1246

Covalent modifications of PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 and PRO1246 are included within the scope of this invention. One type of covalent modification includes reacting targeted amino acid residues of a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C- terminal residues of the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246. Derivatization with bifunctional agents is useful, for instance, for crosslinking PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 to a water-insoluble support matrix or surface for use in the method for purifying anti-PRO172, anti-PRO175, anti-PRO178, anti-

PRO188, anti-PRO356, anti-PRO179, anti-PRO197, anti-PRO198, anti-PRO182, anti-PRO195, anti-PRO200, anti-PRO211, anti-PRO217, anti-PRO219, anti-PRO221, anti-PRO224, anti-PRO228, anti-PRO245, anti-PRO246, anti-PRO258, anti-PRO261, anti-PRO272, anti-PRO301, anti-PRO322, anti-PRO328, anti-PRO331, anti-PRO364, anti-PRO366, anti-PRO535, anti-PRO819, anti-PRO826, anti-PRO1160, anti-PRO1186 or anti-PRO1246 antibodies, and vice-versa. Commonly used crosslinking agents include, e.g., 1,1-bis(diazoacetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis(succinimidylpropionate), bifunctional maleimides such as bis-N-maleimido-1,8-octane and agents such as methyl-3-[(p-azidophenyl)dithio]propioimide.

Other modifications include deamidation of glutaminyl and asparaginyl residues to the corresponding glutamyl and aspartyl residues, respectively, hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the  $\alpha$ -amino groups of lysine, arginine, and histidine side chains [T.E. Creighton, Proteins: Structure and Molecular Properties, W.H. Freeman & Co., San Francisco, pp. 79-86 (1983)], acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

Another type of covalent modification of the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide included within the scope of this invention comprises altering the native glycosylation pattern of the polypeptide. "Altering the native glycosylation pattern" is intended for purposes herein to mean deleting one or more carbohydrate moieties found in native sequence PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 (either by removing the underlying glycosylation site or by deleting the glycosylation by chemical and/or enzymatic means), and/or adding one or more glycosylation sites that are not present in the native sequence PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246. In addition, the phrase includes qualitative changes in the glycosylation of the native proteins, involving a change in the nature and proportions of the various carbohydrate moieties present.

Addition of glycosylation sites to the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide may be accomplished by altering the amino acid sequence. The alteration may be made, for example, by the addition of, or substitution by, one or more serine or threonine residues to the native sequence PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535,



PRO819, PRO826, PRO1160, PRO1186 or PRO1246 (for O-linked glycosylation sites). The PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 amino acid  
 5 sequence may optionally be altered through changes at the DNA level, particularly by mutating the DNA encoding the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide at preselected bases such that codons are generated that will translate into the desired amino  
 10 acids.

Another means of increasing the number of carbohydrate moieties on the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide is by  
 15 chemical or enzymatic coupling of glycosides to the polypeptide. Such methods are described in the art, *e.g.*, in WO 87/05330 published 11 September 1987, and in Aplin and Wriston, CRC Crit. Rev. Biochem., pp. 259-306 (1981).

Removal of carbohydrate moieties present on the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366,  
 20 PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide may be accomplished chemically or enzymatically or by mutational substitution of codons encoding for amino acid residues that serve as targets for glycosylation. Chemical deglycosylation techniques are known in the art and described, for instance, by Hakimuddin, *et al.*, Arch. Biochem. Biophys., 259:52 (1987) and by Edge *et al.*, Anal. Biochem., 118:131 (1981). Enzymatic cleavage of carbohydrate moieties on polypeptides can be achieved by the use of a variety of endo- and  
 25 exo-glycosidases as described by Thotakura *et al.*, Meth. Enzymol., 138:350 (1987).

Another type of covalent modification of PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 comprises linking the PRO172, PRO175, PRO178, PRO188,  
 30 PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide to one of a variety of nonproteinaceous polymers, *e.g.*, polyethylene glycol (PEG), polypropylene glycol, or polyoxyalkylenes, in the manner set forth in U.S. Patent Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337.

The PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160,

PRO1186 or PRO1246 of the present invention may also be modified in a way to form a chimeric molecule comprising PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160,

5 PRO1186 or PRO1246 fused to another, heterologous polypeptide or amino acid sequence.

In one embodiment, such a chimeric molecule comprises a fusion of the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 with a tag polypeptide which provides an epitope to which an anti-tag antibody can selectively bind. The epitope tag is generally placed at the amino- or carboxyl- terminus of the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246. The presence of such epitope-tagged forms of the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 can be detected using an antibody against the tag polypeptide. Also, provision of the epitope tag enables the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 to be readily purified by affinity purification using an anti-tag antibody or another type of affinity matrix that binds to the epitope tag. Various tag polypeptides and their respective antibodies are well known in the art. Examples include poly-histidine (poly-His) or poly-histidine-glycine (poly-His-gly) tags; the flu HA tag polypeptide and its antibody 12CA5 [Field *et al.*, Mol. Cell. Biol., 8:2159-2165 (1988)]; the c-myc tag and the 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies thereto [Evan *et al.*, Molecular and Cellular Biology, 5:3610-3616 (1985)]; and the Herpes Simplex virus glycoprotein D (gD) tag and its antibody [Paborsky *et al.*, Protein Engineering, 3(6):547-553 (1990)]. Other tag polypeptides include the Flag-peptide [Hopp *et al.*, BioTechnology, 6:1204-1210 (1988)]; the KT3 epitope peptide [Martin *et al.*, Science, 255:192-194 (1992)]; an  $\alpha$ -tubulin epitope peptide [Skinner *et al.*, J. Biol. Chem., 266:15163-15166 (1991)]; and the T7 gene 10 protein peptide tag [Lutz-Freyermuth *et al.*, Proc. Natl. Acad. Sci. USA, 87:6393-6397 (1990)].

In an alternative embodiment, the chimeric molecule may comprise a fusion of the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 with an immunoglobulin or a particular region of an immunoglobulin. For a bivalent form of the chimeric molecule (also referred to as an "immunoadhesin"), such a fusion could be to the Fc region of an IgG molecule. The Ig fusions

preferably include the substitution of a soluble (transmembrane domain deleted or inactivated) form of a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide in place of at least one variable region within an Ig molecule. In a particularly preferred embodiment, the immunoglobulin fusion includes the hinge, CH2 and CH3, or the hinge, CH1, CH2 and CH3 regions of an IgG1 molecule. For the production of immunoglobulin fusions *see* also, US Patent No. 5,428,130 issued June 27, 1995.

C. Preparation of the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 and PRO1246

The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246. In particular, cDNAs encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptides have been identified and isolated, as disclosed in further detail in the Examples below. It is noted that proteins produced in separate expression rounds may be given different PRO numbers but the UNQ number is unique for any given DNA and the encoded protein, and will not be changed. However, for sake of simplicity, in the present specification the protein encoded by DNA35916-1161, DNA19355-1150, DNA23339-1130, DNA28497-1130, DNA47470-1130-P1, DNA16451-1078, DNA22780-1078, DNA33457-1078, DNA27865-1091, DNA26847-1395, DNA29101-1122, DNA32292-1131, DNA33094-1131, DNA32290-1164, DNA33089-1132, DNA33221-1133, DNA33092-1202, DNA35638-1141, DNA35639-1172, DNA35918-1174, DNA33473-1176, DNA40620-1183, DNA40628-1216, DNA48336-1309, DNA40587-1231, DNA40981-1234, DNA47365-1206, DNA33085-1110, DNA49143-1429, DNA57695-1340, DNA57694-1341, DNA62872-1509, DNA60621-1516 or DNA64885-1549, as well as all further native homologues and variants included in the foregoing definition of PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246, will be referred to as "PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246", respectively, regardless of their origin or mode of preparation.

The description below relates primarily to production of PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptides by culturing cells transformed or transfected with a vector containing nucleic acid encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptides. It is, of course, contemplated that alternative methods that are well known in the art may be employed to prepare PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246. For instance, the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide sequence, or portions thereof, may be produced by direct peptide synthesis using solid-phase techniques. *See, e.g.,* Stewart *et al.*, Solid-Phase Peptide Synthesis (W.H. Freeman Co.: San Francisco, CA, 1969); Merrifield, *J. Am. Chem. Soc.*, **85**: 2149-2154 (1963). *In vitro* protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be accomplished, for instance, with an Applied Biosystems Peptide Synthesizer (Foster City, CA) using manufacturer's instructions. Various portions of PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 may be chemically synthesized separately and combined using chemical or enzymatic methods to produce the full-length PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide.

i. Isolation of DNA Encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246

DNA encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide may be obtained from a cDNA library prepared from tissue believed

to possess the mRNA encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 and to express it at a detectable level. Accordingly, DNAs encoding

5 human PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 can be conveniently obtained from cDNA libraries prepared from human tissues, such as described in the Examples. The gene encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198,

10 PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide may also be obtained from a genomic library or by oligonucleotide synthesis.

Libraries can be screened with probes (such as antibodies to the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221,

15 PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide or oligonucleotides of at least about 20-80 bases) designed to identify the gene of interest or the protein encoded by it. Screening the cDNA or genomic library with the selected probe may be conducted using standard procedures, such as described

20 in Sambrook *et al.*, *supra*. An alternative means to isolate the gene encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 is to use PCR methodology. Sambrook *et al.*, *supra*; Dieffenbach *et al.*, PCR Primer: A Laboratory Manual (New York: Cold Spring Harbor

25 Laboratory Press, 1995).

The Examples below describe techniques for screening a cDNA library. The oligonucleotide sequences selected as probes should be of sufficient length and sufficiently unambiguous that false positives are minimized. The oligonucleotide is preferably labeled such that it can be detected upon hybridization to DNA in the library being screened. Methods of labeling are well known in the art, and include the use of radiolabels like <sup>32</sup>P-labeled ATP,

30 biotinylation, or enzyme labeling. Hybridization conditions, including moderate stringency and high stringency, are provided in Sambrook *et al.*, *supra*.

Sequences identified in such library screening methods can be compared and aligned to other known sequences deposited and available in public databases such as GenBank or other private sequence databases. Sequence identity (at either the amino acid or nucleotide level) within defined regions of the molecule or across the

35 full-length sequence can be determined through sequence alignment using computer software programs such as ALIGN, DNASTar, and INHERIT, which employ various algorithms to measure homology.

Nucleic acid having protein coding sequence may be obtained by screening selected cDNA or genomic

libraries using the deduced amino acid sequence disclosed herein for the first time, and, if necessary, using conventional primer extension procedures as described in Sambrook *et al.*, *supra*, to detect precursors and processing intermediates of mRNA that may not have been reverse-transcribed into cDNA.

## ii. Selection and Transformation of Host Cells

- 5 Host cells are transfected or transformed with expression or cloning vectors described herein for PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting  
10 transformants, or amplifying the genes encoding the desired sequences. The culture conditions, such as media, temperature, pH, and the like, can be selected by the skilled artisan without undue experimentation. In general, principles, protocols, and practical techniques for maximizing the productivity of cell cultures can be found in Mammalian Cell Biotechnology: A Practical Approach, M. Butler, ed. (IRL Press, 1991) and Sambrook *et al.*, *supra*.
- 15 Methods of transfection are known to the ordinarily skilled artisan, for example, CaPO<sub>4</sub> treatment and electroporation. Depending on the host cell used, transformation is performed using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described in Sambrook *et al.*, *supra*, or electroporation is generally used for prokaryotes or other cells that contain substantial cell-wall barriers. Infection with *Agrobacterium tumefaciens* is used for transformation of certain plant cells, as described by Shaw  
20 *et al.*, Gene, **23**: 315 (1983) and WO 89/05859 published 29 June 1989. For mammalian cells without such cell walls, the calcium phosphate precipitation method of Graham and van der Eb, Virology, **52**:456-457 (1978) can be employed. General aspects of mammalian cell host system transformations have been described in U.S. Patent No. 4,399,216. Transformations into yeast are typically carried out according to the method of Van Solingen *et al.*, J. Bact., **130**: 946 (1977) and Hsiao *et al.*, Proc. Natl. Acad. Sci. (USA), **76**: 3829 (1979). However, other methods  
25 for introducing DNA into cells, such as by nuclear microinjection, electroporation, bacterial protoplast fusion with intact cells, or polycations, *e.g.*, polybrene or polyornithine, may also be used. For various techniques for transforming mammalian cells, *see*, Keown *et al.*, Methods in Enzymology, **185**: 527-537 (1990) and Mansour *et al.*, Nature, **336**: 348-352 (1988).

- Suitable host cells for cloning or expressing the DNA in the vectors herein include prokaryote, yeast, or higher  
30 eukaryote cells. Suitable prokaryotes include, but are not limited to, eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacteriaceae such as *E. coli*. Various *E. coli* strains are publicly available, such as *E. coli* K12 strain MM294 (ATCC 31,446); *E. coli* X1776 (ATCC 31,537); *E. coli* strain W3110 (ATCC 27,325); and K5 772 (ATCC 53,635). Other suitable prokaryotic host cells include Enterobacteriaceae such as *Escherichia*, *e.g.*, *E. coli*, *Enterobacter*, *Erwinia*, *Klebsiella*, *Proteus*, *Salmonella*, *e.g.*, *Salmonella typhimurium*,  
35 *Serratia*, *e.g.*, *Serratia marcescans*, and *Shigella*, as well as *Bacilli* such as *B. subtilis* and *B. licheniformis* (*e.g.*, *B. licheniformis* 41P disclosed in DD 266,710 published 12 April 1989), *Pseudomonas* such as *P. aeruginosa*, and

*Streptomyces*. These examples are illustrative rather than limiting. Strain W3110 is one particularly preferred host or parent host because it is a common host strain for recombinant DNA product fermentations. Preferably, the host cell secretes minimal amounts of proteolytic enzymes. For example, strain W3110 may be modified to effect a genetic mutation in the genes encoding proteins endogenous to the host, with examples of such hosts including *E. coli* W3110 strain 1A2, which has the complete genotype *tonA*; *E. coli* W3110 strain 9E4, which has the complete genotype *tonA ptr3*; *E. coli* W3110 strain 27C7 (ATCC 55,244), which has the complete genotype *tonA ptr3 phoA E15 (argF-lac)169 degP ompT kan'*; *E. coli* W3110 strain 37D6, which has the complete genotype *tonA ptr3 phoA E15 (argF-lac)169 degP ompT rbs7 ilvG kan'*; *E. coli* W3110 strain 40B4, which is strain 37D6 with a non-kanamycin resistant *degP* deletion mutation; and an *E. coli* strain having mutant periplasmic protease disclosed in U.S. Patent No. 4,946,783 issued 7 August 1990. Alternatively, *in vitro* methods of cloning, e.g., PCR or other nucleic acid polymerase reactions, are suitable.

In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for vectors encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246. *Saccharomyces cerevisiae* is a commonly used lower eukaryotic host microorganism. Others include *Schizosaccharomyces pombe* (Beach and Nurse, Nature, 290: 140 [1981]; EP 139,383 published 2 May 1985); *Kluyveromyces* hosts (U.S. Patent No. 4,943,529; Fleer *et al.*, Bio/Technology, 9: 968-975 (1991)) such as, e.g., *K. lactis* (MW98-8C, CBS683, CBS4574; Louvencourt *et al.*, J. Bacteriol., 737 [1983]), *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickerhamii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K. drosophilum* (ATCC 36,906; Van den Berg *et al.*, Bio/Technology, 8: 135 (1990)), *K. thermotolerans*, and *K. marxianus*; *yarrowia* (EP 402,226); *Pichia pastoris* (EP 183,070; Sreekrishna *et al.*, J. Basic Microbiol., 28: 265-278 [1988]); *Candida*; *Trichoderma reesia* (EP 244,234); *Neurospora crassa* (Case *et al.*, Proc. Natl. Acad. Sci. USA, 76: 5259-5263 [1979]); *Schwanniomyces* such as *Schwanniomyces occidentalis* (EP 394,538 published 31 October 1990); and filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, *Tolyptocladium* (WO 91/00357 published 10 January 1991), and *Aspergillus* hosts such as *A. nidulans* (Ballance *et al.*, Biochem. Biophys. Res. Commun., 112: 284-289 [1983]; Tilburn *et al.*, Gene, 26: 205-221 [1983]; Yelton *et al.*, Proc. Natl. Acad. Sci. USA, 81: 1470-1474 [1984]) and *A. niger* (Kelly and Hynes, EMBO J., 4: 475-479 [1985]). Methylophilic yeasts are suitable herein and include, but are not limited to, yeast capable of growth on methanol selected from the genera consisting of *Hansenula*, *Candida*, *Kloeckera*, *Pichia*, *Saccharomyces*, *Torulopsis*, and *Rhodotorula*. A list of specific species that are exemplary of this class of yeasts may be found in C. Anthony, The Biochemistry of Methylophilic, 269 (1982).

Suitable host cells for the expression of nucleic acid encoding glycosylated PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 are derived from multicellular organisms. Examples of invertebrate cells include insect cells such as *Drosophila* S2 and *Spodoptera*

Sf9, as well as plant cells. Examples of useful mammalian host cell lines include Chinese hamster ovary (CHO) and COS cells. More specific examples include monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham *et al.*, J. Gen. Virol., 36: 59 (1977)); Chinese hamster ovary cells/-DHFR (CHO, Urlaub and Chasin, Proc. Natl. Acad. Sci. USA, 77:4216 (1980)); mouse sertoli cells (TM4, Mather, Biol. Reprod., 23:243-251 (1980)); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); and mouse mammary tumor (MMT 060562, ATCC CCL51). The selection of the appropriate host cell is deemed to be within the skill in the art.

### iii. Selection and Use of a Replicable Vector

The nucleic acid (*e.g.*, cDNA or genomic DNA) encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 may be inserted into a replicable vector for cloning (amplification of the DNA) or for expression. Various vectors are publicly available. The vector may, for example, be in the form of a plasmid, cosmid, viral particle, or phage. The appropriate nucleic acid sequence may be inserted into the vector by a variety of procedures. In general, DNA is inserted into an appropriate restriction endonuclease site(s) using techniques known in the art. Vector components generally include, but are not limited to, one or more of a signal sequence if the sequence is to be secreted, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence. Construction of suitable vectors containing one or more of these components employs standard ligation techniques that are known to the skilled artisan.

The PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 may be produced recombinantly not only directly, but also as a fusion polypeptide with a heterologous polypeptide, which may be a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide. In general, the signal sequence may be a component of the vector, or it may be a part of the DNA encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 that is inserted into the vector. The signal sequence may be a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders. For yeast secretion the signal sequence may be, *e.g.*, the yeast invertase leader, alpha factor leader (including *Saccharomyces* and *Kluyveromyces*  $\alpha$ -factor leaders, the latter described in U.S. Patent No. 5,010,182), or acid phosphatase leader, the *C. albicans* glucoamylase leader (EP 362,179 published 4 April 1990), or the signal described in WO 90/13646 published 15 November 1990. In mammalian cell expression, mammalian signal sequences may be used to direct secretion of the protein, such as



signal sequences from secreted polypeptides of the same or related species, as well as viral secretory leaders.

Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the  $2\mu$  plasmid origin is suitable for yeast, and various viral origins (SV40, polyoma, adenovirus, VSV, or BPV) are useful for cloning vectors in mammalian cells.

Expression and cloning vectors will typically contain a selection gene, also termed a selectable marker. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, *e.g.*, ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, or (c) supply critical nutrients not available from complex media, *e.g.*, the gene encoding D-alanine racemase for *Bacilli*.

An example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the nucleic acid encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246, such as DHFR or thymidine kinase. An appropriate host cell when wild-type DHFR is employed is the CHO cell line deficient in DHFR activity, prepared and propagated as described by Urlaub *et al.*, Proc. Natl. Acad. Sci. USA, 77: 4216 (1980). A suitable selection gene for use in yeast is the *trp1* gene present in the yeast plasmid YRp7. Stinchcomb *et al.*, Nature, 282: 39 (1979); Kingsman *et al.*, Gene, 7: 141 (1979); Tschemper *et al.*, Gene, 10: 157 (1980). The *trp1* gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1. Jones, Genetics, 85: 12 (1977).

Expression and cloning vectors usually contain a promoter operably linked to the nucleic acid sequence encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 to direct mRNA synthesis. Promoters recognized by a variety of potential host cells are well known. Promoters suitable for use with prokaryotic hosts include the  $\beta$ -lactamase and lactose promoter systems (Chang *et al.*, Nature, 275: 615 (1978); Goeddel *et al.*, Nature, 281: 544 (1979)), alkaline phosphatase, a tryptophan (*trp*) promoter system (Goeddel, Nucleic Acids Res., 8: 4057 (1980); EP 36,776), and hybrid promoters such as the *tac* promoter. deBoer *et al.*, Proc. Natl. Acad. Sci. USA, 80: 21-25 (1983). Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246.

Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase (Hitzeman *et al.*, J. Biol. Chem., 255: 2073 (1980)) or other glycolytic enzymes (Hess *et al.*, J. Adv. Enzyme Reg., 7: 149 (1968); Holland, Biochemistry, 17: 4900 (1978)), such as enolase, glyceraldehyde-

3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

5 Other yeast promoters that are inducible promoters having the additional advantage of transcription controlled by growth conditions are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in EP 73,657.

10 PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 nucleic acid transcription from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus (UK 2,211,504 published 5 July 1989), adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, 15 hepatitis-B virus, and Simian Virus 40 (SV40); by heterologous mammalian promoters, e.g., the actin promoter or an immunoglobulin promoter; and by heat-shock promoters, provided such promoters are compatible with the host cell systems.

Transcription of a DNA encoding the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, 20 PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 by higher eukaryotes may be increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp, that act on a promoter to increase its transcription. Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin,  $\alpha$ -fetoprotein, and insulin). Typically, however, one will use an enhancer from 25 a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. The enhancer may be spliced into the vector at a position 5' or 3' to the sequence coding for PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, 30 PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246, but is preferably located at a site 5' from the promoter.

Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated 35 regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221,

PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246.

- Still other methods, vectors, and host cells suitable for adaptation to the synthesis of PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 in recombinant vertebrate cell culture are described in Gething *et al.*, Nature, 293: 620-625 (1981); Mantei *et al.*, Nature, 281: 40-46 (1979); EP 117,060; and EP 117,058.

#### iv. Detecting Gene Amplification/Expression

- Gene amplification and/or expression may be measured in a sample directly, for example, by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA (Thomas, Proc. Natl. Acad. Sci. USA, 77:5201-5205 (1980)), dot blotting (DNA analysis), or *in situ* hybridization, using an appropriately labeled probe, based on the sequences provided herein. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes. The antibodies in turn may be labeled and the assay may be carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected.

- Gene expression, alternatively, may be measured by immunological methods, such as immunohistochemical staining of cells or tissue sections and assay of cell culture or body fluids, to quantitate directly the expression of gene product. Antibodies useful for immunohistochemical staining and/or assay of sample fluids may be either monoclonal or polyclonal, and may be prepared in any mammal. Conveniently, the antibodies may be prepared against a native-sequence PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide or against a synthetic peptide based on the DNA sequences provided herein or against exogenous sequence fused to DNA encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 and encoding a specific antibody epitope.

#### v. Purification of Polypeptide

- Forms of PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptides may be recovered from culture medium or from host cell lysates. If membrane-bound, it can be released from the membrane using a suitable detergent solution (*e.g.*, TRITON-X™ 100) or by enzymatic cleavage. Cells employed in expression of nucleic acid encoding the PRO172, PRO175, PRO178,

PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide can be disrupted by various physical or chemical means, such as freeze-thaw cycling, sonication, mechanical disruption, or cell-lysing agents.

It may be desired to purify the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide from recombinant cell proteins or polypeptides. The following procedures are exemplary of suitable purification procedures: by fractionation on an ion-exchange column; ethanol precipitation; reverse phase HPLC; chromatography on silica or on a cation-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75; protein A Sepharose columns to remove contaminants such as IgG; and metal chelating columns to bind epitope-tagged forms of the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide. Various methods of protein purification may be employed and such methods are known in the art and described, for example, in Deutscher, Methods in Enzymology, 182 (1990); Scopes, Protein Purification: Principles and Practice (Springer-Verlag: New York, 1982). The purification step(s) selected will depend, for example, on the nature of the production process used and the particular PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 produced.

D. Uses of the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptides

i. Assays for Cardiovascular, Endothelial, and Angiogenic Activity

Various assays can be used to test the polypeptide herein for cardiovascular, endothelial, and angiogenic activity. Such assays include those provided in the Examples below.

Assays for testing for endothelin antagonist activity, as disclosed in U.S. Pat. No. 5,773,414, include a rat heart ventricle binding assay where the polypeptide is tested for its ability to inhibit iodinated endothelin-1 binding in a receptor assay, an endothelin receptor binding assay testing for intact cell binding of radiolabeled endothelin-1 using rabbit renal artery vascular smooth muscle cells, an inositol phosphate accumulation assay where functional activity is determined in Rat-1 cells by measuring intra-cellular levels of second messengers, an arachidonic acid release

assay that measures the ability of added compounds to reduce endothelin-stimulated arachidonic acid release in cultured vascular smooth muscles, *in vitro* (isolated vessel) studies using endothelium from male New Zealand rabbits, and *in vivo* studies using male Sprague-Dawley rats.

Assays for tissue generation activity include, without limitation, those described in WO 95/16035 (bone, cartilage, tendon); WO 95/05846 (nerve, neuronal), and WO 91/07491 (skin, endothelium).

Assays for wound-healing activity include, for example, those described in Winter, Epidermal Wound Healing, Maibach, HI and Rovee, DT, eds. (Year Book Medical Publishers, Inc., Chicago), pp. 71-112, as modified by the article of Eaglstein and Mertz, J. Invest. Dermatol., 71: 382-384 (1978).

An assay to screen for a test molecule relating to a PRO polypeptide that binds an endothelin B<sub>1</sub> (ETB<sub>1</sub>) receptor polypeptide and modulates signal transduction activity involves providing a host cell transformed with a DNA encoding endothelin B<sub>1</sub> receptor polypeptide, exposing the cells to the test candidate, and measuring endothelin B<sub>1</sub> receptor signal transduction activity, as described, *e.g.*, in U.S. Pat. No. 5,773,223.

There are several cardiac hypertrophy assays. *In vitro* assays include induction of spreading of adult rat cardiac myocytes. In this assay, ventricular myocytes are isolated from a single (male Sprague-Dawley) rat, essentially following a modification of the procedure described in detail by Piper *et al.*, "Adult ventricular rat heart muscle cells" in Cell Culture Techniques in Heart and Vessel Research, H.M. Piper, ed. (Berlin: Springer-Verlag, 1990), pp. 36-60. This procedure permits the isolation of adult ventricular myocytes and the long-term culture of these cells in the rod-shaped phenotype. Phenylephrine and Prostaglandin F<sub>2α</sub> (PGF<sub>2α</sub>) have been shown to induce a spreading response in these adult cells. The inhibition of myocyte spreading induced by PGF<sub>2α</sub> or PGF<sub>2α</sub> analogs (*e.g.*, fluprostenol) and phenylephrine by various potential inhibitors of cardiac hypertrophy is then tested.

One example of an *in vivo* assay is a test for inhibiting cardiac hypertrophy induced by fluprostenol *in vivo*. This pharmacological model tests the ability of the PRO polypeptide to inhibit cardiac hypertrophy induced in rats (*e.g.*, male Wistar or Sprague-Dawley) by subcutaneous injection of fluprostenol (an agonist analog of PGF<sub>2α</sub>). It is known that rats with pathologic cardiac hypertrophy induced by myocardial infarction have chronically elevated levels of extractable PGF<sub>2α</sub> in their myocardium. Lai *et al.*, Am. J. Physiol. (Heart Circ. Physiol.), 271: H2197-H2208 (1996). Accordingly, factors that can inhibit the effects of fluprostenol on myocardial growth *in vivo* are potentially useful for treating cardiac hypertrophy. The effects of the PRO polypeptide on cardiac hypertrophy are determined by measuring the weight of heart, ventricles, and left ventricle (normalized by body weight) relative to fluprostenol-treated rats not receiving the PRO polypeptide.

Another example of an *in vivo* assay is the pressure-overload cardiac hypertrophy assay. For *in vivo* testing it is common to induce pressure-overload cardiac hypertrophy by constriction of the abdominal aorta of test animals. In a typical protocol, rats (*e.g.*, male Wistar or Sprague-Dawley) are treated under anesthesia, and the abdominal aorta of each rat is narrowed down just below the diaphragm. Beznak M., Can. J. Biochem. Physiol., 33: 985-94 (1955). The aorta is exposed through a surgical incision, and a blunted needle is placed next to the vessel. The aorta is constricted with a ligature of silk thread around the needle, which is immediately removed and which reduces the lumen of the aorta to the diameter of the needle. This approach is described, for example, in Rossi *et al.*, Am. Heart J., 124: 700-709 (1992) and O'Rourke and Reibel, P.S.E.M.B., 200: 95-100 (1992).

In yet another *in vivo* assay, the effect on cardiac hypertrophy following experimentally induced myocardial infarction (MI) is measured. Acute MI is induced in rats by left coronary artery ligation and confirmed by electrocardiographic examination. A sham-operated group of animals is also prepared as control animals. Earlier data have shown that cardiac hypertrophy is present in the group of animals with MI, as evidenced by an 18% increase in heart weight-to-body weight ratio. Lai *et al.*, *supra*. Treatment of these animals with candidate blockers of cardiac hypertrophy, *e.g.*, PRO polypeptide, provides valuable information about the therapeutic potential of the candidates tested. One further such assay test for induction of cardiac hypertrophy is disclosed in U.S. Pat. No. 5,773,415, using Sprague-Dawley rats.

For cancer, a variety of well-known animal models can be used to further understand the role of the genes identified herein in the development and pathogenesis of tumors, and to test the efficacy of candidate therapeutic agents, including antibodies and other antagonists of the native PRO polypeptides, such as small-molecule antagonists. The *in vivo* nature of such models makes them particularly predictive of responses in human patients. Animal models of tumors and cancers (*e.g.*, breast cancer, colon cancer, prostate cancer, lung cancer, etc.) include both non-recombinant and recombinant (transgenic) animals. Non-recombinant animal models include, for example, rodent, *e.g.*, murine models. Such models can be generated by introducing tumor cells into syngeneic mice using standard techniques, *e.g.*, subcutaneous injection, tail vein injection, spleen implantation, intraperitoneal implantation, implantation under the renal capsule, or orthotopic implantation, *e.g.*, colon cancer cells implanted in colonic tissue. See, *e.g.*, PCT publication No. WO 97/33551, published September 18, 1997. Probably the most often used animal species in oncological studies are immunodeficient mice and, in particular, nude mice. The observation that the nude mouse with thymic hypo/aplasia could successfully act as a host for human tumor xenografts has led to its widespread use for this purpose. The autosomal recessive *nu* gene has been introduced into a very large number of distinct congenic strains of nude mouse, including, for example, ASW, A/He, AKR, BALB/c, B10.LP, C17, C3H, C57BL, C57, CBA, DBA, DDD, I/st, NC, NFR, NFS, NFS/N, NZB, NZC, NZW, P, RIII, and SJL. In addition, a wide variety of other animals with inherited immunological defects other than the nude mouse have been bred and used as recipients of tumor xenografts. For further details see, *e.g.*, The Nude Mouse in Oncology Research, E. Boven and B. Winograd, eds. (CRC Press, Inc., 1991).

The cells introduced into such animals can be derived from known tumor/cancer cell lines, such as any of the above-listed tumor cell lines, and, for example, the B104-1-1 cell line (stable NIH-3T3 cell line transfected with the *neu* protooncogene); *ras*-transfected NIH-3T3 cells; Caco-2 (ATCC HTB-37); or a moderately well-differentiated grade II human colon adenocarcinoma cell line, HT-29 (ATCC HTB-38); or from tumors and cancers. Samples of tumor or cancer cells can be obtained from patients undergoing surgery, using standard conditions involving freezing and storing in liquid nitrogen. Karmali *et al.*, Br. J. Cancer, 48: 689-696 (1983).

Tumor cells can be introduced into animals such as nude mice by a variety of procedures. The subcutaneous (s.c.) space in mice is very suitable for tumor implantation. Tumors can be transplanted s.c. as solid blocks, as needle biopsies by use of a trochar, or as cell suspensions. For solid-block or trochar implantation, tumor tissue fragments of suitable size are introduced into the s.c. space. Cell suspensions are freshly prepared from primary tumors or stable tumor cell lines, and injected subcutaneously. Tumor cells can also be injected as subdermal

implants. In this location, the inoculum is deposited between the lower part of the dermal connective tissue and the s.c. tissue.

Animal models of breast cancer can be generated, for example, by implanting rat neuroblastoma cells (from which the *neu* oncogene was initially isolated), or *neu*-transformed NIH-3T3 cells into nude mice, essentially as described by Drebin *et al.* Proc. Nat. Acad. Sci. USA, 83: 9129-9133 (1986).

Similarly, animal models of colon cancer can be generated by passaging colon cancer cells in animals, *e.g.*, nude mice, leading to the appearance of tumors in these animals. An orthotopic transplant model of human colon cancer in nude mice has been described, for example, by Wang *et al.*, Cancer Research, 54: 4726-4728 (1994) and Too *et al.*, Cancer Research, 55: 681-684 (1995). This model is based on the so-called "METAMOUSE"<sup>TM</sup> sold by AntiCancer, Inc., (San Diego, California).

Tumors that arise in animals can be removed and cultured *in vitro*. Cells from the *in vitro* cultures can then be passaged to animals. Such tumors can serve as targets for further testing or drug screening. Alternatively, the tumors resulting from the passage can be isolated and RNA from pre-passage cells and cells isolated after one or more rounds of passage analyzed for differential expression of genes of interest. Such passaging techniques can be performed with any known tumor or cancer cell lines.

For example, Meth A, CMS4, CMS5, CMS21, and WEHI-164 are chemically induced fibrosarcomas of BALB/c female mice (DeLeo *et al.*, J. Exp. Med., 146: 720 (1977)), which provide a highly controllable model system for studying the anti-tumor activities of various agents. Palladino *et al.*, J. Immunol., 138: 4023-4032 (1987). Briefly, tumor cells are propagated *in vitro* in cell culture. Prior to injection into the animals, the cell lines are washed and suspended in buffer, at a cell density of about  $10 \times 10^6$  to  $10 \times 10^7$  cells/ml. The animals are then infected subcutaneously with 10 to 100  $\mu$ l of the cell suspension, allowing one to three weeks for a tumor to appear.

In addition, the Lewis lung (3LL) carcinoma of mice, which is one of the most thoroughly studied experimental tumors, can be used as an investigational tumor model. Efficacy in this tumor model has been correlated with beneficial effects in the treatment of human patients diagnosed with small-cell carcinoma of the lung (SCCL). This tumor can be introduced in normal mice upon injection of tumor fragments from an affected mouse or of cells maintained in culture. Zupi *et al.*, Br. J. Cancer, 41: suppl. 4, 30 (1980). Evidence indicates that tumors can be started from injection of even a single cell and that a very high proportion of infected tumor cells survive. For further information about this tumor model *see*, Zacharski, Haemostasis, 16: 300-320 (1986).

One way of evaluating the efficacy of a test compound in an animal model with an implanted tumor is to measure the size of the tumor before and after treatment. Traditionally, the size of implanted tumors has been measured with a slide caliper in two or three dimensions. The measure limited to two dimensions does not accurately reflect the size of the tumor; therefore, it is usually converted into the corresponding volume by using a mathematical formula. However, the measurement of tumor size is very inaccurate. The therapeutic effects of a drug candidate can be better described as treatment-induced growth delay and specific growth delay. Another important variable in the description of tumor growth is the tumor volume doubling time. Computer programs for the calculation and description of tumor growth are also available, such as the program reported by Rygaard and Spang-Thomsen, Proc. 6th Int. Workshop on Immune-Deficient Animals, Wu and Sheng eds. (Basel, 1989), p. 301.

It is noted, however, that necrosis and inflammatory responses following treatment may actually result in an increase in tumor size, at least initially. Therefore, these changes need to be carefully monitored, by a combination of a morphometric method and flow cytometric analysis.

Further, recombinant (transgenic) animal models can be engineered by introducing the coding portion of the PRO gene identified herein into the genome of animals of interest, using standard techniques for producing transgenic animals. Animals that can serve as a target for transgenic manipulation include, without limitation, mice, rats, rabbits, guinea pigs, sheep, goats, pigs, and non-human primates, *e.g.*, baboons, chimpanzees and monkeys. Techniques known in the art to introduce a transgene into such animals include pronucleic microinjection (U.S. Patent No. 4,873,191); retrovirus-mediated gene transfer into germ lines (*e.g.*, Van der Putten *et al.*, Proc. Natl. Acad. Sci. USA, 82: 6148-615 (1985)); gene targeting in embryonic stem cells (Thompson *et al.*, Cell, 56: 313-321 (1989)); electroporation of embryos (Lo, Mol. Cell. Biol., 3: 1803-1814 (1983)); and sperm-mediated gene transfer. Lavitrano *et al.*, Cell, 57: 717-73 (1989). For a review, *see* for example, U.S. Patent No. 4,736,866.

For the purpose of the present invention, transgenic animals include those that carry the transgene only in part of their cells ("mosaic animals"). The transgene can be integrated either as a single transgene, or in concatamers, *e.g.*, head-to-head or head-to-tail tandems. Selective introduction of a transgene into a particular cell type is also possible by following, for example, the technique of Lasko *et al.*, Proc. Natl. Acad. Sci. USA, 89: 6232-636 (1992).

The expression of the transgene in transgenic animals can be monitored by standard techniques. For example, Southern blot analysis or PCR amplification can be used to verify the integration of the transgene. The level of mRNA expression can then be analyzed using techniques such as *in situ* hybridization, Northern blot analysis, PCR, or immunocytochemistry. The animals are further examined for signs of tumor or cancer development.

Alternatively, "knock-out" animals can be constructed that have a defective or altered gene encoding a PRO polypeptide identified herein, as a result of homologous recombination between the endogenous gene encoding the PRO polypeptide and altered genomic DNA encoding the same polypeptide introduced into an embryonic cell of the animal. For example, cDNA encoding a particular PRO polypeptide can be used to clone genomic DNA encoding that polypeptide in accordance with established techniques. A portion of the genomic DNA encoding a particular PRO polypeptide can be deleted or replaced with another gene, such as a gene encoding a selectable marker that can be used to monitor integration. Typically, several kilobases of unaltered flanking DNA (both at the 5' and 3' ends) are included in the vector. *See, e.g.*, Thomas and Capecchi, Cell, 51: 503 (1987) for a description of homologous recombination vectors. The vector is introduced into an embryonic stem cell line (*e.g.*, by electroporation) and cells in which the introduced DNA has homologously recombined with the endogenous DNA are selected. *See, e.g.*, Li *et al.*, Cell, 69: 915 (1992). The selected cells are then injected into a blastocyst of an animal (*e.g.*, a mouse or rat) to form aggregation chimeras. *See, e.g.*, Bradley, in Teratocarcinomas and Embryonic Stem Cells: A Practical Approach. E. J. Robertson, ed. (IRL: Oxford, 1987), pp. 113-152. A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term to create a "knock-out" animal. Progeny harboring the homologously recombined DNA in their germ cells can be identified by standard techniques and used to breed animals in which all cells of the animal contain the homologously recombined DNA. Knockout animals can be characterized, for instance, by their ability to defend against certain



pathological conditions and by their development of pathological conditions due to absence of the PRO polypeptide.

The efficacy of antibodies specifically binding the PRO polypeptides identified herein, and other drug candidates, can be tested also in the treatment of spontaneous animal tumors. A suitable target for such studies is the feline oral squamous cell carcinoma (SCC). Feline oral SCC is a highly invasive, malignant tumor that is the most common oral malignancy of cats, accounting for over 60% of the oral tumors reported in this species. It rarely metastasizes to distant sites, although this low incidence of metastasis may merely be a reflection of the short survival times for cats with this tumor. These tumors are usually not amenable to surgery, primarily because of the anatomy of the feline oral cavity. At present, there is no effective treatment for this tumor. Prior to entry into the study, each cat undergoes complete clinical examination and biopsy, and is scanned by computed tomography (CT). Cats diagnosed with sublingual oral squamous cell tumors are excluded from the study. The tongue can become paralyzed as a result of such tumor, and even if the treatment kills the tumor, the animals may not be able to feed themselves. Each cat is treated repeatedly, over a longer period of time. Photographs of the tumors will be taken daily during the treatment period, and at each subsequent recheck. After treatment, each cat undergoes another CT scan. CT scans and thoracic radiograms are evaluated every 8 weeks thereafter. The data are evaluated for differences in survival, response, and toxicity as compared to control groups. Positive response may require evidence of tumor regression, preferably with improvement of quality of life and/or increased life span.

In addition, other spontaneous animal tumors, such as fibrosarcoma, adenocarcinoma, lymphoma, chondroma, or leiomyosarcoma of dogs, cats, and baboons can also be tested. Of these, mammary adenocarcinoma in dogs and cats is a preferred model as its appearance and behavior are very similar to those in humans. However, the use of this model is limited by the rare occurrence of this type of tumor in animals.

Other *in vitro* and *in vivo* cardiovascular, endothelial, and angiogenic tests known in the art are also suitable herein.

#### ii. Tissue Distribution

The results of the cardiovascular, endothelial, and angiogenic assays herein can be verified by further studies, such as by determining mRNA expression in various human tissues.

As noted before, gene amplification and/or gene expression in various tissues may be measured by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA (Thomas, Proc. Natl. Acad. Sci. USA, 77:5201-5205 (1980)), dot blotting (DNA analysis), or *in situ* hybridization, using an appropriately labeled probe, based on the sequences provided herein. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes.

Gene expression in various tissues, alternatively, may be measured by immunological methods, such as immunohistochemical staining of tissue sections and assay of cell culture or body fluids, to quantitate directly the expression of gene product. Antibodies useful for immunohistochemical staining and/or assay of sample fluids may be either monoclonal or polyclonal, and may be prepared in any mammal. Conveniently, the antibodies may be prepared against a native-sequence PRO polypeptide or against a synthetic peptide based on the DNA sequences

provided herein or against exogenous sequence fused to PRO DNA and encoding a specific antibody epitope. General techniques for generating antibodies, and special protocols for *in situ* hybridization are provided hereinbelow.

iii. Antibody Binding Studies

5 The results of the cardiovascular, endothelial, and angiogenic study can be further verified by antibody binding studies, in which the ability of anti-PRO antibodies to inhibit the effect of the PRO polypeptides on endothelial cells or other cells used in the cardiovascular, endothelial, and angiogenic assays is tested. Exemplary antibodies include polyclonal, monoclonal, humanized, bispecific, and heteroconjugate antibodies, the preparation of which will be described hereinbelow.

10 Antibody binding studies may be carried out in any known assay method, such as competitive binding assays, direct and indirect sandwich assays, and immunoprecipitation assays. Zola, Monoclonal Antibodies: A Manual of Techniques (CRC Press, Inc., 1987), pp.147-158.

Competitive binding assays rely on the ability of a labeled standard to compete with the test sample analyte for binding with a limited amount of antibody. The amount of target protein in the test sample is inversely  
15 proportional to the amount of standard that becomes bound to the antibodies. To facilitate determining the amount of standard that becomes bound, the antibodies preferably are insolubilized before or after the competition, so that the standard and analyte that are bound to the antibodies may conveniently be separated from the standard and analyte that remain unbound.

Sandwich assays involve the use of two antibodies, each capable of binding to a different immunogenic  
20 portion, or epitope, of the protein to be detected. In a sandwich assay, the test sample analyte is bound by a first antibody that is immobilized on a solid support, and thereafter a second antibody binds to the analyte, thus forming an insoluble three-part complex. See, e.g., US Pat No. 4,376,110. The second antibody may itself be labeled with a detectable moiety (direct sandwich assays) or may be measured using an anti-immunoglobulin antibody that is labeled with a detectable moiety (indirect sandwich assay). For example, one type of sandwich assay is an ELISA  
25 assay, in which case the detectable moiety is an enzyme.

For immunohistochemistry, the tissue sample may be fresh or frozen or may be embedded in paraffin and fixed with a preservative such as formalin, for example.

iv. Cell-Based Tumor Assays

Cell-based assays and animal models for cardiovascular, endothelial, and angiogenic disorders, such as tumors,  
30 can be used to verify the findings of a cardiovascular, endothelial, and angiogenic assay herein, and further to understand the relationship between the genes identified herein and the development and pathogenesis of undesirable cardiovascular, endothelial, and angiogenic cell growth. The role of gene products identified herein in the development and pathology of undesirable cardiovascular, endothelial, and angiogenic cell growth, e.g., tumor cells, can be tested by using cells or cells lines that have been identified as being stimulated or inhibited by the PRO  
35 polypeptide herein. Such cells include, for example, those set forth in the Examples below.

In a different approach, cells of a cell type known to be involved in a particular cardiovascular, endothelial, and angiogenic disorder are transfected with the cDNAs herein, and the ability of these cDNAs to induce excessive growth or inhibit growth is analyzed. If the cardiovascular, endothelial, and angiogenic disorder is cancer, suitable tumor cells include, for example, stable tumor cells lines such as the B104-1-1 cell line (stable NIH-3T3 cell line transfected with the *neu* protooncogene) and *ras*-transfected NIH-3T3 cells, which can be transfected with the desired gene and monitored for tumorigenic growth. Such transfected cell lines can then be used to test the ability of poly- or monoclonal antibodies or antibody compositions to inhibit tumorigenic cell growth by exerting cytostatic or cytotoxic activity on the growth of the transformed cells, or by mediating antibody-dependent cellular cytotoxicity (ADCC). Cells transfected with the coding sequences of the genes identified herein can further be used to identify drug candidates for the treatment of cardiovascular, endothelial, and angiogenic disorders such as cancer.

In addition, primary cultures derived from tumors in transgenic animals (as described above) can be used in the cell-based assays herein, although stable cell lines are preferred. Techniques to derive continuous cell lines from transgenic animals are well known in the art. See, e.g., Small *et al.*, Mol. Cell. Biol., 5: 642-648 (1985).

#### v. Gene Therapy

The PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide herein and polypeptidyl agonists and antagonists may be employed in accordance with the present invention by expression of such polypeptides *in vivo*, which is often referred to as gene therapy.

There are two major approaches to getting the nucleic acid (optionally contained in a vector) into the patient's cells: *in vivo* and *ex vivo*. For *in vivo* delivery the nucleic acid is injected directly into the patient, usually at the sites where the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide is required, *i.e.*, the site of synthesis of the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide, if known, and the site (*e.g.*, wound) where biological activity of PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide is needed. For *ex vivo* treatment, the patient's cells are removed, the nucleic acid is introduced into these isolated cells, and the modified cells are administered to the patient either directly or, for example, encapsulated within porous membranes that are implanted into the patient (*see, e.g.*, U.S. Pat. Nos. 4,892,538 and 5,283,187). There are a variety of techniques

available for introducing nucleic acids into viable cells. The techniques vary depending upon whether the nucleic acid is transferred into cultured cells *in vitro*, or transferred *in vivo* in the cells of the intended host. Techniques suitable for the transfer of nucleic acid into mammalian cells *in vitro* include the use of liposomes, electroporation, microinjection, transduction, cell fusion, DEAE-dextran, the calcium phosphate precipitation method, etc.

- 5 Transduction involves the association of a replication-defective, recombinant viral (preferably retroviral) particle with a cellular receptor, followed by introduction of the nucleic acids contained by the particle into the cell. A commonly used vector for *ex vivo* delivery of the gene is a retrovirus.

- 10 The currently preferred *in vivo* nucleic acid transfer techniques include transfection with viral or non-viral vectors (such as adenovirus, lentivirus, Herpes simplex I virus, or adeno-associated virus (AAV)) and lipid-based systems (useful lipids for lipid-mediated transfer of the gene are, for example, DOTMA, DOPE, and DC-Chol; *see, e.g.,* Tonkinson *et al.*, Cancer Investigation, 14(1): 54-65 (1996)). The most preferred vectors for use in gene therapy are viruses, most preferably adenoviruses, AAV, lentiviruses, or retroviruses. A viral vector such as a retroviral vector includes at least one transcriptional promoter/enhancer or locus-defining element(s), or other elements that control gene expression by other means such as alternate splicing, nuclear RNA export, or
- 15 post-translational modification of messenger. In addition, a viral vector such as a retroviral vector includes a nucleic acid molecule that, when transcribed in the presence of a gene encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide, is operably linked
- 20 thereto and acts as a translation initiation sequence. Such vector constructs also include a packaging signal, long terminal repeats (LTRs) or portions thereof, and positive and negative strand primer binding sites appropriate to the virus used (if these are not already present in the viral vector). In addition, such vector typically includes a signal sequence for secretion of the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide from a host cell in which it is placed. Preferably the signal sequence for this purpose is a mammalian signal sequence, most preferably the native signal sequence for the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide. Optionally, the vector construct may also include a signal that directs polyadenylation, as well as one or more restriction sites and a translation termination sequence. By way of example, such vectors will typically include a 5' LTR, a tRNA binding site, a packaging signal, an origin of second-strand DNA synthesis, and a 3' LTR or a portion thereof. Other vectors can be used that are non-viral, such as cationic lipids, polylysine, and
- 35 dendrimers.

In some situations, it is desirable to provide the nucleic acid source with an agent that targets the target cells, such as an antibody specific for a cell-surface membrane protein or the target cell, a ligand for a receptor on the

target cell, etc. Where liposomes are employed, proteins that bind to a cell-surface membrane protein associated with endocytosis may be used for targeting and/or to facilitate uptake, *e.g.*, capsid proteins or fragments thereof tropic for a particular cell type, antibodies for proteins that undergo internalization in cycling, and proteins that target intracellular localization and enhance intracellular half-life. The technique of receptor-mediated endocytosis is described, for example, by Wu *et al.*, J. Biol. Chem., **262**: 4429-4432 (1987); and Wagner *et al.*, Proc. Natl. Acad. Sci. USA, **87**: 3410-3414 (1990). For a review of the currently known gene marking and gene therapy protocols, see, Anderson *et al.*, Science, **256**: 808-813 (1992). See also WO 93/25673 and the references cited therein.

Suitable gene therapy and methods for making retroviral particles and structural proteins can be found in, *e.g.*, U.S. Pat. No. 5,681,746.

#### 10                   vi. Use of Gene as Diagnostic

This invention is also related to the use of the gene encoding the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide as a diagnostic.

15   Detection of a mutated form of the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide will allow a diagnosis of a cardiovascular, endothelial, and angiogenic disease or a susceptibility to a cardiovascular, endothelial, and angiogenic disease, such as a tumor,

20   since mutations in the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide may cause tumors.

Individuals carrying mutations in the genes encoding a human PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide may be detected at the DNA level by a variety of techniques. Nucleic acids for diagnosis may be obtained from a patient's cells, such as from blood, urine, saliva, tissue biopsy, and autopsy material. The genomic DNA may be used directly for

30   detection or may be amplified enzymatically by using PCR (Saiki *et al.*, Nature, **324**: 163-166 (1986)) prior to analysis. RNA or cDNA may also be used for the same purpose. As an example, PCR primers complementary to the nucleic acid encoding the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide can be used to identify and analyze PRO172, PRO175,

35   PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217,

PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide mutations. For example, deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to  
 5 radiolabeled RNA encoding the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide, or alternatively, radiolabeled antisense DNA sequences encoding the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195,  
 10 PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase A digestion or by differences in melting temperatures.

Genetic testing based on DNA sequence differences may be achieved by detection of alteration in  
 15 electrophoretic mobility of DNA fragments in gels with or without denaturing agents. Small sequence deletions and insertions can be visualized by high resolution gel electrophoresis. DNA fragments of different sequences may be distinguished on denaturing formamide gradient gels in which the mobilities of different DNA fragments are retarded in the gel at different positions according to their specific melting or partial melting temperatures. *See, e.g., Myers et al., Science, 230: 1242 (1985).*

20 Sequence changes at specific locations may also be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method, for example, Cotton *et al.*, Proc. Natl. Acad. Sci. USA, 85: 4397-4401 (1985).

Thus, the detection of a specific DNA sequence may be achieved by methods such as hybridization, RNase protection, chemical cleavage, direct DNA sequencing, or the use of restriction enzymes, *e.g.*, restriction fragment  
 25 length polymorphisms (RFLP), and Southern blotting of genomic DNA.

#### vii. Use to Detect PRO Polypeptide Levels

In addition to more conventional gel-electrophoresis and DNA sequencing, mutations can also be detected by *in situ* analysis.

30 Expression of nucleic acid encoding the PRO polypeptide may be linked to vascular disease or neovascularization associated with tumor formation. If the PRO polypeptide has a signal sequence and the mRNA is highly expressed in endothelial cells and to a lesser extent in smooth muscle cells, this indicates that the PRO polypeptide is present in serum. Accordingly, an anti-PRO polypeptide antibody could be used to diagnose vascular disease or neovascularization associated with tumor formation, since an altered level of this PRO polypeptide may be indicative of such disorders.

35 A competition assay may be employed wherein antibodies specific to the PRO polypeptide are attached to a solid support and the labeled PRO polypeptide and a sample derived from the host are passed over the solid support

and the amount of label detected attached to the solid support can be correlated to a quantity of the PRO polypeptide in the sample.

viii. Chromosome Mapping

5 The sequences of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome. Moreover, there is a current need for identifying particular sites on the chromosome. Few chromosome marking reagents based on actual sequence data (repeat polymorphisms) are presently available for marking chromosomal location. The mapping of DNAs to chromosomes according to the present invention is an important first step in correlating those sequences with genes associated with disease.

10 Briefly, sequences can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp) from the cDNA. Computer analysis for the 3'- untranslated region is used to rapidly select primers that do not span more than one exon in the genomic DNA, thus complicating the amplification process. These primers are then used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the primer will yield an amplified fragment.

15 PCR mapping of somatic cell hybrids is a rapid procedure for assigning a particular DNA to a particular chromosome. Using the present invention with the same oligonucleotide primers, sublocalization can be achieved with panels of fragments from specific chromosomes or pools of large genomic clones in an analogous manner. Other mapping strategies that can similarly be used to map to its chromosome include *in situ* hybridization, prescreening with labeled flow-sorted chromosomes, and preselection by hybridization to construct chromosome-specific cDNA libraries.

20 Fluorescence *in situ* hybridization (FISH) of a cDNA clone to a metaphase chromosomal spread can be used to provide a precise chromosomal location in one step. This technique can be used with cDNA as short as 500 or 600 bases; however, clones larger than 2,000 bp have a higher likelihood of binding to a unique chromosomal location with sufficient signal intensity for simple detection. FISH requires use of the clones from which the gene encoding the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, 25 PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide was derived, and the longer the better. For example, 2,000 bp is good, 4,000 bp is better, and more than 4,000 is probably not necessary to get good results a reasonable percentage of the time. For a review of this technique, see, Verma *et al.*, Human Chromosomes: a Manual of Basic Techniques (Pergamon Press, New York, 1988).

30 Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available online through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region is then identified through linkage analysis (coinheritance of physically adjacent genes).

Next, it is necessary to determine the differences in the cDNA or genomic sequence between affected and unaffected individuals. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

With current resolution of physical mapping and genetic mapping techniques, a cDNA precisely localized to a chromosomal region associated with the disease could be one of between 50 and 500 potential causative genes. (This assumes 1 megabase mapping resolution and one gene per 20 kb).

ix. Screening Assays for Drug Candidates

This invention encompasses methods of screening compounds to identify those that mimic the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide (agonists) or prevent the effect of the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide (antagonists). Screening assays for antagonist drug candidates are designed to identify compounds that bind or complex with the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide encoded by the genes identified herein, or otherwise interfere with the interaction of the encoded polypeptides with other cellular proteins. Such screening assays will include assays amenable to high-throughput screening of chemical libraries, making them particularly suitable for identifying small molecule drug candidates.

The assays can be performed in a variety of formats, including protein-protein binding assays, biochemical screening assays, immunoassays, and cell-based assays, which are well characterized in the art.

All assays for antagonists are common in that they call for contacting the drug candidate with a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide encoded by a nucleic acid identified herein under conditions and for a time sufficient to allow these two components to interact.

In binding assays, the interaction is binding and the complex formed can be isolated or detected in the reaction mixture. In a particular embodiment, the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide encoded by the gene identified herein or the drug candidate is immobilized on a solid phase, *e.g.*, on a microtiter plate, by covalent or non-covalent attachments. Non-



covalent attachment generally is accomplished by coating the solid surface with a solution of the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide and drying. Alternatively, an immobilized antibody, *e.g.*, a monoclonal antibody, specific for the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide to be immobilized can be used to anchor it to a solid surface. The assay is performed by adding the non-immobilized component, which may be labeled by a detectable label, to the immobilized component, *e.g.*, the coated surface containing the anchored component. When the reaction is complete, the non-reacted components are removed, *e.g.*, by washing, and complexes anchored on the solid surface are detected. When the originally non-immobilized component carries a detectable label, the detection of label immobilized on the surface indicates that complexing occurred. Where the originally non-immobilized component does not carry a label, complexing can be detected, for example, by using a labeled antibody specifically binding the immobilized complex.

If the candidate compound interacts with but does not bind to a particular PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide encoded by a gene identified herein, its interaction with that polypeptide can be assayed by methods well known for detecting protein-protein interactions. Such assays include traditional approaches, such as, *e.g.*, cross-linking, co-immunoprecipitation, and co-purification through gradients or chromatographic columns. In addition, protein-protein interactions can be monitored by using a yeast-based genetic system described by Fields and co-workers (Fields and Song, *Nature (London)*, **340**: 245-246 (1989); Chien *et al.*, *Proc. Natl. Acad. Sci. USA*, **88**: 9578-9582 (1991)) as disclosed by Chevray and Nathans, *Proc. Natl. Acad. Sci. USA*, **89**: 5789-5793 (1991). Many transcriptional activators, such as yeast GAL4, consist of two physically discrete modular domains, one acting as the DNA-binding domain, the other one functioning as the transcription-activation domain. The yeast expression system described in the foregoing publications (generally referred to as the "two-hybrid system") takes advantage of this property, and employs two hybrid proteins, one in which the target protein is fused to the DNA-binding domain of GAL4, and another, in which candidate activating proteins are fused to the activation domain. The expression of a GAL1-*lacZ* reporter gene under control of a GAL4-activated promoter depends on reconstitution of GAL4 activity via protein-protein interaction. Colonies containing interacting polypeptides are detected with a chromogenic substrate for  $\beta$ -galactosidase. A complete kit (MATCHMAKER™) for identifying protein-protein interactions between two specific proteins using the two-hybrid technique is commercially available from Clontech. This system can also be extended to map protein domains involved in specific protein interactions as well as to pinpoint amino acid residues that are crucial for these interactions.

Compounds that interfere with the interaction of a gene encoding a PRO172, PRO175, PRO178, PRO188,

PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide identified herein and other intra- or extracellular components can be tested as follows: usually a reaction mixture is prepared containing the product of the gene and the intra- or extracellular component under conditions and for a time allowing for the interaction and binding of the two products. To test the ability of a candidate compound to inhibit binding, the reaction is run in the absence and in the presence of the test compound. In addition, a placebo may be added to a third reaction mixture, to serve as positive control. The binding (complex formation) between the test compound and the intra- or extracellular component present in the mixture is monitored as described hereinabove.

5 The formation of a complex in the control reaction(s) but not in the reaction mixture containing the test compound indicates that the test compound interferes with the interaction of the test compound and its reaction partner.

10

If the PRO polypeptide has the ability to stimulate the proliferation of endothelial cells in the presence of the co-mitogen ConA, then one example of a screening method takes advantage of this ability. Specifically, in the proliferation assay, human umbilical vein endothelial cells are obtained and cultured in 96-well flat-bottomed culture plates (Costar, Cambridge, MA) and supplemented with a reaction mixture appropriate for facilitating proliferation of the cells, the mixture containing Con-A (Calbiochem, La Jolla, CA). Con-A and the compound to be screened are added and after incubation at 37°C, cultures are pulsed with <sup>3</sup>H-thymidine and harvested onto glass fiber filters (phD; Cambridge Technology, Watertown, MA). Mean <sup>3</sup>H-thymidine incorporation (cpm) of triplicate cultures is determined using a liquid scintillation counter (Beckman Instruments, Irvine, CA). Significant <sup>3</sup>(H)-thymidine incorporation indicates stimulation of endothelial cell proliferation.

15

20

To assay for antagonists, the assay described above is performed; however, in this assay the PRO polypeptide is added along with the compound to be screened and the ability of the compound to inhibit <sup>3</sup>(H)thymidine incorporation in the presence of the PRO polypeptide indicates that the compound is an antagonist to the PRO polypeptide. Alternatively, antagonists may be detected by combining the PRO polypeptide and a potential antagonist with membrane-bound PRO polypeptide receptors or recombinant receptors under appropriate conditions for a competitive inhibition assay. The PRO polypeptide can be labeled, such as by radioactivity, such that the number of PRO polypeptide molecules bound to the receptor can be used to determine the effectiveness of the potential antagonist. The gene encoding the receptor can be identified by numerous methods known to those of skill in the art, for example, ligand panning and FACS sorting. Coligan *et al.*, Current Protocols in Immun., 1(2): Chapter 5 (1991). Preferably, expression cloning is employed wherein polyadenylated RNA is prepared from a cell responsive to the PRO polypeptide and a cDNA library created from this RNA is divided into pools and used to transfect COS cells or other cells that are not responsive to the PRO polypeptide. Transfected cells that are grown on glass slides are exposed to the labeled PRO polypeptide. The PRO polypeptide can be labeled by a variety of means including iodination or inclusion of a recognition site for a site-specific protein kinase. Following fixation and incubation, the slides are subjected to autoradiographic analysis. Positive pools are identified and sub-pools are prepared and re-transfected using an interactive sub-pooling and re-screening process, eventually yielding a single clone that encodes the putative receptor.

25

30

35

As an alternative approach for receptor identification, the labeled PRO polypeptide can be photoaffinity-linked with cell membrane or extract preparations that express the receptor molecule. Cross-linked material is resolved by PAGE and exposed to X-ray film. The labeled complex containing the receptor can be excised, resolved into peptide fragments, and subjected to protein micro-sequencing. The amino acid sequence obtained from micro-sequencing would be used to design a set of degenerate oligonucleotide probes to screen a cDNA library to identify the gene encoding the putative receptor.

In another assay for antagonists, mammalian cells or a membrane preparation expressing the receptor would be incubated with the labeled PRO polypeptide in the presence of the candidate compound. The ability of the compound to enhance or block this interaction could then be measured.

The compositions useful in the treatment of cardiovascular, endothelial, and angiogenic disorders include, without limitation, antibodies, small organic and inorganic molecules, peptides, phosphopeptides, antisense and ribozyme molecules, triple-helix molecules, etc., that inhibit the expression and/or activity of the target gene product.

More specific examples of potential antagonists include an oligonucleotide that binds to the fusions of immunoglobulin with a PRO polypeptide, and, in particular, antibodies including, without limitation, poly- and monoclonal antibodies and antibody fragments, single-chain antibodies, anti-idiotypic antibodies, and chimeric or humanized versions of such antibodies or fragments, as well as human antibodies and antibody fragments. Alternatively, a potential antagonist may be a closely related protein, for example, a mutated form of the PRO polypeptide that recognizes the receptor but imparts no effect, thereby competitively inhibiting the action of the PRO polypeptide.

Another potential PRO polypeptide antagonist or agonist is an antisense RNA or DNA construct prepared using antisense technology, where, *e.g.*, an antisense RNA or DNA molecule acts to block directly the translation of mRNA by hybridizing to targeted mRNA and preventing protein translation. Antisense technology can be used to control gene expression through triple-helix formation or antisense DNA or RNA, both of which methods are based on binding of a polynucleotide to DNA or RNA. For example, the 5' coding portion of the polynucleotide sequence, which encodes the mature PRO polypeptides herein, is used to design an antisense RNA oligonucleotide of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription (triple helix - *see*, Lee *et al.*, Nucl. Acids Res., **6**:3073 (1979); Cooney *et al.*, Science, **241**: 456 (1988); Dervan *et al.*, Science, **251**:1360 (1991)), thereby preventing transcription and the production of the PRO polypeptide. The antisense RNA oligonucleotide hybridizes to the mRNA *in vivo* and blocks translation of the mRNA molecule into the PRO polypeptide (antisense - Okano, Neurochem., **56**:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression (CRC Press: Boca Raton, FL, 1988). The oligonucleotides described above can also be delivered to cells such that the antisense RNA or DNA may be expressed *in vivo* to inhibit production of the PRO polypeptide. When antisense DNA is used, oligodeoxyribonucleotides derived from the translation-initiation site, *e.g.*, between about -10 and +10 positions of the target gene nucleotide sequence, are preferred.

Antisense RNA or DNA molecules are generally at least about 5 bases in length, about 10 bases in length,

about 15 bases in length, about 20 bases in length, about 25 bases in length, about 30 bases in length, about 35 bases in length, about 40 bases in length, about 45 bases in length, about 50 bases in length, about 55 bases in length, about 60 bases in length, about 65 bases in length, about 70 bases in length, about 75 bases in length, about 80 bases in length, about 85 bases in length, about 90 bases in length, about 95 bases in length, about 100 bases in length, or more.

Potential antagonists include small molecules that bind to the active site, the receptor binding site, or growth factor or other relevant binding site of the PRO polypeptide, thereby blocking the normal biological activity of the PRO polypeptide. Examples of small molecules include, but are not limited to, small peptides or peptide-like molecules, preferably soluble peptides, and synthetic non-peptidyl organic or inorganic compounds.

Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. Ribozymes act by sequence-specific hybridization to the complementary target RNA, followed by endonucleolytic cleavage. Specific ribozyme cleavage sites within a potential RNA target can be identified by known techniques. For further details *see, e.g.*, Rossi, Current Biology, 4: 469-471 (1994), and PCT publication No. WO 97/33551 (published September 18, 1997).

Nucleic acid molecules in triple-helix formation used to inhibit transcription should be single-stranded and composed of deoxynucleotides. The base composition of these oligonucleotides is designed such that it promotes triple-helix formation via Hoogsteen base-pairing rules, which generally require sizeable stretches of purines or pyrimidines on one strand of a duplex. For further details *see, e.g.*, PCT publication No. WO 97/33551, *supra*.

These small molecules can be identified by any one or more of the screening assays discussed hereinabove and/or by any other screening techniques well known for those skilled in the art.

x. Types of Cardiovascular, Endothelial, and Angiogenic Disorders to be Treated

The PRO polypeptides, or agonists or antagonists thereto, that have activity in the cardiovascular, angiogenic, and endothelial assays described herein, and/or whose gene product has been found to be localized to the cardiovascular system, are likely to have therapeutic uses in a variety of cardiovascular, endothelial, and angiogenic disorders, including systemic disorders that affect vessels, such as diabetes mellitus. Their therapeutic utility could include diseases of the arteries, capillaries, veins, and/or lymphatics. Examples of treatments hereunder include treating muscle wasting disease, treating osteoporosis, aiding in implant fixation to stimulate the growth of cells around the implant and therefore facilitate its attachment to its intended site, increasing IGF stability in tissues or in serum, if applicable, and increasing binding to the IGF receptor (since IGF has been shown *in vitro* to enhance human marrow erythroid and granulocytic progenitor cell growth).

The PRO polypeptides or agonists or antagonists thereto may also be employed to stimulate erythropoiesis or granulopoiesis, to stimulate wound healing or tissue regeneration and associated therapies concerned with re-growth of tissue, such as connective tissue, skin, bone, cartilage, muscle, lung, or kidney, to promote angiogenesis, to stimulate or inhibit migration of endothelial cells, and to proliferate the growth of vascular smooth muscle and endothelial cell production. The increase in angiogenesis mediated by the PRO polypeptide or antagonist would be beneficial to ischemic tissues and to collateral coronary development in the heart subsequent to coronary stenosis.

Antagonists are used to inhibit the action of such polypeptides, for example, to limit the production of excess connective tissue during wound healing or pulmonary fibrosis if the PRO polypeptide promotes such production. This would include treatment of acute myocardial infarction and heart failure.

Moreover, the present invention concerns the treatment of cardiac hypertrophy, regardless of the underlying cause, by administering a therapeutically effective dose of the PRO polypeptide, or agonist or antagonist thereto. If the objective is the treatment of human patients, the PRO polypeptide preferably is recombinant human PRO polypeptide (rhPRO polypeptide). The treatment for cardiac hypertrophy can be performed at any of its various stages, which may result from a variety of diverse pathologic conditions, including myocardial infarction, hypertension, hypertrophic cardiomyopathy, and valvular regurgitation. The treatment extends to all stages of the progression of cardiac hypertrophy, with or without structural damage of the heart muscle, regardless of the underlying cardiac disorder.

The decision of whether to use the molecule itself or an agonist thereof for any particular indication, as opposed to an antagonist to the molecule, would depend mainly on whether the molecule herein promotes cardiovascularization, genesis of endothelial cells, or angiogenesis or inhibits these conditions. For example, if the molecule promotes angiogenesis, an antagonist thereof would be useful for treatment of disorders where it is desired to limit or prevent angiogenesis. Examples of such disorders include vascular tumors such as haemangioma, tumor angiogenesis, neovascularization in the retina, choroid, or cornea, associated with diabetic retinopathy or premature infant retinopathy or macular degeneration and proliferative vitreoretinopathy, rheumatoid arthritis, Crohn's disease, atherosclerosis, ovarian hyperstimulation, psoriasis, endometriosis associated with neovascularization, restenosis subsequent to balloon angioplasty, scar tissue overproduction, for example, that seen in a keloid that forms after surgery, fibrosis after myocardial infarction, or fibrotic lesions associated with pulmonary fibrosis.

If, however, the molecule inhibits angiogenesis, it would be expected to be used directly for treatment of the above conditions.

On the other hand, if the molecule stimulates angiogenesis it would be used itself (or an agonist thereof) for indications where angiogenesis is desired such as peripheral vascular disease, hypertension, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, arterial restenosis, thrombophlebitis, lymphangitis, lymphedema, wound healing and tissue repair, ischemia reperfusion injury, angina, myocardial infarctions such as acute myocardial infarctions, chronic heart conditions, heart failure such as congestive heart failure, and osteoporosis.

If, however, the molecule inhibits angiogenesis, an antagonist thereof would be used for treatment of those conditions where angiogenesis is desired.

Specific types of diseases are described below, where the PRO polypeptide herein or antagonists thereof may serve as useful for vascular-related drug targeting or as therapeutic targets for the treatment or prevention of the disorders. Atherosclerosis is a disease characterized by accumulation of plaques of intimal thickening in arteries, due to accumulation of lipids, proliferation of smooth muscle cells, and formation of fibrous tissue within the arterial wall. The disease can affect large, medium, and small arteries in any organ. Changes in endothelial and vascular smooth muscle cell function are known to play an important role in modulating the accumulation and

regression of these plaques.

Hypertension is characterized by raised vascular pressure in the systemic arterial, pulmonary arterial, or portal venous systems. Elevated pressure may result from or result in impaired endothelial function and/or vascular disease.

- 5        Inflammatory vasculitides include giant cell arteritis, Takayasu's arteritis, polyarteritis nodosa (including the microangiopathic form), Kawasaki's disease, microscopic polyangiitis, Wegener's granulomatosis, and a variety of infectious-related vascular disorders (including Henoch-Schonlein purpura). Altered endothelial cell function has been shown to be important in these diseases.

- 10        Reynaud's disease and Reynaud's phenomenon are characterized by intermittent abnormal impairment of the circulation through the extremities on exposure to cold. Altered endothelial cell function has been shown to be important in this disease.

Aneurysms are saccular or fusiform dilatations of the arterial or venous tree that are associated with altered endothelial cell and/or vascular smooth muscle cells.

- 15        Arterial restenosis (restenosis of the arterial wall) may occur following angioplasty as a result of alteration in the function and proliferation of endothelial and vascular smooth muscle cells.

Thrombophlebitis and lymphangitis are inflammatory disorders of veins and lymphatics, respectively, that may result from, and/or in, altered endothelial cell function. Similarly, lymphedema is a condition involving impaired lymphatic vessels resulting from endothelial cell function.

- 20        The family of benign and malignant vascular tumors are characterized by abnormal proliferation and growth of cellular elements of the vascular system. For example, lymphangiomas are benign tumors of the lymphatic system that are congenital, often cystic, malformations of the lymphatics that usually occur in newborns. Cystic tumors tend to grow into the adjacent tissue. Cystic tumors usually occur in the cervical and axillary region. They can also occur in the soft tissue of the extremities. The main symptoms are dilated, sometimes reticular, structured lymphatics and lymphocysts surrounded by connective tissue. Lymphangiomas are assumed to be caused by  
25        improperly connected embryonic lymphatics or their deficiency. The result is impaired local lymph drainage. Griener *et al.*, Lymphology, 4: 140-144 (1971).

- Another use for the PRO polypeptides herein or antagonists thereto is in the prevention of tumor angiogenesis, which involves vascularization of a tumor to enable it to grow and/or metastasize. This process is dependent on the growth of new blood vessels. Examples of neoplasms and related conditions that involve tumor angiogenesis  
30        include breast carcinomas, lung carcinomas, gastric carcinomas, esophageal carcinomas, colorectal carcinomas, liver carcinomas, ovarian carcinomas, thecomas, arrhenoblastomas, cervical carcinomas, endometrial carcinoma, endometrial hyperplasia, endometriosis, fibrosarcomas, choriocarcinoma, head and neck cancer, nasopharyngeal carcinoma, laryngeal carcinomas, hepatoblastoma, Kaposi's sarcoma, melanoma, skin carcinomas, hemangioma, cavernous hemangioma, hemangioblastoma, pancreas carcinomas, retinoblastoma, astrocytoma, glioblastoma.  
35        Schwannoma, oligodendroglioma, medulloblastoma, neuroblastomas, rhabdomyosarcoma, osteogenic sarcoma, leiomyosarcomas, urinary tract carcinomas, thyroid carcinomas, Wilms' tumor, renal cell carcinoma, prostate carcinoma, abnormal vascular proliferation associated with phakomatoses, edema (such as that associated with brain

tumors), and Meigs' syndrome.

Age-related macular degeneration (AMD) is a leading cause of severe visual loss in the elderly population. The exudative form of AMD is characterized by choroidal neovascularization and retinal pigment epithelial cell detachment. Because choroidal neovascularization is associated with a dramatic worsening in prognosis, the PRO polypeptide or antagonist thereto is expected to be useful in reducing the severity of AMD.

Healing of trauma such as wound healing and tissue repair is also a targeted use for the PRO polypeptides herein or their antagonists. Formation and regression of new blood vessels is essential for tissue healing and repair. This category includes bone, cartilage, tendon, ligament, and/or nerve tissue growth or regeneration, as well as wound healing and tissue repair and replacement, and in the treatment of burns, incisions, and ulcers. A PRO polypeptide or antagonist thereof that induces cartilage and/or bone growth in circumstances where bone is not normally formed has application in the healing of bone fractures and cartilage damage or defects in humans and other animals. Such a preparation employing a PRO polypeptide or antagonist thereof may have prophylactic use in closed as well as open fracture reduction and also in the improved fixation of artificial joints. *De novo* bone formation induced by an osteogenic agent contributes to the repair of congenital, trauma-induced, or oncologic, resection-induced craniofacial defects, and also is useful in cosmetic plastic surgery.

PRO polypeptides or antagonists thereto may also be useful to promote better or faster closure of non-healing wounds, including without limitation pressure ulcers, ulcers associated with vascular insufficiency, surgical and traumatic wounds, and the like.

It is expected that a PRO polypeptide or antagonist thereto may also exhibit activity for generation or regeneration of other tissues, such as organs (including, for example, pancreas, liver, intestine, kidney, skin, or endothelium), muscle (smooth, skeletal, or cardiac), and vascular (including vascular endothelium) tissue, or for promoting the growth of cells comprising such tissues. Part of the desired effects may be by inhibition or modulation of fibrotic scarring to allow normal tissue to regenerate.

A PRO polypeptide herein or antagonist thereto may also be useful for gut protection or regeneration and treatment of lung or liver fibrosis, reperfusion injury in various tissues, and conditions resulting from systemic cytokine damage. Also, the PRO polypeptide or antagonist thereto may be useful for promoting or inhibiting differentiation of tissues described above from precursor tissues or cells, or for inhibiting the growth of tissues described above.

A PRO polypeptide or antagonist thereto may also be used in the treatment of periodontal diseases and in other tooth-repair processes. Such agents may provide an environment to attract bone-forming cells, stimulate growth of bone-forming cells, or induce differentiation of progenitors of bone-forming cells. A PRO polypeptide herein or an antagonist thereto may also be useful in the treatment of osteoporosis or osteoarthritis, such as through stimulation of bone and/or cartilage repair or by blocking inflammation or processes of tissue destruction (collagenase activity, osteoclast activity, etc.) mediated by inflammatory processes, since blood vessels play an important role in the regulation of bone turnover and growth.

Another category of tissue regeneration activity that may be attributable to the PRO polypeptide herein or antagonist thereto is tendon/ligament formation. A protein that induces tendon/ligament-like tissue or other tissue

formation in circumstances where such tissue is not normally formed has application in the healing of tendon or ligament tears, deformities, and other tendon or ligament defects in humans and other animals. Such a preparation may have prophylactic use in preventing damage to tendon or ligament tissue, as well as use in the improved fixation of tendon or ligament to bone or other tissues, and in repairing defects to tendon or ligament tissue. *De novo* tendon/ligament-like tissue formation induced by a composition of the PRO polypeptide herein or antagonist thereto contributes to the repair of congenital, trauma-induced, or other tendon or ligament defects of other origin, and is also useful in cosmetic plastic surgery for attachment or repair of tendons or ligaments. The compositions herein may provide an environment to attract tendon- or ligament-forming cells, stimulate growth of tendon- or ligament-forming cells, induce differentiation of progenitors of tendon- or ligament-forming cells, or induce growth of tendon/ligament cells or progenitors *ex vivo* for return *in vivo* to effect tissue repair. The compositions herein may also be useful in the treatment of tendinitis, carpal tunnel syndrome, and other tendon or ligament defects. The compositions may also include an appropriate matrix and/or sequestering agent as a carrier as is well known in the art.

The PRO polypeptide or its antagonist may also be useful for proliferation of neural cells and for regeneration of nerve and brain tissue, *i.e.*, for the treatment of central and peripheral nervous system disease and neuropathies, as well as mechanical and traumatic disorders, that involve degeneration, death, or trauma to neural cells or nerve tissue. More specifically, a PRO polypeptide or its antagonist may be used in the treatment of diseases of the peripheral nervous system, such as peripheral nerve injuries, peripheral neuropathy and localized neuropathies, and central nervous system diseases, such as Alzheimer's, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome. Further conditions that may be treated in accordance with the present invention include mechanical and traumatic disorders, such as spinal cord disorders, head trauma, and cerebrovascular diseases such as stroke. Peripheral neuropathies resulting from chemotherapy or other medical therapies may also be treatable using a PRO polypeptide herein or antagonist thereto.

Ischemia-reperfusion injury is another indication. Endothelial cell dysfunction may be important in both the initiation of, and in regulation of the sequelae of events that occur following ischemia-reperfusion injury.

Rheumatoid arthritis is a further indication. Blood vessel growth and targeting of inflammatory cells through the vasculature is an important component in the pathogenesis of rheumatoid and sero-negative forms of arthritis.

A PRO polypeptide or its antagonist may also be administered prophylactically to patients with cardiac hypertrophy, to prevent the progression of the condition, and avoid sudden death, including death of asymptomatic patients. Such preventative therapy is particularly warranted in the case of patients diagnosed with massive left ventricular cardiac hypertrophy (a maximal wall thickness of 35 mm or more in adults, or a comparable value in children), or in instances when the hemodynamic burden on the heart is particularly strong.

A PRO polypeptide or its antagonist may also be useful in the management of atrial fibrillation, which develops in a substantial portion of patients diagnosed with hypertrophic cardiomyopathy.

Further indications include angina, myocardial infarctions such as acute myocardial infarctions, and heart failure such as congestive heart failure. Additional non-neoplastic conditions include psoriasis, diabetic and other proliferative retinopathies including retinopathy of prematurity, retrolental fibroplasia, neovascular glaucoma,



thyroid hyperplasias (including Grave's disease), corneal and other tissue transplantation, chronic inflammation, lung inflammation, nephrotic syndrome, preeclampsia, ascites, pericardial effusion (such as that associated with pericarditis), and pleural effusion.

5 In view of the above, the PRO polypeptides or agonists or antagonists thereof described herein, which are shown to alter or impact endothelial cell function, proliferation, and/or form, are likely to play an important role in the etiology and pathogenesis of many or all of the disorders noted above, and as such can serve as therapeutic targets to augment or inhibit these processes or for vascular-related drug targeting in these disorders.

xi. Administration Protocols, Schedules, Doses, and Formulations

10 The molecules herein and agonists and antagonists thereto are pharmaceutically useful as a prophylactic and therapeutic agent for various disorders and diseases as set forth above.

Therapeutic compositions of the PRO polypeptides or agonists or antagonists are prepared for storage by mixing the desired molecule having the appropriate degree of purity with optional pharmaceutically acceptable carriers, excipients, or stabilizers (Remington's Pharmaceutical Sciences, 16th edition, Osol, A. ed. (1980)), in the form of lyophilized formulations or aqueous solutions. Acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrins; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (*e.g.*, Zn-protein complexes); and/or non-ionic surfactants such as TWEEN<sup>TM</sup>, PLURONICS<sup>TM</sup> or polyethylene glycol (PEG).

25 Additional examples of such carriers include ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts, or electrolytes such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, and polyethylene glycol. Carriers for topical or gel-based forms of antagonist include polysaccharides such as sodium carboxymethylcellulose or methylcellulose, polyvinylpyrrolidone, polyacrylates, polyoxyethylene-polyoxypropylene-block polymers, polyethylene glycol, and wood wax alcohols. For all administrations, conventional depot forms are suitably used. Such forms include, for example, microcapsules, nano-capsules, liposomes, plasters, inhalation forms, nose sprays, sublingual tablets, and sustained-release preparations. The PRO polypeptides or agonists or antagonists will typically be formulated in such vehicles at a concentration of about 0.1 mg/ml to 100 mg/ml.

35 Another formulation comprises incorporating a PRO polypeptide or antagonist thereof into formed articles.

Such articles can be used in modulating endothelial cell growth and angiogenesis. In addition, tumor invasion and metastasis may be modulated with these articles.

PRO polypeptide or antagonist to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes, prior to or following lyophilization and reconstitution. PRO polypeptide ordinarily will be stored in lyophilized form or in solution if administered systemically. If in lyophilized form, PRO polypeptide or antagonist thereto is typically formulated in combination with other ingredients for reconstitution with an appropriate diluent at the time for use. An example of a liquid formulation of PRO polypeptide or antagonist is a sterile, clear, colorless unpreserved solution filled in a single-dose vial for subcutaneous injection. Preserved pharmaceutical compositions suitable for repeated use may contain, for example, depending mainly on the indication and type of polypeptide:

- a) PRO polypeptide or agonist or antagonist thereto;
- b) a buffer capable of maintaining the pH in a range of maximum stability of the polypeptide or other molecule in solution, preferably about 4-8;
- c) a detergent/surfactant primarily to stabilize the polypeptide or molecule against agitation-induced aggregation;
- d) an isotonifier;
- e) a preservative selected from the group of phenol, benzyl alcohol and a benzethonium halide, *e.g.*, chloride; and
- f) water.

If the detergent employed is non-ionic, it may, for example, be polysorbates (*e.g.*, POLYSORBATE™ (TWEEN™) 20, 80, etc.) or poloxamers (*e.g.*, POLOXAMER™ 188). The use of non-ionic surfactants permits the formulation to be exposed to shear surface stresses without causing denaturation of the polypeptide. Further, such surfactant-containing formulations may be employed in aerosol devices such as those used in a pulmonary dosing, and needleless jet injector guns (*see, e.g.*, EP 257,956).

An isotonifier may be present to ensure isotonicity of a liquid composition of the PRO polypeptide or antagonist thereto, and includes polyhydric sugar alcohols, preferably trihydric or higher sugar alcohols, such as glycerin, erythritol, arabitol, xylitol, sorbitol, and mannitol. These sugar alcohols can be used alone or in combination. Alternatively, sodium chloride or other appropriate inorganic salts may be used to render the solutions isotonic.

The buffer may, for example, be an acetate, citrate, succinate, or phosphate buffer depending on the pH desired. The pH of one type of liquid formulation of this invention is buffered in the range of about 4 to 8, preferably about physiological pH.

The preservatives phenol, benzyl alcohol and benzethonium halides, *e.g.*, chloride, are known antimicrobial agents that may be employed.

Therapeutic PRO polypeptide compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle. The formulations are preferably administered as repeated intravenous (*i.v.*), subcutaneous (*s.c.*), or intramuscular (*i.m.*)

injections, or as aerosol formulations suitable for intranasal or intrapulmonary delivery (for intrapulmonary delivery see, e.g., EP 257,956).

PRO polypeptide can also be administered in the form of sustained-released preparations. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the protein, which matrices are in the form of shaped articles, e.g., films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (e.g., poly(2-hydroxyethyl-methacrylate) as described by Langer *et al.*, J. Biomed. Mater. Res., **15**: 167-277 (1981) and Langer, Chem. Tech., **12**: 98-105 (1982) or poly(vinylalcohol)), polylactides (U.S. Patent No. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma ethyl-L-glutamate (Sidman *et al.*, Biopolymers, **22**: 547-556 (1983)), non-degradable ethylene-vinyl acetate (Langer *et al.*, *supra*),  
degradable lactic acid-glycolic acid copolymers such as the Lupron Depot™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid (EP 133,988).

While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated proteins remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37°C, resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for protein stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S-S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

Sustained-release PRO polypeptide compositions also include liposomally entrapped PRO polypeptides. Liposomes containing the PRO polypeptide are prepared by methods known *per se*: DE 3,218,121; Epstein *et al.*, Proc. Natl. Acad. Sci. USA, **82**: 3688-3692 (1985); Hwang *et al.*, Proc. Natl. Acad. Sci. USA, **77**: 4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese patent application 83-118008; U.S. Patent Nos. 4,485,045 and 4,544,545; and EP 102,324. Ordinarily the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. % cholesterol, the selected proportion being adjusted for the optimal therapy.

The therapeutically effective dose of PRO polypeptide or antagonist thereto will, of course, vary depending on such factors as the pathological condition to be treated (including prevention), the method of administration, the type of compound being used for treatment, any co-therapy involved, the patient's age, weight, general medical condition, medical history, etc., and its determination is well within the skill of a practicing physician. Accordingly, it will be necessary for the therapist to titer the dosage and modify the route of administration as required to obtain the maximal therapeutic effect. If the PRO polypeptide has a narrow host range, for the treatment of human patients formulations comprising human PRO polypeptide, more preferably native-sequence human PRO polypeptide, are preferred. The clinician will administer PRO polypeptide until a dosage is reached that achieves the desired effect for treatment of the condition in question. For example, if the objective is the treatment of CHF, the amount would be one that inhibits the progressive cardiac hypertrophy associated with this condition. The progress of this therapy is easily monitored by echo cardiography. Similarly, in patients with hypertrophic cardiomyopathy, PRO

polypeptide can be administered on an empirical basis.

With the above guidelines, the effective dose generally is within the range of from about 0.001 to about 1.0 mg/kg, more preferably about 0.01-1.0 mg/kg, most preferably about 0.01-0.1 mg/kg.

For non-oral use in treating human adult hypertension, it is advantageous to administer PRO polypeptide in the form of an injection at about 0.01 to 50 mg, preferably about 0.05 to 20 mg, most preferably 1 to 20 mg, per kg body weight, 1 to 3 times daily by intravenous injection. For oral administration, a molecule based on the PRO polypeptide is preferably administered at about 5 mg to 1 g, preferably about 10 to 100 mg, per kg body weight, 1 to 3 times daily. It should be appreciated that endotoxin contamination should be kept minimally at a safe level, for example, less than 0.5 ng/mg protein. Moreover, for human administration, the formulations preferably meet sterility, pyrogenicity, general safety, and purity as required by FDA Office and Biologics standards.

The dosage regimen of a pharmaceutical composition containing PRO polypeptide to be used in tissue regeneration will be determined by the attending physician considering various factors that modify the action of the polypeptides, *e.g.*, amount of tissue weight desired to be formed, the site of damage, the condition of the damaged tissue, the size of a wound, type of damaged tissue (*e.g.*, bone), the patient's age, sex, and diet, the severity of any infection, time of administration, and other clinical factors. The dosage may vary with the type of matrix used in the reconstitution and with inclusion of other proteins in the pharmaceutical composition. For example, the addition of other known growth factors, such as IGF-I, to the final composition may also affect the dosage. Progress can be monitored by periodic assessment of tissue/bone growth and/or repair, for example, X-rays, histomorphometric determinations, and tetracycline labeling.

The route of PRO polypeptide or antagonist or agonist administration is in accord with known methods, *e.g.*, by injection or infusion by intravenous, intramuscular, intracerebral, intraperitoneal, intracerebrospinal, subcutaneous, intraocular, intraarticular, intrasynovial, intrathecal, oral, topical, or inhalation routes, or by sustained-release systems as noted below. The PRO polypeptide or antagonists thereof also are suitably administered by intratumoral, peritumoral, intralesional, or perilesional routes, to exert local as well as systemic therapeutic effects. The intraperitoneal route is expected to be particularly useful, for example, in the treatment of ovarian tumors.

If a peptide or small molecule is employed as an antagonist or agonist, it is preferably administered orally or non-orally in the form of a liquid or solid to mammals.

Examples of pharmacologically acceptable salts of molecules that form salts and are useful hereunder include alkali metal salts (*e.g.*, sodium salt, potassium salt), alkaline earth metal salts (*e.g.*, calcium salt, magnesium salt), ammonium salts, organic base salts (*e.g.*, pyridine salt, triethylamine salt), inorganic acid salts (*e.g.*, hydrochloride, sulfate, nitrate), and salts of organic acid (*e.g.*, acetate, oxalate, p-toluenesulfonate).

For compositions herein that are useful for bone, cartilage, tendon, or ligament regeneration, the therapeutic method includes administering the composition topically, systemically, or locally as an implant or device. When administered, the therapeutic composition for use is in a pyrogen-free, physiologically acceptable form. Further, the composition may desirably be encapsulated or injected in a viscous form for delivery to the site of bone, cartilage, or tissue damage. Topical administration may be suitable for wound healing and tissue repair. Preferably,

for bone and/or cartilage formation, the composition would include a matrix capable of delivering the protein-containing composition to the site of bone and/or cartilage damage, providing a structure for the developing bone and cartilage and preferably capable of being resorbed into the body. Such matrices may be formed of materials presently in use for other implanted medical applications.

5 The choice of matrix material is based on biocompatibility, biodegradability, mechanical properties, cosmetic appearance, and interface properties. The particular application of the compositions will define the appropriate formulation. Potential matrices for the compositions may be biodegradable and chemically defined calcium sulfate, tricalcium phosphate, hydroxyapatite, polylactic acid, polyglycolic acid, and polyanhydrides. Other potential materials are biodegradable and biologically well-defined, such as bone or dermal collagen. Further matrices are  
10 comprised of pure proteins or extracellular matrix components. Other potential matrices are nonbiodegradable and chemically defined, such as sintered hydroxyapatite, bioglass, aluminates, or other ceramics. Matrices may be comprised of combinations of any of the above-mentioned types of material, such as polylactic acid and hydroxyapatite or collagen and tricalcium phosphate. The bioceramics may be altered in composition, such as in calcium-aluminate-phosphate and processing to alter pore size, particle size, particle shape, and biodegradability.

15 One specific embodiment is a 50:50 (mole weight) copolymer of lactic acid and glycolic acid in the form of porous particles having diameters ranging from 150 to 800 microns. In some applications, it will be useful to utilize a sequestering agent, such as carboxymethyl cellulose or autologous blood clot, to prevent the polypeptide compositions from disassociating from the matrix.

One suitable family of sequestering agents is cellulosic materials such as alkylcelluloses (including  
20 hydroxyalkylcelluloses), including methylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose, and carboxymethylcellulose, one preferred being cationic salts of carboxymethylcellulose (CMC). Other preferred sequestering agents include hyaluronic acid, sodium alginate, poly(ethylene glycol), polyoxyethylene oxide, carboxyvinyl polymer, and poly(vinyl alcohol). The amount of sequestering agent useful herein is 0.5-20 wt%, preferably 1-10 wt%, based on total formulation weight, which  
25 represents the amount necessary to prevent desorption of the polypeptide (or its antagonist) from the polymer matrix and to provide appropriate handling of the composition, yet not so much that the progenitor cells are prevented from infiltrating the matrix, thereby providing the polypeptide (or its antagonist) the opportunity to assist the osteogenic activity of the progenitor cells.

#### xii. Combination Therapies

30 The effectiveness of the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide or an agonist or antagonist thereof in preventing or treating the disorder in question may be improved by administering the active agent serially or in combination with another  
35 agent that is effective for those purposes, either in the same composition or as separate compositions.

For example, for treatment of cardiac hypertrophy, PRO polypeptide therapy can be combined with the

administration of inhibitors of known cardiac myocyte hypertrophy factors, *e.g.*, inhibitors of  $\alpha$ -adrenergic agonists such as phenylephrine; endothelin-1 inhibitors such as BOSENTAN<sup>TM</sup> and MOXONODIN<sup>TM</sup>; inhibitors to CT-1 (US Pat. No. 5,679,545); inhibitors to LIF; ACE inhibitors; des-aspartate-angiotensin I inhibitors (U.S. Pat. No. 5,773,415), and angiotensin II inhibitors.

5 For treatment of cardiac hypertrophy associated with hypertension, the PRO polypeptide can be administered in combination with  $\beta$ -adrenergic receptor blocking agents, *e.g.*, propranolol, timolol, tertalolol, carteolol, nadolol, betaxolol, penbutolol, acetobutolol, atenolol, metoprolol, or carvedilol; ACE inhibitors, *e.g.*, quinapril, captopril, enalapril, ramipril, benazepril, fosinopril, or lisinopril; diuretics, *e.g.*, chlorothiazide, hydrochlorothiazide, hydroflumethazide, methylchlorothiazide, benzthiazide, dichlorphenamide, acetazolamide, or indapamide; and/or  
10 calcium channel blockers, *e.g.*, diltiazem, nifedipine, verapamil, or nicardipine. Pharmaceutical compositions comprising the therapeutic agents identified herein by their generic names are commercially available, and are to be administered following the manufacturers' instructions for dosage, administration, adverse effects, contraindications, etc. *See, e.g.*, Physicians' Desk Reference (Medical Economics Data Production Co.: Montvale, N.J., 1997), 51th Edition.

15 Preferred candidates for combination therapy in the treatment of hypertrophic cardiomyopathy are  $\beta$ -adrenergic-blocking drugs (*e.g.*, propranolol, timolol, tertalolol, carteolol, nadolol, betaxolol, penbutolol, acetobutolol, atenolol, metoprolol, or carvedilol), verapamil, diltiazem, or diltiazem. Treatment of hypertrophy associated with high blood pressure may require the use of antihypertensive drug therapy, using calcium channel blockers, *e.g.*, diltiazem, nifedipine, verapamil, or nicardipine;  $\beta$ -adrenergic blocking agents; diuretics, *e.g.*,  
20 chlorothiazide, hydrochlorothiazide, hydroflumethazide, methylchlorothiazide, benzthiazide, dichlorphenamide, acetazolamide, or indapamide; and/or ACE-inhibitors, *e.g.*, quinapril, captopril, enalapril, ramipril, benazepril, fosinopril, or lisinopril.

For other indications, PRO polypeptides or their antagonists may be combined with other agents beneficial to the treatment of the bone and/or cartilage defect, wound, or tissue in question. These agents include various  
25 growth factors such as EGF, PDGF, TGF- $\alpha$  or TGF- $\beta$ , IGF, FGF, and CTGF.

In addition, PRO polypeptides or their antagonists used to treat cancer may be combined with cytotoxic, chemotherapeutic, or growth-inhibitory agents as identified above. Also, for cancer treatment, the PRO polypeptide or antagonist thereof is suitably administered serially or in combination with radiological treatments, whether involving irradiation or administration of radioactive substances.

30 The effective amounts of the therapeutic agents administered in combination with the PRO polypeptide or antagonist thereof will be at the physician's or veterinarian's discretion. Dosage administration and adjustment is done to achieve maximal management of the conditions to be treated. For example, for treating hypertension, these amounts ideally take into account use of diuretics or digitalis, and conditions such as hyper- or hypotension, renal impairment, etc. The dose will additionally depend on such factors as the type of the therapeutic agent to be used  
35 and the specific patient being treated. Typically, the amount employed will be the same dose as that used, if the given therapeutic agent is administered without the PRO polypeptide.

xiii. Articles of Manufacture

An article of manufacture such as a kit containing PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide or agonists or antagonists thereof useful for the diagnosis or treatment of the disorders described above comprises at least a container and a label. Suitable containers include, for example, bottles, vials, syringes, and test tubes. The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition that is effective for diagnosing or treating the condition and may have a sterile access port (for example, the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). The active agent in the composition is the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide or an agonist or antagonist thereto. The label on, or associated with, the container indicates that the composition is used for diagnosing or treating the condition of choice. The article of manufacture may further comprise a second container comprising a pharmaceutically-acceptable buffer, such as phosphate-buffered saline, Ringer's solution, and dextrose solution. It may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, syringes, and package inserts with instructions for use. The article of manufacture may also comprise a second or third container with another active agent as described above.

E. Antibodies

Some of the most promising drug candidates according to the present invention are antibodies and antibody fragments that may inhibit the production or the gene product of the genes identified herein and/or reduce the activity of the gene products.

i. Polyclonal Antibodies

Methods of preparing polyclonal antibodies are known to the skilled artisan. Polyclonal antibodies can be raised in a mammal, for example, by one or more injections of an immunizing agent and, if desired, an adjuvant. Typically, the immunizing agent and/or adjuvant will be injected in the mammal by multiple subcutaneous or intraperitoneal injections. The immunizing agent may include the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide or a fusion protein thereof. It may be useful to conjugate the immunizing agent to a protein known to be immunogenic in the mammal being immunized. Examples of such immunogenic proteins include, but are not limited to, keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. Examples of adjuvants that may be employed

include Freund's complete adjuvant and MPL-TDM adjuvant (monophosphoryl Lipid A or synthetic trehalose dicorynomycolate). The immunization protocol may be selected by one skilled in the art without undue experimentation.

ii. Monoclonal Antibodies

5 The anti-PRO172, anti-PRO175, anti-PRO178, anti-PRO188, anti-PRO356, anti-PRO179, anti-PRO197, anti-PRO198, anti-PRO182, anti-PRO195, anti-PRO200, anti-PRO211, anti-PRO217, anti-PRO219, anti-PRO221, anti-PRO224, anti-PRO228, anti-PRO245, anti-PRO246, anti-PRO258, anti-PRO261, anti-PRO272, anti-PRO301, anti-PRO322, anti-PRO328, anti-PRO331, anti-PRO364, anti-PRO366, anti-PRO535, anti-PRO819, anti-PRO826, anti-PRO1160, anti-PRO1186 or anti-PRO1246 antibodies may, alternatively, be monoclonal antibodies. Monoclonal  
10 antibodies may be prepared using hybridoma methods, such as those described by Kohler and Milstein, Nature, 256:495 (1975). In a hybridoma method, a mouse, hamster, or other appropriate host animal is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes may be immunized *in vitro*.

The immunizing agent will typically include the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide or a fusion protein thereof. Generally, either peripheral blood lymphocytes ("PBLs") are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an  
15 immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell. Goding, Monoclonal Antibodies: Principles and Practice (New York: Academic Press, 1986), pp. 59-103. Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of rodent, bovine, and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells may be cultured in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused,  
20 immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells.

Preferred immortalized cell lines are those that fuse efficiently, support stable high-level expression of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. More  
30 preferred immortalized cell lines are murine myeloma lines, which can be obtained, for instance, from the Salk Institute Cell Distribution Center, San Diego, California and the American Type Culture Collection, Manassas, Virginia. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies. Kozbor, J. Immunol., 133:3001 (1984); Brodeur *et al.*, Monoclonal Antibody Production Techniques and Applications (Marcel Dekker, Inc.: New York, 1987) pp. 51-63.

35 The culture medium in which the hybridoma cells are cultured can then be assayed for the presence of monoclonal antibodies directed against the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197,



PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide. Preferably, the binding specificity of monoclonal antibodies produced by the hybridoma cells is determined by immunoprecipitation or by an *in vitro* binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA). Such techniques and assays are known in the art. The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson and Pollard, Anal. Biochem., 107:220 (1980).

After the desired hybridoma cells are identified, the clones may be subcloned by limiting dilution procedures and grown by standard methods. Goding, *supra*. Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium and RPMI-1640 medium. Alternatively, the hybridoma cells may be grown *in vivo* as ascites in a mammal.

The monoclonal antibodies secreted by the subclones may be isolated or purified from the culture medium or ascites fluid by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

The monoclonal antibodies may also be made by recombinant DNA methods, such as those described in U.S. Patent No. 4,816,567. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (*e.g.*, by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells of the invention serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also may be modified, for example, by substituting the coding sequence for human heavy- and light-chain constant domains in place of the homologous murine sequences (U.S. Patent No. 4,816,567; Morrison *et al.*, *supra*) or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. Such a non-immunoglobulin polypeptide can be substituted for the constant domains of an antibody of the invention, or can be substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody.

The antibodies may be monovalent antibodies. Methods for preparing monovalent antibodies are well known in the art. For example, one method involves recombinant expression of immunoglobulin light chain and modified heavy chain. The heavy chain is truncated generally at any point in the Fc region so as to prevent heavy-chain crosslinking. Alternatively, the relevant cysteine residues are substituted with another amino acid residue or are deleted so as to prevent crosslinking.

*In vitro* methods are also suitable for preparing monovalent antibodies. Digestion of antibodies to produce fragments thereof, particularly Fab fragments, can be accomplished using routine techniques known in the art.

### iii. Human and Humanized Antibodies

The anti-PRO172, anti-PRO175, anti-PRO178, anti-PRO188, anti-PRO356, anti-PRO179, anti-PRO197, anti-

PRO198, anti-PRO182, anti-PRO195, anti-PRO200, anti-PRO211, anti-PRO217, anti-PRO219, anti-PRO221, anti-PRO224, anti-PRO228, anti-PRO245, anti-PRO246, anti-PRO258, anti-PRO261, anti-PRO272, anti-PRO301, anti-PRO322, anti-PRO328, anti-PRO331, anti-PRO364, anti-PRO366, anti-PRO535, anti-PRO819, anti-PRO826, anti-PRO1160, anti-PRO1186 or anti-PRO1246 antibodies may further comprise humanized antibodies or human antibodies. Humanized forms of non-human (*e.g.*, murine) antibodies are chimeric immunoglobulins, immunoglobulin chains, or fragments thereof (such as Fv, Fab, Fab', F(ab')<sub>2</sub>, or other antigen-binding subsequences of antibodies) that contain minimal sequence derived from non-human immunoglobulin. Humanized antibodies include human immunoglobulins (recipient antibody) in which residues from a CDR of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat, or rabbit having the desired specificity, affinity, and capacity. In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies may also comprise residues that are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin, and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody preferably also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. Jones *et al.*, Nature, 321: 522-525 (1986); Riechmann *et al.*, Nature, 332: 323-329 (1988); Presta, Curr. Op. Struct. Biol., 2:593-596 (1992).

Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source that is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers (Jones *et al.*, Nature, 321: 522-525 (1986); Riechmann *et al.*, Nature, 332: 323-327 (1988); Verhoeven *et al.*, Science, 239: 1534-1536 (1988)), by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Patent No. 4,816,567), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

Human antibodies can also be produced using various techniques known in the art, including phage display libraries. Hoogenboom and Winter, J. Mol. Biol., 227: 381 (1991); Marks *et al.*, J. Mol. Biol., 222: 581 (1991). The techniques of Cole *et al.* and Boerner *et al.* are also available for the preparation of human monoclonal antibodies. Cole *et al.*, Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, p. 77 (1985) and Boerner *et al.*, J. Immunol., 147(1): 86-95 (1991). Similarly, human antibodies can be made by introducing human immunoglobulin loci into transgenic animals, *e.g.*, mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed that closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach is described, for example, in U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; and

5,661,016, and in the following scientific publications: Marks *et al.*, Bio/Technology, 10: 779-783 (1992); Lonberg *et al.*, Nature, 368: 856-859 (1994); Morrison, Nature, 368: 812-813 (1994); Fishwild *et al.*, Nature Biotechnology, 14: 845-851 (1996); Neuberger, Nature Biotechnology, 14: 826 (1996); Lonberg and Huszar, Intern. Rev. Immunol., 13: 65-93 (1995).

5           iv.       Bispecific Antibodies

Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, 10 PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide, the other one is for any other antigen, and preferably for a cell-surface protein or receptor or receptor subunit.

Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy-chain/light-chain pairs, where 15 the two heavy chains have different specificities. Milstein and Cuello, Nature, 305: 537-539 (1983). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of ten different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule is usually accomplished by affinity chromatography steps. Similar procedures are disclosed in WO 93/08829, published 13 May 1993, and in Traunecker *et al.*, EMBO J., 10: 3655-3659 (1991).

20       Antibody variable domains with the desired binding specificities (antibody-antigen combining sites) can be fused to immunoglobulin constant-domain sequences. The fusion preferably is with an immunoglobulin heavy-chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light-chain binding present in at least one of the fusions. DNAs encoding the immunoglobulin heavy-chain fusions and, if desired, the immunoglobulin light 25 chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. For further details of generating bispecific antibodies, *see*, for example, Suresh *et al.*, Methods in Enzymology, 121: 210 (1986).

v.       Heteroconjugate Antibodies

Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for 30 example, been proposed to target immune-system cells to unwanted cells (U.S. Patent No. 4,676,980), and for treatment of HIV infection. WO 91/00360; WO 92/200373; EP 03089. It is contemplated that the antibodies may be prepared *in vitro* using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins may be constructed using a disulfide-exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4- 35 mercaptobutyrimidate and those disclosed, for example, in U.S. Patent No. 4,676,980.

vi. Effector Function Engineering

It may be desirable to modify the antibody of the invention with respect to effector function, so as to enhance, e.g., the effectiveness of the antibody in treating cancer. For example, cysteine residue(s) may be introduced into the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated may have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). See, Caron *et al.*, J. Exp. Med., 176: 1191-1195 (1992) and Shopes, J. Immunol., 148: 2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor activity may also be prepared using heterobifunctional cross-linkers as described in Wolff *et al.*, Cancer Research, 53: 2560-2565 (1993). Alternatively, an antibody can be engineered that has dual Fc regions and may thereby have enhanced complement lysis and ADCC capabilities. See, Stevenson *et al.*, Anti-Cancer Drug Design, 3: 219-230 (1989).

vii. Immunoconjugates

The invention also pertains to immunoconjugates comprising an antibody conjugated to a cytotoxic agent such as a chemotherapeutic agent, toxin (e.g., an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (i.e., a radioconjugate).

Chemotherapeutic agents useful in the generation of such immunoconjugates have been described above. Enzymatically active toxins and fragments thereof that can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolaca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, croton, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes. A variety of radionuclides are available for the production of radioconjugated antibodies. Examples include  $^{212}\text{Bi}$ ,  $^{131}\text{I}$ ,  $^{131}\text{In}$ ,  $^{90}\text{Y}$ , and  $^{186}\text{Re}$ .

Conjugates of the antibody and cytotoxic agent are made using a variety of bifunctional protein-coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCl), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta *et al.*, Science, 238: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See, WO94/11026.

In another embodiment, the antibody may be conjugated to a "receptor" (such as streptavidin) for utilization in tumor pretargeting wherein the antibody-receptor conjugate is administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (e.g., avidin) that is conjugated to a cytotoxic agent (e.g., a radionucleotide).

viii. Immunoliposomes

The antibodies disclosed herein may also be formulated as immunoliposomes. Liposomes containing the antibody are prepared by methods known in the art, such as described in Epstein *et al.*, Proc. Natl. Acad. Sci. USA, 82: 3688 (1985); Hwang *et al.*, Proc. Natl. Acad. Sci. USA, 77: 4030 (1980); and U.S. Pat. Nos. 4,485,045 and 4,544,545. Liposomes with enhanced circulation time are disclosed in U.S. Patent No. 5,013,556.

Particularly useful liposomes can be generated by the reverse-phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol, and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of the antibody of the present invention can be conjugated to the liposomes as described in Martin *et al.*, J. Biol. Chem., 257: 286-288 (1982) via a disulfide-interchange reaction. A chemotherapeutic agent (such as Doxorubicin) is optionally contained within the liposome. See, Gabizon *et al.*, J. National Cancer Inst., 81(19): 1484 (1989).

ix. Pharmaceutical Compositions of Antibodies

Antibodies specifically binding a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide identified herein, as well as other molecules identified by the screening assays disclosed hereinbefore, can be administered for the treatment of various disorders as noted above and below in the form of pharmaceutical compositions.

If the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide is intracellular and whole antibodies are used as inhibitors, internalizing antibodies are preferred. However, lipofections or liposomes can also be used to deliver the antibody, or an antibody fragment, into cells. Where antibody fragments are used, the smallest inhibitory fragment that specifically binds to the binding domain of the target protein is preferred. For example, based upon the variable-region sequences of an antibody, peptide molecules can be designed that retain the ability to bind the target protein sequence. Such peptides can be synthesized chemically and/or produced by recombinant DNA technology. See, *e.g.*, Marasco *et al.*, Proc. Natl. Acad. Sci. USA, 90: 7889-7893 (1993).

The formulation herein may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. Alternatively, or in addition, the composition may comprise an agent that enhances its function, such as, for example, a cytotoxic agent, cytokine, chemotherapeutic agent, or growth-inhibitory agent. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

The active ingredients may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-

(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles, and nanocapsules) or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences, *supra*.

5 The formulations to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, *e.g.*, films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919),  
10 copolymers of L-glutamic acid and  $\gamma$  ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT<sup>TM</sup> (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated antibodies remain in the body for a long time, they  
15 may denature or aggregate as a result of exposure to moisture at 37°C, resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S-S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific  
20 polymer matrix compositions.

x. Methods of Treatment using the Antibody

It is contemplated that the antibodies to a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535,  
25 PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide may be used to treat various cardiovascular, endothelial, and angiogenic conditions as noted above.

The antibodies are administered to a mammal, preferably a human, in accord with known methods, such as intravenous administration as a bolus or by continuous infusion over a period of time, by intramuscular, intraperitoneal, intracerebrospinal, subcutaneous, intra-articular, intrasynovial, intrathecal, oral, topical, or  
30 inhalation routes. Intravenous administration of the antibody is preferred.

Other therapeutic regimens may be combined with the administration of the antibodies of the instant invention as noted above. For example, if the antibodies are to treat cancer, the patient to be treated with such antibodies may also receive radiation therapy. Alternatively, or in addition, a chemotherapeutic agent may be administered to the patient. Preparation and dosing schedules for such chemotherapeutic agents may be used according to  
35 manufacturers' instructions or as determined empirically by the skilled practitioner. Preparation and dosing schedules for such chemotherapy are also described in Chemotherapy Service, Ed., M.C. Perry (Williams &

Wilkins: Baltimore, MD, 1992). The chemotherapeutic agent may precede, or follow administration of the antibody, or may be given simultaneously therewith. The antibody may be combined with an anti-estrogen compound such as tamoxifen or EVISTA<sup>TM</sup> or an anti-progesterone such as onapristone (*see*, EP 616812) in dosages known for such molecules.

- 5 If the antibodies are used for treating cancer, it may be desirable also to administer antibodies against other tumor-associated antigens, such as antibodies that bind to one or more of the ErbB2, EGFR, ErbB3, ErbB4, or VEGF receptor(s). These also include the agents set forth above. Also, the antibody is suitably administered serially or in combination with radiological treatments, whether involving irradiation or administration of radioactive substances. Alternatively, or in addition, two or more antibodies binding the same or two or more  
10 different antigens disclosed herein may be co-administered to the patient. Sometimes, it may be beneficial also to administer one or more cytokines to the patient. In a preferred embodiment, the antibodies herein are co-administered with a growth-inhibitory agent. For example, the growth-inhibitory agent may be administered first, followed by an antibody of the present invention. However, simultaneous administration or administration of the antibody of the present invention first is also contemplated. Suitable dosages for the growth-inhibitory agent are  
15 those presently used and may be lowered due to the combined action (synergy) of the growth-inhibitory agent and the antibody herein.

- In one embodiment, vascularization of tumors is attacked in combination therapy. The anti-PRO polypeptide antibody and another antibody (*e.g.*, anti-VEGF) are administered to tumor-bearing patients at therapeutically effective doses as determined, for example, by observing necrosis of the tumor or its metastatic foci, if any. This  
20 therapy is continued until such time as no further beneficial effect is observed or clinical examination shows no trace of the tumor or any metastatic foci. Then TNF is administered, alone or in combination with an auxiliary agent such as alpha-, beta-, or gamma-interferon, anti-HER2 antibody, heregulin, anti-heregulin antibody, D-factor, interleukin-1 (IL-1), interleukin-2 (IL-2), granulocyte-macrophage colony stimulating factor (GM-CSF), or agents that promote microvascular coagulation in tumors, such as anti-protein C antibody, anti-protein S antibody, or C4b  
25 binding protein (*see*, WO 91/01753, published 21 February 1991), or heat or radiation.

- Since the auxiliary agents will vary in their effectiveness, it is desirable to compare their impact on the tumor by matrix screening in conventional fashion. The administration of anti-PRO polypeptide antibody and TNF is repeated until the desired clinical effect is achieved. Alternatively, the anti-PRO polypeptide antibody is administered together with TNF and, optionally, auxiliary agent(s). In instances where solid tumors are found in  
30 the limbs or in other locations susceptible to isolation from the general circulation, the therapeutic agents described herein are administered to the isolated tumor or organ. In other embodiments, a FGF or PDGF antagonist, such as an anti-FGF or an anti-PDGF neutralizing antibody, is administered to the patient in conjunction with the anti-PRO polypeptide antibody. Treatment with anti-PRO polypeptide antibodies preferably may be suspended during periods of wound healing or desirable neovascularization.

- 35 For the prevention or treatment of cardiovascular, endothelial, and angiogenic disorder, the appropriate dosage of an antibody herein will depend on the type of disorder to be treated, as defined above, the severity and course of the disease, whether the antibody is administered for preventive or therapeutic purposes, previous therapy, the

patient's clinical history and response to the antibody, and the discretion of the attending physician. The antibody is suitably administered to the patient at one time or over a series of treatments.

For example, depending on the type and severity of the disorder, about 1  $\mu\text{g/kg}$  to 50 mg/kg (*e.g.*, 0.1-20 mg/kg) of antibody is an initial candidate dosage for administration to the patient, whether, for example, by one or more separate administrations, or by continuous infusion. A typical daily or weekly dosage might range from about 1  $\mu\text{g/kg}$  to 100 mg/kg or more, depending on the factors mentioned above. For repeated administrations over several days or longer, depending on the condition, the treatment is repeated or sustained until a desired suppression of disorder symptoms occurs. However, other dosage regimens may be useful. The progress of this therapy is easily monitored by conventional techniques and assays, including, for example, radiographic tumor imaging.

#### xi. Articles of Manufacture with Antibodies

An article of manufacture containing a container with the antibody and a label is also provided. Such articles are described above, wherein the active agent is an anti-PRO172, anti-PRO175, anti-PRO178, anti-PRO188, anti-PRO356, anti-PRO179, anti-PRO197, anti-PRO198, anti-PRO182, anti-PRO195, anti-PRO200, anti-PRO211, anti-PRO217, anti-PRO219, anti-PRO221, anti-PRO224, anti-PRO228, anti-PRO245, anti-PRO246, anti-PRO258, anti-PRO261, anti-PRO272, anti-PRO301, anti-PRO322, anti-PRO328, anti-PRO331, anti-PRO364, anti-PRO366, anti-PRO535, anti-PRO819, anti-PRO826, anti-PRO1160, anti-PRO1186 or anti-PRO1246 antibody.

#### xii. Diagnosis and Prognosis of Tumors using Antibodies

If the indication for which the antibodies are used is cancer, while cell-surface proteins, such as growth receptors over expressed in certain tumors, are excellent targets for drug candidates or tumor (*e.g.*, cancer) treatment, the same proteins along with PRO polypeptides find additional use in the diagnosis and prognosis of tumors. For example, antibodies directed against the PRO polypeptides may be used as tumor diagnostics or prognostics.

For example, antibodies, including antibody fragments, can be used qualitatively or quantitatively to detect the expression of genes including the gene encoding the PRO polypeptide. The antibody preferably is equipped with a detectable, *e.g.*, fluorescent label, and binding can be monitored by light microscopy, flow cytometry, fluorimetry, or other techniques known in the art. Such binding assays are performed essentially as described above.

*In situ* detection of antibody binding to the marker gene products can be performed, for example, by immunofluorescence or immunoelectron microscopy. For this purpose, a histological specimen is removed from the patient, and a labeled antibody is applied to it, preferably by overlaying the antibody on a biological sample. This procedure also allows for determining the distribution of the marker gene product in the tissue examined. It will be apparent to those skilled in the art that a wide variety of histological methods are readily available for *in situ* detection.

The following Examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.



The disclosures of all patent and literature references cited in the present specification are hereby incorporated by reference in their entirety.

### EXAMPLES

Commercially available reagents referred to in the Examples were used according to manufacturer's instructions unless otherwise indicated. The source of those cells identified in the following Examples, and throughout the specification, by ATCC accession numbers is the American Type Culture Collection, Manassas, VA. Unless otherwise noted, the present invention uses standard procedures of recombinant DNA technology, such as those described hereinabove and in the following textbooks: Sambrook *et al.*, *supra*; Ausubel *et al.*, Current Protocols in Molecular Biology (Green Publishing Associates and Wiley Interscience, N.Y., 1989); Innis *et al.*, PCR Protocols: A Guide to Methods and Applications (Academic Press, Inc.: N.Y., 1990); Harlow *et al.*, Antibodies: A Laboratory Manual (Cold Spring Harbor Press: Cold Spring Harbor, 1988); Gait, Oligonucleotide Synthesis (IRL Press: Oxford, 1984); Freshney, Animal Cell Culture, 1987; Coligan *et al.*, Current Protocols in Immunology, 1991.

#### EXAMPLE 1

##### Extracellular Domain Homology Screening to Identify Novel Polypeptides and cDNA Encoding Therefor

The extracellular domain (ECD) sequences (including the secretion signal sequence, if any) from about 950 known secreted proteins from the Swiss-Prot public database were used to search EST databases. The EST databases included public databases (*e.g.*, GenBank), and proprietary databases (*e.g.*, LIFESEQ®, Incyte Pharmaceuticals, Palo Alto, CA). The search was performed using the computer program BLAST or BLAST-2 [Altschul *et al.*, Methods in Enzymology, 266:460-480 (1996)] as a comparison of the ECD protein sequences to a 6 frame translation of the EST sequences. Those comparisons with a BLAST score of 70 (or in some cases, 90) or greater that did not encode known proteins were clustered and assembled into consensus DNA sequences with the program "phrap" (Phil Green, University of Washington, Seattle, Washington).

Using this extracellular domain homology screen, consensus DNA sequences were assembled relative to other identified EST sequences using phrap. In addition, the consensus DNA sequences obtained were often (but not always) extended using repeated cycles of BLAST and phrap to extend the consensus sequence as far as possible using the sources of EST sequences discussed above.

Based upon the consensus sequences obtained as described above, oligonucleotides were then synthesized and used to identify by PCR a cDNA library that contained the sequence of interest and for use as probes to isolate a clone of the full-length coding sequence for a PRO polypeptide. Forward and reverse PCR primers generally range from 20 to 30 nucleotides and are often designed to give a PCR product of about 100-1000 bp in length. The probe sequences are typically 40-55 bp in length. In some cases, additional oligonucleotides are synthesized when the consensus sequence is greater than about 1-1.5 kbp. In order to screen several libraries for a full-length clone, DNA from the libraries was screened by PCR amplification, as per Ausubel *et al.*, Current Protocols in Molecular Biology, *supra*, with the PCR primer pair. A positive library was then used to isolate clones encoding the gene of interest using the probe oligonucleotide and one of the primer pairs.

The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with oligo dT containing a NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes *et al.*, Science, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

## EXAMPLE 2

### Isolation of cDNA clones by Amylase Screening

#### 1. Preparation of oligo dT primed cDNA library

mRNA was isolated from a human tissue of interest using reagents and protocols from Invitrogen, San Diego, CA (Fast Track 2). This RNA was used to generate an oligo dT primed cDNA library in the vector pRK5D using reagents and protocols from Life Technologies, Gaithersburg, MD (Super Script Plasmid System). In this procedure, the double stranded cDNA was sized to greater than 1000 bp and the SalI/NotI linked cDNA was cloned into XhoI/NotI cleaved vector. pRK5D is a cloning vector that has an sp6 transcription initiation site followed by an SfiI restriction enzyme site preceding the XhoI/NotI cDNA cloning sites.

#### 2. Preparation of random primed cDNA library

A secondary cDNA library was generated in order to preferentially represent the 5' ends of the primary cDNA clones. Sp6 RNA was generated from the primary library (described above), and this RNA was used to generate a random primed cDNA library in the vector pSST-AMY.0 using reagents and protocols from Life Technologies (Super Script Plasmid System, referenced above). In this procedure the double stranded cDNA was sized to 500-1000 bp, linked with blunt to NotI adaptors, cleaved with SfiI, and cloned into SfiI/NotI cleaved vector. pSST-AMY.0 is a cloning vector that has a yeast alcohol dehydrogenase promoter preceding the cDNA cloning sites and the mouse amylase sequence (the mature sequence without the secretion signal) followed by the yeast alcohol dehydrogenase terminator, after the cloning sites. Thus, cDNAs cloned into this vector that are fused in frame with amylase sequence will lead to the secretion of amylase from appropriately transfected yeast colonies.

#### 3. Transformation and Detection

DNA from the library described in paragraph 2 above was chilled on ice to which was added electrocompetent DH10B bacteria (Life Technologies, 20 ml). The bacteria and vector mixture was then electroporated as recommended by the manufacturer. Subsequently, SOC media (Life Technologies, 1 ml) was added and the mixture was incubated at 37°C for 30 minutes. The transformants were then plated onto 20 standard 150 mm LB plates containing ampicillin and incubated for 16 hours (37°C). Positive colonies were scraped off the plates and the DNA was isolated from the bacterial pellet using standard protocols, *e.g.*, CsCl-gradient. The purified DNA was then carried on to the yeast protocols below.

The yeast methods were divided into three categories: (1) Transformation of yeast with the plasmid/cDNA

combined vector; (2) Detection and isolation of yeast clones secreting amylase; and (3) PCR amplification of the insert directly from the yeast colony and purification of the DNA for sequencing and further analysis.

The yeast strain used was HD56-5A (ATCC-90785). This strain has the following genotype: MAT alpha, ura3-52, leu2-3, leu2-112, his3-11, his3-15, MAL<sup>+</sup>, SUC<sup>+</sup>, GAL<sup>+</sup>. Preferably, yeast mutants can be employed that have deficient post-translational pathways. Such mutants may have translocation deficient alleles in *sec71*, *sec72*, *sec62*, with truncated *sec71* being most preferred. Alternatively, antagonists (including antisense nucleotides and/or ligands) which interfere with the normal operation of these genes, other proteins implicated in this post translation pathway (e.g., SEC61p, SEC72p, SEC62p, SEC63p, TDJ1p or SSA1p-4p) or the complex formation of these proteins may also be preferably employed in combination with the amylase-expressing yeast.

Transformation was performed based on the protocol outlined by Gietz *et al.*, Nucl. Acid. Res., 20:1425 (1992). Transformed cells were then inoculated from agar into YEPD complex media broth (100 ml) and grown overnight at 30°C. The YEPD broth was prepared as described in Kaiser *et al.*, Methods in Yeast Genetics, Cold Spring Harbor Press, Cold Spring Harbor, NY, p. 207 (1994). The overnight culture was then diluted to about  $2 \times 10^6$  cells/ml (approx. OD<sub>600</sub>=0.1) into fresh YEPD broth (500 ml) and regrown to  $1 \times 10^7$  cells/ml (approx. OD<sub>600</sub>=0.4-0.5).

The cells were then harvested and prepared for transformation by transfer into GS3 rotor bottles in a Sorval GS3 rotor at 5,000 rpm for 5 minutes, the supernatant discarded, and then resuspended into sterile water, and centrifuged again in 50 ml falcon tubes at 3,500 rpm in a Beckman GS-6KR centrifuge. The supernatant was discarded and the cells were subsequently washed with LiAc/TE (10 ml, 10 mM Tris-HCl, 1 mM EDTA pH 7.5, 100 mM Li<sub>2</sub>OOCCH<sub>3</sub>), and resuspended into LiAc/TE (2.5 ml).

Transformation took place by mixing the prepared cells (100 µl) with freshly denatured single stranded salmon testes DNA (Lofstrand Labs, Gaithersburg, MD) and transforming DNA (1 µg, vol. < 10 µl) in microfuge tubes. The mixture was mixed briefly by vortexing, then 40% PEG/TE (600 µl, 40% polyethylene glycol-4000, 10 mM Tris-HCl, 1 mM EDTA, 100 mM Li<sub>2</sub>OOCCH<sub>3</sub>, pH 7.5) was added. This mixture was gently mixed and incubated at 30°C while agitating for 30 minutes. The cells were then heat shocked at 42°C for 15 minutes, and the reaction vessel centrifuged in a microfuge at 12,000 rpm for 5-10 seconds, decanted and resuspended into TE (500 µl, 10 mM Tris-HCl, 1 mM EDTA pH 7.5) followed by recentrifugation. The cells were then diluted into TE (1 ml) and aliquots (200 µl) were spread onto the selective media previously prepared in 150 mm growth plates (VWR).

Alternatively, instead of multiple small reactions, the transformation was performed using a single, large scale reaction, wherein reagent amounts were scaled up accordingly.

The selective media used was a synthetic complete dextrose agar lacking uracil (SCD-Ura) prepared as described in Kaiser *et al.*, Methods in Yeast Genetics, Cold Spring Harbor Press, Cold Spring Harbor, NY, p. 208-210 (1994). Transformants were grown at 30°C for 2-3 days.

The detection of colonies secreting amylase was performed by including red starch in the selective growth media. Starch was coupled to the red dye (Reactive Red-120, Sigma) as per the procedure described by Biely *et al.*, Anal. Biochem., 172:176-179 (1988). The coupled starch was incorporated into the SCD-Ura agar plates at a final concentration of 0.15% (w/v), and was buffered with potassium phosphate to a pH of 7.0 (50-100 mM final

concentration).

The positive colonies were picked and streaked across fresh selective media (onto 150 mm plates) in order to obtain well isolated and identifiable single colonies. Well isolated single colonies positive for amylase secretion were detected by direct incorporation of red starch into buffered SCD-Ura agar. Positive colonies were determined by their ability to break down starch resulting in a clear halo around the positive colony visualized directly.

#### 4. Isolation of DNA by PCR Amplification

When a positive colony was isolated, a portion of it was picked by a toothpick and diluted into sterile water (30  $\mu$ l) in a 96 well plate. At this time, the positive colonies were either frozen and stored for subsequent analysis or immediately amplified. An aliquot of cells (5  $\mu$ l) was used as a template for the PCR reaction in a 25  $\mu$ l volume containing: 0.5  $\mu$ l Klentaq (Clontech, Palo Alto, CA); 4.0  $\mu$ l 10 mM dNTP's (Perkin Elmer-Cetus); 2.5  $\mu$ l Klentaq buffer (Clontech); 0.25  $\mu$ l forward oligo 1; 0.25  $\mu$ l reverse oligo 2; 12.5  $\mu$ l distilled water.

The sequence of the forward oligonucleotide 1 was:

5'-TGTAACGACGGCCAGTTAAATAGACCTGCAATTATTAATCT-3' (SEQ ID NO:3)

The sequence of the reverse oligonucleotide 2 was:

5'-CAGGAAACAGCTATGACCACCTGCACACCTGCAAATCCATT-3' (SEQ ID NO:4)

PCR was then performed as follows:

- |                  |          |                  |
|------------------|----------|------------------|
| a.               | Denature | 92°C, 5 minutes  |
| b. 3 cycles of:  | Denature | 92°C, 30 seconds |
|                  | Anneal   | 59°C, 30 seconds |
|                  | Extend   | 72°C, 60 seconds |
| c. 3 cycles of:  | Denature | 92°C, 30 seconds |
|                  | Anneal   | 57°C, 30 seconds |
|                  | Extend   | 72°C, 60 seconds |
| d. 25 cycles of: | Denature | 92°C, 30 seconds |
|                  | Anneal   | 55°C, 30 seconds |
|                  | Extend   | 72°C, 60 seconds |
| e.               | Hold     | 4°C              |

The underlined regions of the oligonucleotides annealed to the ADH promoter region and the amylase region, respectively, and amplified a 307 bp region from vector pSST-AMY.0 when no insert was present. Typically, the first 18 nucleotides of the 5' end of these oligonucleotides contained annealing sites for the sequencing primers. Thus, the total product of the PCR reaction from an empty vector was 343 bp. However, signal sequence-fused cDNA resulted in considerably longer nucleotide sequences.

Following the PCR, an aliquot of the reaction (5  $\mu$ l) was examined by agarose gel electrophoresis in a 1% agarose gel using a Tris-Borate-EDTA (TBE) buffering system as described by Sambrook *et al.*, *supra*. Clones resulting in a single strong PCR product larger than 400 bp were further analyzed by DNA sequencing after purification with a 96 Qiaquick PCR clean-up column (Qiagen Inc., Chatsworth, CA).

## EXAMPLE 3

Isolation of cDNA Clones Using Signal Algorithm Analysis

Various polypeptide-encoding nucleic acid sequences were identified by applying a proprietary signal sequence finding algorithm developed by Genentech, Inc., (South San Francisco, CA) upon ESTs as well as clustered and assembled EST fragments from public (*e.g.*, GenBank) and/or private (LIFESEQ®, Incyte Pharmaceuticals, Inc., Palo Alto, CA) databases. The signal sequence algorithm computes a secretion signal score based on the character of the DNA nucleotides surrounding the first and optionally the second methionine codon(s) (ATG) at the 5'-end of the sequence or sequence fragment under consideration. The nucleotides following the first ATG must code for at least 35 unambiguous amino acids without any stop codons. If the first ATG has the required amino acids, the second is not examined. If neither meets the requirement, the candidate sequence is not scored. In order to determine whether the EST sequence contains an authentic signal sequence, the DNA and corresponding amino acid sequences surrounding the ATG codon are scored using a set of seven sensors (evaluation parameters) known to be associated with secretion signals. Use of this algorithm resulted in the identification of numerous polypeptide-encoding nucleic acid sequences.

## EXAMPLE 4

Isolation of cDNA clones Encoding PRO172

A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA28765. Based on the DNA28765 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO172. PCR primers (forward and reverse) were synthesized based upon the DNA28765 sequence. Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA28765 sequence.

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification, as per Ausubel *et al.*, Current Protocols in Molecular Biology, *supra*, with the PCR primer pair. A positive library was then used to isolate clones encoding the PRO172 gene using the probe oligonucleotide and one of the PCR primers.

The oligonucleotide sequences used in the above procedure were the following:

28765.p

5'-AAATCTGTGAATTGAGTGCCATGGACCTGTTGCGGACGGCCCTTGCTT-3'

(SEQ ID NO:5)

28765.f

5'-GGATCTCGAGAACAGCTACTCC-3' (SEQ ID NO:6)

28765.r

5'-TCGTCCACGTTGTCGTACATG-3' (SEQ ID NO:7)

RNA for construction of the cDNA libraries was isolated from human fetal kidney tissue. The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with NotI site, linked with blunt to Sall

hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes *et al.*, Science, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

5 DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO172 [herein designated as DNA35916-1161] (Figures 1A-1B, SEQ ID NO:1) and the derived protein sequence for PRO172.

10 The entire nucleotide sequence of DNA35916-1161 is shown in Figures 1A-1B (SEQ ID NO:1). Clone DNA35916-1161 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 38-40 and ending at the stop codon at nucleotide positions 2207-2209 (Figures 1A-1B). The predicted polypeptide precursor is 723 amino acids long (Figure 2; SEQ ID NO:2).

Analysis of the full-length PRO172 sequence shown in Figure 2 (SEQ ID NO:2) evidenced the presence of a variety of important polypeptide domains as shown in Figure 2, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO172 sequence (Figure 2; SEQ ID NO:2) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 19; a transmembrane domain from about amino acid 544 to about amino acid 565; an N-glycosylation site from about amino acid 475 to about amino acid 479; a cAMP- and cGMP-dependent protein kinase phosphorylation site from about amino acid 658 to about amino acid 662; tyrosine kinase phosphorylation sites from about amino acid 174 to about amino acid 183 and from about amino acid 250 to about amino acid 259; N-myristoylation sites from about amino acid 35 to about amino acid 41, from about amino acid 38 to about amino acid 44, from about amino acid 96 to about amino acid 102, from about amino acid 97 to about amino acid 103, from about amino acid 260 to about amino acid 266, from about amino acid 279 to about amino acid 285, from about amino acid 280 to about amino acid 286, from about amino acid 299 to about amino acid 305, from about amino acid 308 to about amino acid 314, from about amino acid 326 to about amino acid 332, from about amino acid 338 to about amino acid 344, from about amino acid 376 to about amino acid 382, from about amino acid 385 to about amino acid 391, from about amino acid 510 to about amino acid 516, from about amino acid 674 to about amino acid 680, from about amino acid 681 to about amino acid 687, and from about amino acid 693 to about amino acid 699; aspartic acid and asparagine hydroxylation sites from about amino acid 314 to about amino acid 353, from about amino acid 418 to about amino acid 430, and from about amino acid 456 to about amino acid 468; a prokaryotic membrane lipoprotein lipid attachment site from about amino acid 550 to about amino acid 561; and EGF-like domain cysteine pattern signatures from about amino acid 241 to about amino acid 253, from about amino acid 272 to about amino acid 284, from about amino acid 312 to about amino acid 324, from about amino acid 350 to about amino acid 362, from about amino acid 389 to about amino acid 401, from about amino acid 427 to about amino acid 439, from about amino acid 465 to about amino acid 477, and from about amino acid 503 to about amino acid 515. Clone DNA35916-1161 has been deposited with ATCC on October 28, 1997 and is assigned ATCC deposit no. 209419.

35 An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 2 (SEQ ID NO:2), evidenced 89% amino acid sequence identity between the PRO172 amino acid sequence and the delta-I mouse protein.

## EXAMPLE 5

Isolation of cDNA clones encoding PRO175

5 A cDNA clone (DNA 19355-1150) encoding a native human PRO175 polypeptide was identified using a yeast screen, in a human umbilical vein endothelium cell cDNA library that preferentially represents the 5' ends of the primary cDNA clones.

A full length clone was identified that contained a single open reading frame with an apparent translational initiation site at nucleotide positions 21-23, and a stop signal at nucleotide positions 552-554 (Figures 3A-3B, SEQ ID NO:8). The predicted polypeptide precursor is 177 amino acids long, and has a calculated molecular weight of approximately 20,308 daltons.

10 Analysis of the full-length PRO175 sequence shown in Figure 4 (SEQ ID NO:9) evidenced the presence of a variety of important polypeptide domains as shown in Figure 4, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO175 sequence (Figure 4; SEQ ID NO:9) evidences a type II transmembrane protein typology and the presence of the following: a signal peptide from about amino acid 1 to about amino acid 25; a transmembrane domain from about amino acid 26 to  
15 about amino acid 51; an extracellular domain from about amino acid 52 to about amino acid 177; potential N-linked glycosylation sites from about amino acid 129 to about amino acid 133 and from about amino acid 161 to about amino acid 165; and an N-myristoylation site from about amino acid 18 to about amino acid 24. Clone DNA 19355-1150 has been deposited with ATCC on November 18, 1997 and is assigned ATCC deposit no. 209466.

20 An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 4 (SEQ ID NO:9), evidenced significant identity between the PRO175 amino acid sequence and several members of the TNF cytokine family, and particularly, to human Apo-2L (19.8%), Fas/Apo1-ligand (19.0%), TNF-alpha (20.6%) and Lymphotoxin- $\alpha$  (17.5%). Most of the amino acid sequence identity is found in the regions corresponding to the beta-strands in the crystal structure of TNF- $\alpha$  [Banner *et al.*, *Cell*, **73**:431-435 (1993); Eck *et al.*, *J. Biol. Chem.*, **264**:17595-605 (1989); Lewit-Bentley *et al.*, *J. Mol. Biol.*,  
25 **199**:389-92 (1988)]. The sequence of strand C is especially conserved in all members of the family. The sequence between the putative transmembrane domain and the first beta-strand of the DNA 19355 polypeptide is relatively short, including 5 residues, as compared to about 30 to about 80 residues in TNF- $\alpha$ , CD95L or Apo-2 ligand.

## EXAMPLE 6

Isolation of cDNA clones encoding PRO178

30 An expressed sequence tag (EST) DNA database (LIFESEQ<sup>®</sup>, Incyte Pharmaceuticals, Palo Alto, CA) was searched and an EST was identified that had homology to PRO179 [identified in EXAMPLE 9 below and designated DNA 16451-1078 (Figures 11A-11B; SEQ ID NO:25)]. To clone PRO178, a human fetal lung library prepared from mRNA purchased from Clontech, Inc., (Palo Alto, CA), catalog # 6528-1 was used, following the manufacturer's instructions.

35 The cDNA libraries used to isolate the cDNA clones encoding human PRO178 were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was

primed with oligo dT containing a NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes *et al.*, Science, 253:1278-1280 (1991)) in the unique XhoI and NotI.

5 Oligonucleotide probes based upon the above described EST sequence were then synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO178. Forward and reverse PCR primers generally range from 20-30 nucleotides and are often designed to give a PCR product of about 100-1000 bp in length. The probe sequences are typically 40-55 bp in length. In order to screen several libraries for a full-length clone, DNA from the libraries was screened  
10 by PCR amplification, as per Ausubel *et al.*, Current Protocols in Molecular Biology, *supra*, with the PCR primer pair. A positive library was then used to isolate clones encoding the gene of interest using the probe oligonucleotide and one of the primer pairs. The oligonucleotide probes used were as follows:

5'-ACGTAGTTCAGTATGGTGTGAGCAGCAACTGGA-3' (SEQ ID NO:12)

5'-AGTCCAGCCTCCACCCTCCAGTTGCT-3' (SEQ ID NO:13)

15 5'-CCCCAGTCCTCCAGGAGAACCAGCA-3' (SEQ ID NO:14)

A cDNA clone was identified and sequenced in entirety. The entire nucleotide sequence of DNA23339-1130 is shown in Figure 5 (SEQ ID NO:10). Clone DNA23339-1130 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 118-120, and a stop codon at nucleotide positions 1528-1530 (Figure 5; SEQ ID NO:10). The predicted polypeptide precursor is 470 amino acids long. The full-length  
20 PRO178 protein is shown in Figure 6 (SEQ ID NO:11).

Analysis of the full-length PRO178 sequence shown in Figure 6 (SEQ ID NO:11) evidences the presence of important polypeptide domains as shown in Figure 6, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO178 sequence (Figure 6; SEQ ID NO:11) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 20;  
25 N-glycosylation sites from about amino acid 58 to about amino acid 62 and from about amino acid 145 to about amino acid 149; a cAMP- and cGMP-dependent protein kinase phosphorylation site from about amino acid 97 to about amino acid 101; a tyrosine kinase phosphorylation site from about amino acid 441 to about amino acid 448; N-myristoylation sites from about amino acid 16 to about amino acid 22, from about amino acid 23 to about amino acid 29, from about amino acid 87 to about amino acid 93, from about amino acid 108 to about amino acid 114,  
30 from about amino acid 121 to about amino acid 127, from about amino acid 125 to about amino acid 131, from about amino acid 129 to about amino acid 135, from about amino acid 187 to about amino acid 193, from about amino acid 293 to about amino acid 299, from about amino acid 353 to about amino acid 359, from about amino acid 378 to about amino acid 384, from about amino acid 445 to about amino acid 451, and from about amino acid 453 to about amino acid 459; a cell attachment sequence from about amino acid 340 to about amino acid 343; and  
35 a fibrinogen beta and gamma chains C-terminal domain signature from about amino acid 418 to about amino acid 431.

Clone DNA23339-1130 has been deposited with ATCC on September 18, 1997 and is assigned ATCC.



deposit no. 209282. It is understood that the deposited clone has the actual correct sequence rather than the representations provided herein.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 6 (SEQ ID NO:11), shows a 23% sequence identity between the PRO178 amino acid sequence and both ligand 1 and ligand 2 of the TIE-2 receptor. The abbreviation "TIE" is an acronym which stands for "tyrosine kinase containing Ig and EGF homology domains" and was coined to designate a new family of receptor tyrosine kinases. The fibrinogen domains of ligand 1 and ligand 2 of the TIE-2 receptor are 64% identical and 40-43% identical to PRO178, respectively.

#### EXAMPLE 7

##### 10 Isolation of cDNA clones encoding PRO188

An expressed sequence tag (EST) DNA database (LIFESEQ<sup>®</sup>, Incyte Pharmaceuticals, Palo Alto, CA) was searched and an EST was identified that had homology to PRO179 [identified in EXAMPLE 9 below and designated DNA16451-1078 (Figures 11A-11B; SEQ ID NO:25)]. To clone PRO188, a human fetal lung library prepared from mRNA purchased from Clontech, Inc., (Palo Alto, CA), catalog # 6528-1 was used, following the manufacturer's instructions.

The cDNA libraries used to isolate the cDNA clones encoding human PRO188 were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with oligo dT containing a NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes *et al.*, Science, 253:1278-1280 (1991)) in the unique XhoI and NotI.

Oligonucleotide probes based upon the above described EST sequence were then synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO188. Forward and reverse PCR primers generally range from 20-30 nucleotides and are often designed to give a PCR product of about 100-1000 bp in length. The probe sequences are typically 40-55 bp in length. In order to screen several libraries for a full-length clone, DNA from the libraries was screened by PCR amplification, as per Ausubel *et al.*, Current Protocols in Molecular Biology, *supra*, with the PCR primer pair. A positive library was then used to isolate clones encoding the gene of interest using the probe oligonucleotide and one of the primer pairs.

30 The oligonucleotide sequences used were as follows:

5'-CAGGTTATCCCAGAGATTTAATGCCACCA-3' (SEQ ID NO:17)

5'-TTGGTGGGAGAAGTTGCCAGATCAGGTGGTGGCA-3' (SEQ ID NO:18)

5'-TTCACACCATAACTGCATTGGTCCA-3' (SEQ ID NO:19)

A cDNA clone was identified and sequenced in entirety. The entire nucleotide sequence of DNA28497-1130 is shown in Figures 7A-7B (SEQ ID NO:15). Clone DNA28497-1130 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 449-451, and a stop codon at nucleotide positions

1922-1924 (Figures 7A-7B; SEQ ID NO:15). The predicted polypeptide precursor is 491 amino acids long. The full-length PRO188 protein is shown in Figure 8 (SEQ ID NO:16).

Analysis of the full-length PRO188 sequence shown in Figure 8 (SEQ ID NO:16) evidences the presence of important polypeptide domains as shown in Figure 8, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO188 sequence (Figure 8; SEQ ID NO:16) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 23; N-glycosylation sites from about amino acid 160 to about amino acid 164 and from about amino acid 188 to about amino acid 192; a cAMP- and cGMP-dependent protein kinase phosphorylation site from about amino acid 120 to about amino acid 124; tyrosine kinase phosphorylation sites from about amino acid 173 to about amino acid 180 and from about amino acid 387 to about amino acid 396; N-myristoylation sites from about amino acid 70 to about amino acid 76, from about amino acid 110 to about amino acid 116, from about amino acid 232 to about amino acid 238, from about amino acid 343 to about amino acid 349, from about amino acid 400 to about amino acid 406, from about amino acid 467 to about amino acid 473, and from about amino acid 475 to about amino acid 481; and a fibrinogen beta and gamma chains C-terminal domain signature from about amino acid 440 to about amino acid 453.

Clone DNA28497-1130 has been deposited with ATCC on September 18, 1997 and is assigned ATCC deposit no. 209279. It is understood that the deposited clone has the actual correct sequence rather than the representations provided herein.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 8 (SEQ ID NO:16), shows a 24% sequence identity between the PRO188 amino acid sequence and both ligand 1 and ligand 2 of the TIE-2 receptor. The abbreviation "TIE" is an acronym which stands for "tyrosine kinase containing Ig and EGF homology domains" and was coined to designate a new family of receptor tyrosine kinases. The fibrinogen domains of ligand 1 and ligand 2 of the TIE-2 receptor are 64% identical and 40-43% identical to PRO188, respectively.

#### EXAMPLE 8

##### Isolation of cDNA clones encoding PRO356

An expressed sequence tag (EST) DNA database (LIFESEQ<sup>®</sup>, Incyte Pharmaceuticals, Palo Alto, CA) was searched and an EST (#2939340) was identified that had homology to PRO179 [identified in EXAMPLE 9 below and designated DNA16451-1078 (Figures 11A-11B; SEQ ID NO:25)]. To clone PRO356, a human fetal lung library prepared from mRNA purchased from Clontech, Inc., (Palo Alto, CA), catalog #6528-1 was used, following the manufacturer's instructions.

The cDNA libraries used to isolate the cDNA clones encoding human PRO356 were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with oligo dT containing a NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes *et al.*, Science, 253:1278-1280 (1991)) in the unique XhoI and NotI.

Oligonucleotide probes based upon the above described EST sequence were then synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO356. Forward and reverse PCR primers generally range from 20-30 nucleotides and are often designed to give a PCR product of about 100-1000 bp in length. The probe sequences are typically 40-55 bp in length. In order to screen several libraries for a full-length clone, DNA from the libraries was screened by PCR amplification, as per Ausubel *et al.*, Current Protocols in Molecular Biology, *supra*, with the PCR primer pair. A positive library was then used to isolate clones encoding the gene of interest using the probe oligonucleotide and one of the primer pairs.

The oligonucleotide sequences used were as follows:

- 10      5'-TTCAGCACCAAGGACAAGGACAATGACAACT-3'      (SEQ ID NO:22)  
          5'-TGTGCACACTTGTCCAAGCAGTTGTCATTGTC-3'      (SEQ ID NO:23)  
          5'-GTAGTACACTCCATTGAGGTTGG-3'      (SEQ ID NO:24)

A cDNA clone was identified and sequenced in entirety. The entire nucleotide sequence of DNA47470-1130-P1 is shown in Figures 9A-9B (SEQ ID NO:20). Clone DNA47470-1130-P1 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 215-217, and a stop codon at nucleotide positions 1253-1255 (Figures 9A-9B; SEQ ID NO:20). The predicted polypeptide precursor is 346 amino acids long. The full-length PRO356 protein is shown in Figure 10 (SEQ ID NO:21).

Analysis of the full-length PRO356 sequence shown in Figure 10 (SEQ ID NO:21) evidences the presence of important polypeptide domains as shown in Figure 10, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO356 sequence (Figure 10; SEQ ID NO:21) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 26; N-glycosylation sites from about amino acid 58 to about amino acid 62, from about amino acid 253 to about amino acid 257 and from about amino acid 267 to about amino acid 271; a glycosaminoglycan attachment site from about amino acid 167 to about amino acid 171; a cAMP- and cGMP-dependent protein kinase phosphorylation site from about amino acid 176 to about amino acid 180; N-myristoylation sites from about amino acid 168 to about amino acid 174, from about amino acid 196 to about amino acid 202, from about amino acid 241 to about amino acid 247, from about amino acid 252 to about amino acid 258, from about amino acid 256 to about amino acid 262, and from about amino acid 327 to about amino acid 333; and a cell attachment sequence from about amino acid 199 to about amino acid 202.

30      Clone DNA47470-1130-P1 has been deposited with ATCC on October 28, 1997 and is assigned ATCC deposit no. 209422. It is understood that the deposited clone has the actual correct sequence rather than the representations provided herein.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 10 (SEQ ID NO:21), shows amino acid sequence identity between the PRO356 amino acid sequence and both TIE-2L1 (32%) and TIE-2L2 (34%). The abbreviation "TIE" is an acronym which stands for "tyrosine kinase containing Ig and EGF homology domains" and was coined to designate a new family of receptor tyrosine kinases.

## EXAMPLE 9

Isolation of cDNA clones encoding PRO179

A cDNA clone (DNA 16451-1078) encoding a native human PRO179 polypeptide was identified using a yeast screen, in a human fetal liver library that preferentially represents the 5'ends of the primary cDNA clones.

5 Clone DNA 16451-1078 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 37-39, and a stop codon at nucleotide positions 1417-1419 (Figures 11A-11B; SEQ ID NO:25). The predicted polypeptide precursor is 460 amino acids long. The full-length PRO179 protein is shown in Figure 12 (SEQ ID NO:26).

10 Analysis of the full-length PRO179 sequence shown in Figure 12 (SEQ ID NO:26) evidences the presence of important polypeptide domains as shown in Figure 12, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO179 sequence (Figure 12; SEQ ID NO:26) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 16; N-glycosylation sites from about amino acid 23 to about amino acid 27, from about amino acid 115 to about amino acid 119, from about amino acid 296 to about amino acid 300, and from about amino acid 357 to about amino acid 361; cAMP- and cGMP-dependent protein kinase phosphorylation sites from about amino acid 100 to about amino acid 104 and from about amino acid 204 to about amino acid 208; a tyrosine kinase phosphphorylation site from about amino acid 342 to about amino acid 348; N-myristoylation sites from about amino acid 279 to about amino acid 285, from about amino acid 352 to about amino acid 358 and from about amino acid 367 to about amino acid 373; and leucine zipper patterns from about amino acid 120 to about amino acid 142 and from about amino acid 127 to about amino acid 149.

20 Clone DNA 16451-1078 has been deposited with ATCC on September 18, 1997 and is assigned ATCC deposit no. 209281. It is understood that the deposited clone has the actual correct sequence rather than the representations provided herein.

25 An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 12 (SEQ ID NO:26), evidenced the presence of a fibrinogen-like domain exhibiting a high degree of sequence homology with the two known human ligands of the TIE-2 receptor (h-TIE-2L1 and h-TIE-2L2). The abbreviation "TIE" is an acronym which stands for "tyrosine kinase containing Ig and EGF homology domains" and was coined to designate a new family of receptor tyrosine kinases. Accordingly, PRO179 has been identified as a novel member of the TIE ligand family.

30

## EXAMPLE 10

Isolation of cDNA clones encoding PRO197

PRO197 was identified by screening the GenBank database using the computer program BLAST (Altschul *et al.*, Methods in Enzymology, 266:460-480 (1996)). The PRO197 sequence was shown to have homology with known EST sequences T08223, AA122061, and M62290. None of the known EST sequences have been identified as full-length sequences, or described as ligands associated with TIE receptors. Following identification, PRO197 was cloned from a human fetal lung library prepared from mRNA purchased from Clontech, Inc., (Palo Alto, CA).

catalog # 6528-1, following the manufacturer's instructions. The library was screened by hybridization with synthetic oligonucleotide probes.

The oligonucleotide sequences used were as follows:

- 5' -ATGAGGTGGCCAAGCCTGCCCCGAAGAAAGAGGC-3' (SEQ ID NO:32)  
 5' -CAACTGGCTGGGCCATCTCGGGCAGCCTCTTCTTCGGG-3' (SEQ ID NO:33)  
 5' -CCCAGCCAGAACTCGCCGTGGGGA-3' (SEQ ID NO:34)

A cDNA clone was identified and sequenced in entirety. The entire nucleotide sequence of DNA22780-1078 is shown in Figure 13 (SEQ ID NO:30). Clone DNA22780-1078 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 23-25, and a stop codon at nucleotide positions 1382-1384 (Figure 13; SEQ ID NO:30). The predicted polypeptide precursor is 453 amino acids long. The full-length PRO197 protein is shown in Figure 14 (SEQ ID NO:31).

Analysis of the full-length PRO197 sequence shown in Figure 14 (SEQ ID NO:31) evidences the presence of important polypeptide domains as shown in Figure 14, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO197 sequence (Figure 14; SEQ ID NO:31) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 13; a transmembrane domain from about amino acid 53 to about amino acid 70; an N-glycosylation site from about amino acid 224 to about amino acid 228; cAMP- and cGMP-dependent protein kinase phosphorylation sites from about amino acid 46 to about amino acid 50 and from about amino acid 118 to about amino acid 122; N-myristoylation sites from about amino acid 50 to about amino acid 56, from about amino acid 129 to about amino acid 135, from about amino acid 341 to about amino acid 347, and from about amino acid 357 to about amino acid 363; and a fibrinogen beta and gamma chains C-terminal domain signature from about amino acid 396 to about amino acid 409.

Clone DNA22780-1078 has been deposited with ATCC on September 18, 1997 and is assigned ATCC deposit no. 209284. It is understood that the deposited clone has the actual correct sequence rather than the representations provided herein.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 14 (SEQ ID NO:31), evidenced homology between the PRO197 amino acid sequence and ligands associated with TIE receptors. The abbreviation "TIE" is an acronym which stands for "tyrosine kinase containing Ig and EGF homology domains" and was coined to designate a new family of receptor tyrosine kinases.

#### EXAMPLE 11

##### Isolation of cDNA clones encoding PRO198

PRO198 was identified by screening the GenBank database using the computer program BLAST (Altschul *et al.*, Methods in Enzymology, 266:460-480 (1996)). The PRO198 sequence was shown to have homology with known EST sequences T57280 and T50719. None of the known EST sequences have been identified as full-length sequences, or described as ligands associated with TIE receptors. Following identification, PRO198 was cloned

from a human fetal lung library prepared from mRNA purchased from Clontech, Inc., (Palo Alto, CA), catalog # 6528-1, following the manufacturer's instructions. The library was screened by hybridization with synthetic oligonucleotide probes.

The oligonucleotide sequences used were as follows:

- 5        5'-TGGTTGGCAAAGGCAAGGTGGCTGACGATCCGG-3'        (SEQ ID NO:37)  
           5'-GTGGCCCTTATCTCTCCTGTACAGCTTCCGGATCGTCAGCCAC-3'        (SEQ ID NO:38)  
           5'-TCCATTCCCACCTATGACGCTGACCCA-3'        (SEQ ID NO:39)

- 10        A cDNA clone was identified and sequenced in entirety. The entire nucleotide sequence of DNA33457-1078 is shown in Figure 15 (SEQ ID NO:35). Clone DNA33457-1078 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 33-35, and a stop codon at nucleotide positions 897-899 (Figure 15; SEQ ID NO:35). The predicted polypeptide precursor is 288 amino acids long. The full-length PRO198 protein is shown in Figure 16 (SEQ ID NO:36).

- 15        Analysis of the full-length PRO198 sequence shown in Figure 16 (SEQ ID NO:36) evidences the presence of important polypeptide domains as shown in Figure 16, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO198 sequence (Figure 16; SEQ ID NO:36) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 16; an N-glycosylation site from about amino acid 178 to about amino acid 182; a glycosaminoglycan attachment site from about amino acid 272 to about amino acid 276; a tyrosine kinase phosphorylation site from about amino acid 188 to about amino acid 197; N-myristoylation sites from about amino acid 16 to about amino acid 20  
 20        22, from about amino acid 89 to about amino acid 95, from about amino acid 144 to about amino acid 150, and from about amino acid 267 to about amino acid 273; and a fibrinogen beta and gamma chains C-terminal domain signature from about amino acid 242 to about amino acid 255.

- 25        Clone DNA33457-1078 has been deposited with ATCC on September 18, 1997 and is assigned ATCC deposit no. 209283. It is understood that the deposited clone has the actual correct sequence rather than the representations provided herein.

- 30        An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 16 (SEQ ID NO:36), evidenced homology between the PRO198 amino acid sequence and ligands associated with TIE receptors. The abbreviation "TIE" is an acronym which stands for "tyrosine kinase containing Ig and EGF homology domains" and was coined to designate a new family of receptor tyrosine kinases.

## EXAMPLE 12

### Isolation of cDNA clones encoding PRO182

- 35        An expressed sequence tag (EST) DNA database (LIFESEQ<sup>®</sup>, Incyte Pharmaceuticals, Palo Alto, CA) was searched and two Incyte EST sequences (Incyte EST no. INC2328985 and Incyte EST no. INC778319) were identified that had approximately 40% homology to a region of the human relaxin nucleic acid sequence and represent sequences within a gene of an insulin-like polypeptide. To clone PRO182, a human uterus library

prepared from mRNA purchased from Clontech, Inc., (Palo Alto, CA), catalog # 65237-1 was used, following the manufacturer's instructions. The cDNA libraries used to isolate the cDNA clones encoding human PRO182 were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with oligo dT containing a NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes *et al.*, Science, 253:1278-1280 (1991)) in the unique XhoI and NotI.

Oligonucleotide probes based upon the above described EST sequences were then synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO182. Forward and reverse PCR primers generally range from 20-30 nucleotides and are often designed to give a PCR product of about 100-1000 bp in length. The probe sequences are typically 40-55 bp in length. In order to screen several libraries for a full-length clone, DNA from the libraries was screened by PCR amplification, as per Ausubel *et al.*, Current Protocols in Molecular Biology, *supra*, with the PCR primer pair. A positive library was then used to isolate clones encoding the gene of interest. The library was screened by hybridization with synthetic oligonucleotide probes based on the ESTs found in the database.

The oligonucleotide sequences used were as follows:

5'-CACATTCAGTCCTCAGCAAAATGAA-3' (SEQ ID NO:42)

5'-GAGAATAAAACAGAGTGAAAATGGAGCCCTTCA TTTTGC-3' (SEQ ID NO:43)

5'-CTCAGCTTGCTGAGCTTGAGGGA-3' (SEQ ID NO:44)

A cDNA clone was identified and sequenced in entirety. The entire nucleotide sequence of DNA27865-1091 is shown in Figure 17 (SEQ ID NO:40). Clone DNA27865-1091 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 39-41, and a stop codon at nucleotide positions 444-446 (Figure 17; SEQ ID NO:40). The predicted polypeptide precursor is 135 amino acids long. The full-length PRO182 protein is shown in Figure 18 (SEQ ID NO:41).

Analysis of the full-length PRO182 sequence shown in Figure 18 (SEQ ID NO:41) evidences the presence of important polypeptide domains as shown in Figure 18, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO182 sequence (Figure 18; SEQ ID NO:41) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 18; a cAMP- and cGMP-dependent protein kinase phosphorylation site from about amino acid 107 to about amino acid 111; N-myristoylation sites from about amino acid 3 to about amino acid 9, from about amino acid 52 to about amino acid 58, from about amino acid 96 to about amino acid 102, and from about amino acid 125 to about amino acid 131; and an insulin family signature from about amino acid 121 to about amino acid 136.

Clone DNA27865-1091 has been deposited with ATCC on September 23, 1997 and is assigned ATCC deposit no. 209296. It is understood that the deposited clone has the actual correct sequence rather than the representations provided herein.

Based on an ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 18 (SEQ ID NO:41), PRO182 was homologous to but clearly different (no exact matches) from any known member of the

insulin family of proteins, and therefore PRO182 constitutes a novel member of the insulin family of proteins.

A phylogenetic analysis shows that PRO182 is closely related to other well characterized human insulin and insulin-like polypeptides. The insulin-like molecules share several characteristics. They are each secreted proteins, and each possesses a similar arrangement of six conserved cysteine residues. Numerous additional amino acids are also generally conserved between members of the family indicating an evolutionary relationship.

### EXAMPLE 13

#### Isolation of cDNA clones encoding PRO195

A clone designated herein as DNA13199\_ABI2 was isolated as described in Example 2 above. The DNA13199\_ABI2 sequence was then compared to a variety of expressed sequence tag (EST) databases which included public EST databases (e.g., GenBank) to identify existing homologies. The homology search was performed using the computer program BLAST or BLAST2 (Altschul *et al.*, Methods in Enzymology, 266:460-480 (1996)). Those comparisons resulting in a BLAST score of 70 (or in some cases, 90) or greater that did not encode known proteins were clustered and assembled into consensus DNA sequences with the program "phrap" (Phil Green, University of Washington, Seattle, Washington). The consensus sequence is herein designated DNA22778.

Oligonucleotide probes were then generated from the DNA22778 molecule and used to screen a human placenta tissue library (LIB89) prepared as described in paragraph 1 of Example 2 above. The cloning vector was pRK5B (pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes *et al.*, Science, 253:1278-1280 (1991)) in the unique XhoI and NotI sites, and the cDNA size cut was less than 2800 bp. The oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO195. Forward and reverse PCR primers generally range from 20 to 30 nucleotides and are often designed to give a PCR product of about 100-1000 bp in length. The probe sequences are typically 40-55 bp in length. In order to screen several libraries for a full-length clone, DNA from the libraries was screened by PCR amplification, as per Ausubel *et al.*, Current Protocols in Molecular Biology, *supra*, with the PCR primer pair. A positive library was then used to isolate clones encoding the gene of interest using the probe oligonucleotide and one of the primer pairs.

A pair of PCR primers (forward and reverse) were synthesized:

forward PCR primer (22778.f)

5'-ACAAGCTGAGCTGCTGTGACAG-3'

(SEQ ID NO:47)

reverse PCR primer (22778.r)

5'-TGATTCTGGCAACCAAGATGGC-3'

(SEQ ID NO:48).

Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA22778 sequence which had the following nucleotide sequence:

hybridization probe (22778.p)

5'-ATGGCCTTGGCCGGAGGTTCTGGGGACCGCTTCGGCTGAAG-3' (SEQ ID NO:49)

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO195 and the derived protein sequence for PRO195. The entire nucleotide sequence of DNA26847-1395 is shown in



Figure 19 (SEQ ID NO:45). Clone DNA26847-1395 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 70-72 and ending at the stop codon at nucleotide positions 1039-1041 (Figure 19). The predicted polypeptide precursor is 323 amino acids long (Figure 20, SEQ ID NO:46). The full-length PRO195 protein shown in Figure 20 has an estimated molecular weight of about 36,223 daltons and a pI of about 5.06.

Analysis of the full-length PRO195 sequence shown in Figure 20 (SEQ ID NO:46) evidences the presence of a variety of important polypeptide domains as shown in Figure 20, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO195 sequence (Figure 20; SEQ ID NO:46), evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 31; a transmembrane domain from about amino acid 241 to about amino acid 260; N-myristoylation sites from about amino acid 28 to about amino acid 34, from about amino acid 29 to about amino acid 35, from about amino acid 31 to about amino acid 37, and from about amino acid 86 to about amino acid 92; and a potential N-glycosylation site from about amino acid 90 to about amino acid 94. The corresponding nucleotides can be routinely determined given the sequences provided herein.

Clone DNA26847-1395 has been deposited with the ATCC on April 14, 1998 and is assigned ATCC deposit no. 209772. Regarding the sequence, it is understood that the deposited clone contains the correct sequence, and the sequences provided herein are based on known sequencing techniques.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 20 (SEQ ID NO:46), evidenced some degree of sequence identity between the PRO195 amino acid sequence and the following Dayhoff sequences: P\_P91380, AF035118\_1, HUMTROPCS\_1, NUOD\_SALTY and E70002.

#### EXAMPLE 14

##### Isolation of cDNA clones encoding PRO200

An expressed sequence tag (EST) DNA database (LIFESEQ®, Incyte Pharmaceuticals, Palo Alto, CA) was searched and Incyte EST clone "INC1302516" was identified that had homology to VEGF. Probes based on the Incyte EST clone "INC1302516" were used to screen a cDNA library derived from the human glioma cell line G61. In particular, Incyte clone "INC1302516" was used to generate the following four probes:

5'-ACTTCTCAGTGTCCATAAGGG-3' (SEQ ID NO:52)  
 5'-GAACTAAAGAGAACCGATACCATTTTCTGGCCAGGTTGTC-3' (SEQ ID NO:53)  
 5'-CACCACAGCGTTTAACCAGG-3' (SEQ ID NO:54)  
 5'-ACAACAGGCACAGTTCCAC-3' (SEQ ID NO:55)

The cDNA libraries used to isolate the cDNA clones encoding human PRO200 were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with oligo dT containing a NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes *et al.*, Science,

253:1278-1280 (1991)) in the unique XhoI and NotI.

A cDNA clone was identified and sequenced in entirety. The entire nucleotide sequence of DNA29101-1122 is shown in Figures 21A-21B (SEQ ID NO:50). Clone DNA29101-1122 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 285-287, and a stop codon at nucleotide positions 1320-1322 (Figures 21A-21B; SEQ ID NO:50). The predicted polypeptide precursor is 345 amino acids long. The full-length PRO200 protein is shown in Figure 22 (SEQ ID NO:51).

Analysis of the full-length PRO200 sequence shown in Figure 22 (SEQ ID NO:51) evidences the presence of important polypeptide domains as shown in Figure 22, wherein the location given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO200 sequence (Figure 22; SEQ ID NO:51) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 14; N-glycosylation sites from about amino acid 25 to about amino acid 29, from about amino acid 55 to about amino acid 59, and from about amino acid 254 to about amino acid 258; N-myristoylation sites from about amino acid 15 to about amino acid 21, from about amino acid 117 to about amino acid 123, from about amino acid 127 to about amino acid 133, from about amino acid 281 to about amino acid 287, from about amino acid 282 to about amino acid 288, and from about amino acid 319 to about amino acid 325; and an amidation site from about amino acid 229 to about amino acid 233.

Clone DNA29101-1122 has been deposited with ATCC on March 5, 1998 and is assigned ATCC deposit no. 209653. It is understood that the deposited clone has the actual correct sequence rather than the representations provided herein.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 22 (SEQ ID NO:51), evidenced some degree of sequence identity between the PRO200 amino acid sequence and the vascular endothelial cell growth factor (VEGF) and the bone morphogenetic protein (VEGF-E polypeptide).

#### EXAMPLE 15

##### Isolation of cDNA clones encoding PRO211

A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA28730. Based on the DNA28730 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO211. PCR primers (forward and reverse) were synthesized based upon the DNA28730 sequence. Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA28730 sequence.

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO211 gene using the probe oligonucleotide and one of the PCR primers.

The oligonucleotide sequences used in the above procedure were the following:

hybridization probe (28730.p)

5'-AGGGAGCACGGACAGTGTGCAGATGTGGACGAGTGCTCACTAGCA-3'

(SEQ ID NO:58)

forward PCR primer (28730.f)

5'-AGAGTGTATCTCTGGCTACGC-3'

(SEQ ID NO:59)

reverse PCR primer (28730.r)

5'-TAAGTCCGGCACATTACAGGTC-3'

(SEQ ID NO:60)

RNA for construction of the cDNA libraries was isolated from human fetal lung tissue. The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes *et al.*, *Science*, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO211 [herein designated as DNA32292-1131] (Figure 23, SEQ ID NO:56) and the derived protein sequence for PRO211.

The entire nucleotide sequence of DNA32292-1131 is shown in Figure 23 (SEQ ID NO:56). Clone DNA32292-1131 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 65-67 and ending at the stop codon at nucleotide positions 1124-1126 (Figure 23). The predicted polypeptide precursor is 353 amino acids long (Figure 24; SEQ ID NO:57). The full-length PRO211 protein shown in Figure 24 has an estimated molecular weight of about 38,190 daltons.

Analysis of the full-length PRO211 sequence shown in Figure 24 (SEQ ID NO:57) evidences the presence of important polypeptide domains as shown in Figure 24, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO211 sequence (Figure 24; SEQ ID NO:57) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 24; N-glycosylation sites from about amino acid 190 to about amino acid 194 and from about amino acid 251 to about amino acid 255; glycosaminoglycan attachment sites from about amino acid 149 to about amino acid 153 and from about amino acid 155 to about amino acid 159; a cAMP- and cGMP-dependent protein kinase phosphorylation site from about amino acid 26 to about amino acid 30; a tyrosine kinase phosphorylation site from about amino acid 303 to about amino acid 310; N-myristoylation sites from about amino acid 44 to about amino acid 50, from about amino acid 54 to about amino acid 60, from about amino acid 55 to about amino acid 61, from about amino acid 81 to about amino acid 87, from about amino acid 150 to about amino acid 156, from about amino acid 158 to about amino acid 164, from about amino acid 164 to about amino acid 170, from about amino acid 252 to about amino acid 258, and from about amino acid 313 to about amino acid 319; an aspartic acid and asparagine hydroxylation site from about amino acid 308 to about amino acid 320; an EGF-like domain cysteine pattern signature from about amino acid 166 to about amino acid 178; and a leucine zipper pattern from about amino acid 94 to about amino acid 116.

Clone DNA32292-1131 has been deposited with ATCC on September 16, 1997 and is assigned ATCC deposit

no. 209258.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 24 (SEQ ID NO:57), evidenced amino acid sequence identity between the PRO211 amino acid sequence and the EGF-like polypeptides.

5

#### EXAMPLE 16

##### Isolation of cDNA clones encoding PRO217

A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA28760. Based on the DNA28760 consensus  
10 sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO217. PCR primers (forward and reverse) were synthesized based upon the DNA28760 sequence. Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA28760 sequence.

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened  
15 by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO217 gene using the probe oligonucleotide and one of the PCR primers.

The oligonucleotide sequences used in the above procedure were the following:

##### hybridization probe (28760.p)

5'-CCCACGATGTATGAATGGTGGACTTTGTGTGACTCCTGGTTTCTGCATC-3'

20 (SEQ ID NO:63)

##### forward PCR primer (28760.f)

5'-AAAGACGCATCTGCGAGTGTCC-3' (SEQ ID NO:64)

##### reverse PCR primer (28760.r)

5'-TGCTGATTTCACACTGCTCTCCC-3' (SEQ ID NO:65)

25 RNA for construction of the cDNA libraries was isolated from human fetal lung tissue. The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not  
30 contain the SfiI site; see, Holmes *et al.*, *Science*, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO217 [herein designated as DNA33094-1131] (Figures 25A-25B, SEQ ID NO:61) and the derived protein sequence for PRO217.

35 The entire nucleotide sequence of DNA33094-1131 is shown in Figures 25A-25B (SEQ ID NO:61). Clone DNA33094-1131 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 146-148 and ending at the stop codon at nucleotide positions 1283-1285 (Figures 25A-25B). The predicted polypeptide precursor is 379 amino acids long (Figure 26; SEQ ID NO:62). The full-length PRO217

protein shown in Figure 26 has an estimated molecular weight of about 41,520 daltons.

Analysis of the full-length PRO217 sequence shown in Figure 26 (SEQ ID NO:62) evidences the presence of important polypeptide domains as shown in Figure 26, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO217 sequence (Figure 26; SEQ ID NO:62) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 28; N-glycosylation sites from about amino acid 88 to about amino acid 92 and from about amino acid 245 to about amino acid 249; a tyrosine kinase phosphorylation site from about amino acid 370 to about amino acid 378; N-myristoylation sites from about amino acid 184 to about amino acid 190, from about amino acid 185 to about amino acid 191, from about amino acid 189 to about amino acid 195, and from about amino acid 315 to about amino acid 321; an ATP/GTP-binding site motif A (P-loop) from about amino acid 285 to about amino acid 293; and EGF-like domain cysteine pattern signatures from about amino acid 198 to about amino acid 210, from about amino acid 230 to about amino acid 242, from about amino acid 262 to about amino acid 274, from about amino acid 294 to about amino acid 306, and from about amino acid 326 to about amino acid 338.

Clone DNA33094-1131 has been deposited with ATCC on September 16, 1997 and is assigned ATCC deposit no. 209256.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 26 (SEQ ID NO:62), evidenced amino acid sequence identity between the PRO217 amino acid sequence and the EGF-like polypeptides.

#### EXAMPLE 17

##### Isolation of cDNA clones encoding PRO219

A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA28729. Based on the DNA28729 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO219. PCR primers (forward and reverse) were synthesized based upon the DNA28729 sequence. Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA28729 sequence.

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO219 gene using the probe oligonucleotide and one of the PCR primers.

The oligonucleotide sequences used in the above procedure were the following:

##### forward PCR primer (28729.f)

5'-GTGACCCTGGTTGTGAATACTCC-3' (SEQ ID NO:68)

##### reverse PCR primer (28730.r)

5'-ACAGCCATGGTCTATAGCTTGG-3' (SEQ ID NO:69)

hybridization probe (28730.p)

5'-GCCTGTCAGTGTCTGAGGGACACGTGCTCCGCAGCGATGGGAAG-3'

(SEQ ID NO:70)

RNA for construction of the cDNA libraries was isolated from human fetal kidney tissue. The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with NotI site, linked with blunt to Sall hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes *et al.*, Science, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO219 [herein designated as DNA32290-1164] (Figures 27A-27B, SEQ ID NO:66) and the derived protein sequence for PRO219.

The entire nucleotide sequence of DNA32290-1164 is shown in Figures 27A-27B (SEQ ID NO:66). Clone DNA32290-1164 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 204-206 and ending at the stop codon at nucleotide positions 2949-2951 (Figures 27A-27B). The predicted polypeptide precursor is 915 amino acids long (Figure 28; SEQ ID NO:67).

Analysis of the full-length PRO219 sequence shown in Figure 28 (SEQ ID NO:67) evidences the presence of important polypeptide domains as shown in Figure 28, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO219 sequence (Figure 28; SEQ ID NO:67) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 23; an N-glycosylation site from about amino acid 221 to about amino acid 225; cAMP- and cGMP-dependent protein kinase phosphorylation sites from about amino acid 115 to about amino acid 119, from about amino acid 606 to about amino acid 610, and from about amino acid 892 to about amino acid 896; N-myristoylation sites from about amino acid 133 to about amino acid 139, from about amino acid 258 to about amino acid 291, from about amino acid 299 to about amino acid 305, from about amino acid 340 to about amino acid 346, from about amino acid 453 to about amino acid 459, from about amino acid 494 to about amino acid 500, from about amino acid 639 to about amino acid 645, from about amino acid 690 to about amino acid 696, from about amino acid 752 to about amino acid 758, and from about amino acid 792 to about amino acid 798; amidation sites from about amino acid 314 to about amino acid 318, from about amino acid 560 to about amino acid 564 and from about amino acid 601 to about amino acid 605; and aspartic acid and asparagine hydroxylation sites from about amino acid 253 to about amino acid 265, from about amino acid 294 to about amino acid 306, from about amino acid 335 to about amino acid 347, from about amino acid 376 to about amino acid 388, from about amino acid 417 to about amino acid 429, from about amino acid 458 to about amino acid 470, from about amino acid 540 to about amino acid 552, and from about amino acid 581 to about amino acid 593.

Clone DNA32290-1164 has been deposited with ATCC on October 16, 1997 and is assigned ATCC deposit no. 209384.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment

analysis of the full-length sequence shown in Figure 28 (SEQ ID NO:67), evidenced amino acid sequence identity between the PRO219 amino acid sequence and the mouse and human matrilin-2 precursor polypeptides.

#### EXAMPLE 18

##### Isolation of cDNA clones encoding PRO221

5 A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA28756. Based on the DNA28756 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO221. PCR primers (forward and reverse) were synthesized based upon the DNA28756 sequence. Additionally, a synthetic  
10 oligonucleotide hybridization probe was constructed from the consensus DNA28756 sequence.

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO221 gene using the probe oligonucleotide and one of the PCR primers.

The oligonucleotide sequences used in the above procedure were the following:

15 forward PCR primer (28756.f)

5'-CCATGTGTCTCCTCCTACAAAG-3' (SEQ ID NO:73)

reverse PCR primer (28756.r)

5'-GGGAATAGATGTGATCTGATTGG-3' (SEQ ID NO:74)

hybridization probe (28756.p)

20 5'-CACCTGTAGCAATGCAAATCTCAAGGAAATACCTAGAGATCTTCCTCCTG-3' (SEQ ID NO:75)

RNA for construction of the cDNA libraries was isolated from human fetal lung tissue. The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with NotI site, linked with blunt to Sall hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined  
25 orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes *et al.*, *Science*, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO221 [herein designated as DNA33089-1132] (Figure 29, SEQ ID NO:71) and the derived protein sequence for PRO221.

The entire nucleotide sequence of DNA33089-1132 is shown in Figure 29 (SEQ ID NO:71). Clone  
30 DNA33089-1132 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 179-181 and ending at the stop codon at nucleotide positions 956-958 (Figure 29). The predicted polypeptide precursor is 259 amino acids long (Figure 30; SEQ ID NO:72). Clone DNA33089-1132 has been deposited with ATCC on September 16, 1997 and is assigned ATCC deposit no. 209262.

Analysis of the full-length PRO221 sequence shown in Figure 30 (SEQ ID NO:72) evidences the presence  
35 of important polypeptide domains as shown in Figure 30, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO221 sequence (Figure

30; SEQ ID NO:72), evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 33; a transmembrane domain from about amino acid 206 to about amino acid 225; N-glycosylation sites from about amino acid 47 to about amino acid 51 and from about amino acid 94 to about amino acid 98; a cAMP- and cGMP-dependent protein kinase phosphorylation site from about amino acid 199 to about amino acid 203; and  
 5 N-myristoylation sites from about amino acid 37 to about amino acid 43, from about amino acid 45 to about amino acid 51 and from about amino acid 110 to about amino acid 116.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 30 (SEQ ID NO:72), evidenced amino acid sequence identity between the PRO221 amino acid sequence and a member of the leucine rich repeat protein superfamily, including  
 10 the SLIT protein.

### EXAMPLE 19

#### Isolation of cDNA clones encoding PRO224

A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA30845. Based on the DNA30845 consensus  
 15 sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO224. PCR primers (forward and reverse) were synthesized based upon the DNA30845 sequence. Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA30845 sequence.

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened  
 20 by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO224 gene using the probe oligonucleotide and one of the PCR primers.

The oligonucleotide sequences used in the above procedure were the following:

#### forward PCR primer

5'-AAGTTCCAGTGCCGCACCACTGGC-3' (SEQ ID NO:78)

#### 25 reverse PCR primer

5'-TTGGTTCCACAGCCGAGCTCGTCG-3' (SEQ ID NO:79)

#### hybridization probe

5'-GAGGAGGAGTGCAGGATTGAGCCATGTACCCAGAAAGGGCAATGCCCACC-3'

(SEQ ID NO:80)

30 RNA for construction of the cDNA libraries was isolated from human fetal liver tissue. The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not  
 35 contain the SfiI site; see, Holmes *et al.*, Science, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO224



[herein designated as DNA33221-1133] (Figure 31, SEQ ID NO:76) and the derived protein sequence for PRO224.

The entire nucleotide sequence of DNA33221-1133 is shown in Figure 31 (SEQ ID NO:76). Clone DNA33221-1133 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 33-35 and ending at the stop codon at nucleotide positions 879-881 (Figure 31). The predicted polypeptide precursor is 282 amino acids long (Figure 32; SEQ ID NO:77).

Analysis of the full-length PRO224 sequence shown in Figure 32 (SEQ ID NO:77) evidences the presence of important polypeptide domains as shown in Figure 32, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO224 sequence (Figure 32; SEQ ID NO:77) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 30; a transmembrane domain from about amino acid 231 to about amino acid 248; N-glycosylation sites from about amino acid 126 to about amino acid 130, from about amino acid 195 to about amino acid 199 and from about amino acid 213 to about amino acid 217; N-myristoylation sites from about amino acid 3 to about amino acid 9, from about amino acid 10 to about amino acid 16, from about amino acid 26 to about amino acid 32, from about amino acid 30 to about amino acid 36, from about amino acid 112 to about amino acid 118, from about amino acid 166 to about amino acid 172, from about amino acid 212 to about amino acid 218, from about amino acid 224 to about amino acid 230, from about amino acid 230 to about amino acid 236, and from about amino acid 263 to about amino acid 269; a prokaryotic membrane lipoprotein lipid attachment site from about amino acid 44 to about amino acid 55; and a leucine zipper pattern from about amino acid 17 to about amino acid 39.

Clone DNA33221-1133 has been deposited with ATCC on September 16, 1997 and is assigned ATCC deposit no. 209263.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 32 (SEQ ID NO:77), evidenced sequence identity of the PRO224 amino acid sequence to very low-density lipoprotein receptors, apolipoprotein E receptor and chicken oocyte receptors P95. PRO224 has amino acid sequence identity to portions of these proteins in the range from 28% to 45%, and overall identity with these proteins in the range from 33% to 39%.

## EXAMPLE 20

### Isolation of cDNA clones encoding PRO228

A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA28758. An EST proprietary to Genentech, Inc., was employed in the consensus assembly, herein designated DNA21951. Based on the DNA28758 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO228. Three PCR primers (forward and reverse) were synthesized based upon the DNA28758 sequence. Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA28758 sequence.

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primers identified above. A positive library was then used to isolate clones

encoding the PRO228 gene using the probe oligonucleotide and one of the PCR primers.

The oligonucleotide sequences used in the above procedure were the following:

forward PCR primer 1

5'-GGTAATGAGCTCCATTACAG-3' (SEQ ID NO:83)

5 forward PCR primer 2

5'-GGAGTAGAAAGCGCATGG-3' (SEQ ID NO:84)

forward PCR primer 3

5'-CACCTGATACCATGAATGGCAG-3' (SEQ ID NO:85)

reverse PCR primer 1

10 5'-CGAGCTCGAATTAATTTCG-3' (SEQ ID NO:86)

reverse PCR primer 2

5'-GGATCTCCTGAGCTCAGG-3' (SEQ ID NO:87)

reverse PCR primer 3

5'-CCTAGTTGAGTGATCCTTGTAAG-3' (SEQ ID NO:88)

15 hybridization probe

5'-ATGAGACCCACACCTCATGCCGCTGTAATCACCTGACACATTTTGCAATT-3'  
(SEQ ID NO:89)

RNA for construction of the cDNA libraries was isolated from human fetal kidney tissue. The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available  
20 reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes *et al.*, Science, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO228  
25 [herein designated as DNA33092-1202] (Figures 33A-33B, SEQ ID NO:81) and the derived protein sequence for PRO228.

The entire nucleotide sequence of DNA33092-1202 is shown in Figures 33A-33B (SEQ ID NO:81). Clone DNA33092-1202 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 24-26 and ending at the stop codon at nucleotide positions 2094-2096 (Figures 33A-33B). The predicted  
30 polypeptide precursor is 690 amino acids long (Figure 34; SEQ ID NO:82).

Analysis of the full-length PRO228 sequence shown in Figure 34 (SEQ ID NO:82) evidences the presence of important polypeptide domains as shown in Figure 34, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO228 sequence (Figure 34; SEQ ID NO:82) evidences the presence of the following: a signal peptide from about amino acid 1 to about  
35 amino acid 19; transmembrane domains from about amino acid 431 to about amino acid 450, from about amino acid 494 to about amino acid 515, from about amino acid 573 to about amino acid 594, from about amino acid 619 to about amino acid 636 and from about amino acid 646 to about amino acid 664; N-glycosylation sites from about

amino acid 15 to about amino acid 19, from about amino acid 21 to about amino acid 25, from about amino acid 64 to about amino acid 68, from about amino acid 74 to about amino acid 78, from about amino acid 127 to about amino acid 131, from about amino acid 177 to about amino acid 181, from about amino acid 188 to about amino acid 192, from about amino acid 249 to about amino acid 253, from about amino acid 381 to about amino acid 385, and from about amino acid 395 to about amino acid 399; a glycosaminoglycan attachment site from about amino acid 49 to about amino acid 53; a cAMP- and cGMP-dependent protein kinase phosphorylation site from about amino acid 360 to about amino acid 364; tyrosine kinase phosphorylation sites from about amino acid 36 to about amino acid 44 and from about amino acid 670 to about amino acid 677; N-myristoylation sites from about amino acid 38 to about amino acid 44, from about amino acid 50 to about amino acid 56, from about amino acid 52 to about amino acid 58, from about amino acid 80 to about amino acid 86, from about amino acid 382 to about amino acid 388, from about amino acid 388 to about amino acid 394, from about amino acid 434 to about amino acid 440, from about amino acid 480 to about amino acid 486, and from about amino acid 521 to about amino acid 527; and an aspartic acid and asparagine hydroxylation site from about amino acid 75 to about amino acid 87.

Clone DNA33092-1202 has been deposited with ATCC on October 28, 1997 and is assigned ATCC deposit no. 209420.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 34 (SEQ ID NO:82), evidenced sequence identity of the PRO228 amino acid sequence to the secretin related proteins CD97 and EMR1 as well as the secretin member, latrophilin, thereby indicating that PRO228 may be a new member of the secretin related proteins.

## EXAMPLE 21

### Isolation of cDNA clones encoding PRO245

A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA30954. Based on the DNA30954 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO245. PCR primers (forward and reverse) were synthesized based upon the DNA30954 sequence. Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA30954 sequence.

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO245 gene using the probe oligonucleotide and one of the PCR primers.

The oligonucleotide sequences used in the above procedure were the following:

#### forward PCR primer

5'-ATCGTTGTGAAGTTAGTGCCCC-3' (SEQ ID NO:92)

#### reverse PCR primer

5'-ACCTGCGATATCCAACAGAATTG-3' (SEQ ID NO:93)

hybridization probe

5'-GGAAGAGGATACAGTCACTCTGGAAGTATTAGTGGCTCCAGCAGTTCC-3'

(SEQ ID NO:94)

RNA for construction of the cDNA libraries was isolated from human fetal liver tissue. The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes *et al.*, Science, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO245 [herein designated as DNA35638-1141] (Figure 35, SEQ ID NO:90) and the derived protein sequence for PRO245.

The entire nucleotide sequence of DNA35638-1141 is shown in Figure 35 (SEQ ID NO:90). Clone DNA35638-1141 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 89-91 and ending at the stop codon at nucleotide positions 1025-1027 (Figure 35). The predicted polypeptide precursor is 312 amino acids long (Figure 36; SEQ ID NO:91).

Analysis of the full-length PRO245 sequence shown in Figure 36 (SEQ ID NO:91) evidences the presence of important polypeptide domains as shown in Figure 36, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO245 sequence (Figure 36; SEQ ID NO:91) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 20; a transmembrane domain from about amino acid 235 to about amino acid 254; N-glycosylation sites from about amino acid 98 to about amino acid 102, from about amino acid 187 to about amino acid 191, from about amino acid 236 to about amino acid 240, and from about amino acid 277 to about amino acid 281; N-myristoylation sites from about amino acid 182 to about amino acid 188, from about amino acid 239 to about amino acid 245, from about amino acid 255 to about amino acid 261, from about amino acid 257 to about amino acid 263, and from about amino acid 305 to about amino acid 311; and an amidation site from about amino acid 226 to about amino acid 230.

Clone DNA35638-1141 has been deposited with ATCC on September 16, 1997 and is assigned ATCC deposit no. 209265.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 36 (SEQ ID NO:91), evidenced 60% sequence identity of the PRO245 amino acid sequence with the human c-myc protein and, therefore, PRO245 may be a new member of the transmembrane protein receptor tyrosine kinase family.

## EXAMPLE 22

Isolation of cDNA clones encoding PRO246

A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA30955. Based on the DNA30955 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence

of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO246. PCR primers (forward and reverse) were synthesized based upon the DNA30955 sequence. Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA30955 sequence.

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened  
5 by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO246 gene using the probe oligonucleotide and one of the PCR primers.

The oligonucleotide sequences used in the above procedure were the following:

forward PCR primer

5'-AGGGTCTCCAGGAGAAAGACTC-3' (SEQ ID NO:97)

10 reverse PCR primer

5'-ATTGTGGGCCTTGCAGACATAGAC-3' (SEQ ID NO:98)

hybridization probe

5'-GGCCACAGCATCAAAACCTTAGAACTCAATGTACTGGTTCCTCCAGCTCC-3'  
(SEQ ID NO:99)

15 RNA for construction of the cDNA libraries was isolated from human fetal liver tissue. The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not  
20 contain the SfiI site; see, Holmes *et al.*, Science, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO246 [herein designated as DNA35639-1172] (Figure 37, SEQ ID NO:95) and the derived protein sequence for PRO246.

The entire nucleotide sequence of DNA35639-1172 is shown in Figure 37 (SEQ ID NO:95). Clone DNA35639-1172 contains a single open reading frame with an apparent translational initiation site at nucleotide  
25 positions 126-128 and ending at the stop codon at nucleotide positions 1296-1298 (Figure 37). The predicted polypeptide precursor is 390 amino acids long (Figure 38; SEQ ID NO:96).

Analysis of the full-length PRO246 sequence shown in Figure 38 (SEQ ID NO:96) evidences the presence of important polypeptide domains as shown in Figure 38, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO246 sequence (Figure  
30 38; SEQ ID NO:96) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 29; a transmembrane domain from about amino acid 244 to about amino acid 265; N-glycosylation sites from about amino acid 108 to about amino acid 112, from about amino acid 169 to about amino acid 173, from about amino acid 213 to about amino acid 217, from about amino acid 236 to about amino acid 240, and from about amino acid 307 to about amino acid 311; N-myristoylation sites from about amino acid 90 to about amino acid 96,  
35 from about amino acid 167 to about amino acid 173, from about amino acid 220 to about amino acid 226, from about amino acid 231 to about amino acid 237, from about amino acid 252 to about amino acid 258, from about amino acid 256 to about amino acid 262, from about amino acid 262 to about amino acid 268, from about amino

acid 308 to about amino acid 314, from about amino acid 363 to about amino acid 369, and from about amino acid 364 to about amino acid 370; and a prokaryotic membrane lipoprotein lipid attachment site from about amino acid 164 to about amino acid 175.

Clone DNA35639-1172 has been deposited with ATCC on October 17, 1997 and is assigned ATCC deposit  
5 no. 209396.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 38 (SEQ ID NO:96), evidenced sequence identity of the PRO246 amino acid sequence to the human cell surface protein HCAR, thereby indicating that PRO246 may be a novel cell surface virus receptor.

10

## EXAMPLE 23

Isolation of cDNA clones encoding PRO258

A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA28746. Based on the DNA28746 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence  
15 of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO258. One forward PCR primer and two reverse PCR primers were synthesized based upon the DNA28746 sequence. Additionally, two synthetic oligonucleotide hybridization probes were constructed from the consensus DNA28746 sequence.

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones  
20 encoding the PRO258 gene using the probe oligonucleotide and one of the PCR primers.

The oligonucleotide sequences used in the above procedure were the following:

forward PCR primer

5'-GCTAGGAATTCCACAGAAGCCC-3' (SEQ ID NO:102)

reverse PCR primer 1

25 5'-AACCTGGAATGTCACCGAGCTG-3' (SEQ ID NO:103)

reverse PCR primer 2

5'-CCTAGCACAGTGACGAGGGACTTGGC-3' (SEQ ID NO:104)

hybridization probe 1

5'-AAGACACAGCCACCCTAAACTGTCAGTCTTCTGGGAGCAAGCCTGCAGCC-3'  
30 (SEQ ID NO:105)

hybridization probe 2

5'-GCCCTGGCAGACGAGGGCGAGTACACCTGCTCAATCTTCACTATGCCTGT-3'  
(SEQ ID NO:106)

RNA for construction of the cDNA libraries was isolated from human fetal lung tissue. The cDNA libraries  
35 used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with NotI site, linked with blunt to SalI

hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes *et al.*, Science, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO258 [herein designated as DNA35918-1174] (Figure 39, SEQ ID NO:100) and the derived protein sequence for PRO258.

The entire nucleotide sequence of DNA35918-1174 is shown in Figure 39 (SEQ ID NO:100). Clone DNA35918-1174 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 147-149 and ending at the stop codon at nucleotide positions 1341-1343 (Figure 39). The predicted polypeptide precursor is 398 amino acids long (Figure 40; SEQ ID NO:101).

Analysis of the full-length PRO258 sequence shown in Figure 40 (SEQ ID NO:101) evidences the presence of important polypeptide domains as shown in Figure 40, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO258 sequence (Figure 40; SEQ ID NO:101) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 20; a transmembrane domain from about amino acid 331 to about amino acid 352; N-glycosylation sites from about amino acid 25 to about amino acid 29 and from about amino acid 290 to about amino acid 294; N-myristoylation sites from about amino acid 2 to about amino acid 8, from about amino acid 23 to about amino acid 29, from about amino acid 156 to about amino acid 162, from about amino acid 218 to about amino acid 224, from about amino acid 295 to about amino acid 301, from about amino acid 298 to about amino acid 304, from about amino acid 306 to about amino acid 312, from about amino acid 334 to about amino acid 340, from about amino acid 360 to about amino acid 366, from about amino acid 385 to about amino acid 391, and from about amino acid 386 to about amino acid 392; and a prokaryotic membrane lipoprotein lipid attachment site from about amino acid 7 to about amino acid 18.

Clone DNA35918-1174 has been deposited with ATCC on October 17, 1997 and is assigned ATCC deposit no. 209402.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 40 (SEQ ID NO:101), evidenced sequence identity of the PRO258 amino acid sequence to the CRTAM and the poliovirus receptor and contains an Ig domain, thereby indicating that PRO258 is a new member of the Ig superfamily.

30

#### EXAMPLE 24

##### Isolation of cDNA clones encoding PRO261

A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA30843. Based on the DNA30843 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO261. PCR primers (forward and reverse) were synthesized based upon the DNA30843 sequence. Additionally, a synthetic

35

oligonucleotide hybridization probe was constructed from the consensus DNA30843 sequence.

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO261 gene using the probe oligonucleotide and one of the PCR primers.

5 The oligonucleotide sequences used in the above procedure were the following:

forward PCR primer

5'-AAAGGTGCGTACCCAGCTGTGCC-3' (SEQ ID NO:109)

reverse PCR primer

5'-TCCAGTCGGCAGAAGCGGTTCTGG-3' (SEQ ID NO:110)

10 hybridization probe

5'-CCTGGTGTGGATGGCTGTGGCTGCTGCCGGGTATGTGCACGGCGGCTGGG-3'

(SEQ ID NO:111)

RNA for construction of the cDNA libraries was isolated from human fetal lung tissue. The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes *et al.*, *Science*, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

20 DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO261 [herein designated as DNA33473-1176] (Figure 41, SEQ ID NO:107) and the derived protein sequence for PRO261.

The entire nucleotide sequence of DNA33473-1176 is shown in Figure 41 (SEQ ID NO:107). Clone DNA33473-1176 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 10-12 and ending at the stop codon at nucleotide positions 760-762 (Figure 41). The predicted polypeptide precursor is 250 amino acids long (Figure 42; SEQ ID NO:108).

Analysis of the full-length PRO261 sequence shown in Figure 42 (SEQ ID NO:108) evidences the presence of important polypeptide domains as shown in Figure 42, wherein the location given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO261 sequence (Figure 42; SEQ ID NO:108) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 23; N-myristoylation sites from about amino acid 3 to about amino acid 9, from about amino acid 49 to about amino acid 55, from about amino acid 81 to about amino acid 87, from about amino acid 85 to about amino acid 91, from about amino acid 126 to about amino acid 132, from about amino acid 164 to about amino acid 170, from about amino acid 166 to about amino acid 172, from about amino acid 167 to about amino acid 173, from about amino acid 183 to about amino acid 189, and from about amino acid 209 to about amino acid 215; an insulin-like growth factor binding proteins signature from about amino acid 49 to about amino acid 65; a von Willebrand C1 domain from about amino acid 107 to about amino acid 124; a thrombospondin 1 site from about amino acid 201 to about amino acid 216, and an IGF binding protein site from about amino acid 49 to about amino acid 58.



Clone DNA33473-1176 has been deposited with ATCC on October 17, 1997 and is assigned ATCC deposit no. 209391.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 42 (SEQ ID NO:108), evidenced sequence identity of the PRO261 amino acid sequence to the CTGF, thereby indicating that PRO261 is a novel growth factor.

#### EXAMPLE 25

##### Isolation of cDNA clones encoding PRO272

A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA36460. Based on the DNA36460 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO272. PCR primers (forward and reverse) were synthesized based upon the DNA36460 sequence. Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA36460 sequence.

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO272 gene using the probe oligonucleotide and one of the PCR primers.

The oligonucleotide sequences used in the above procedure were the following:

##### forward PCR primer (f.1)

5'-CGCAGGCCCTCATGGCCAGG-3' (SEQ ID NO:114)

##### forward PCR primer (f.2)

5'-GAAATCCTGGGTAATTGG-3' (SEQ ID NO:115)

##### reverse PCR primer

5'-GTGCGCGGTGCTCACAGCTCATC-3' (SEQ ID NO:116)

##### hybridization probe

5'-CCCCCTGAGCGACGCTCCCCATGATGACGCCACGGGAACTTC-3' (SEQ ID NO:117)

RNA for construction of the cDNA libraries was isolated from human fetal lung tissue. The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes *et al.*, Science, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO272 [herein designated as DNA40620-1183] (Figure 43, SEQ ID NO:112) and the derived protein sequence for PRO272.

The entire nucleotide sequence of DNA40620-1183 is shown in Figure 43 (SEQ ID NO:112). Clone

DNA40620-1183 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 35-37 and ending at the stop codon at nucleotide positions 1019-1021 (Figure 43). The predicted polypeptide precursor is 328 amino acids long (Figure 44; SEQ ID NO:113).

Analysis of the full-length PRO272 sequence shown in Figure 44 (SEQ ID NO:113) evidences the presence of important polypeptide domains as shown in Figure 44, wherein the location given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO272 sequence (Figure 44; SEQ ID NO:113) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 20; an N-glycosylation site from about amino acid 140 to about amino acid 144; N-myristoylation sites from about amino acid 263 to about amino acid 269 and from about amino acid 311 to about amino acid 317; and an endoplasmic reticulum targeting sequence from about amino acid 325 to about amino acid 330.

Clone DNA40620-1183 has been deposited with ATCC on October 17, 1997 and is assigned ATCC deposit no. 209388.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 44 (SEQ ID NO:113), evidenced sequence identity of the PRO272 amino acid sequence to the human and mouse reticulocalbin proteins, thereby indicating that PRO272 may be a novel reticulocalbin protein.

#### EXAMPLE 26

##### Isolation of cDNA clones encoding PRO301

A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA35936. Based on the DNA35936 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO301. PCR primers (forward and reverse) were synthesized based upon the DNA35936 sequence. Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA35936 sequence.

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO301 gene using the probe oligonucleotide and one of the PCR primers.

The oligonucleotide sequences used in the above procedure were the following:

##### forward PCR primer (35936.f1)

5'-TCGCGGAGCTGTGTTCTGTTTCCC-3' (SEQ ID NO:120)

##### forward PCR primer (35936.f2)

5'-ACACCTGGTTCAAAGATGGG-3' (SEQ ID NO:121)

##### forward PCR primer (35936.f3)

5'-TTGCCTTACTCAGGTGCTAC-3' (SEQ ID NO:122)

##### reverse PCR primer (35936.r1)

5'-TAGGAAGAGTTGCTGAAGGCACGG-3' (SEQ ID NO:123)

reverse PCR primer (35936.r2)

5'-ACTCAGCAGTGGTAGGAAAG-3'

(SEQ ID NO:124)

hybridization probe (35936.p1)

5'-TGATCGCGATGGGGACAAAGGCGCAAGCTCGAGAGGAACTGTTGTGCCT-3'

5 (SEQ ID NO:125)

RNA for construction of the cDNA libraries was isolated from human fetal kidney tissue. The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes *et al.*, *Science*, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO301 [herein designated as DNA40628-1216] (Figure 45, SEQ ID NO:118) and the derived protein sequence for PRO301.

The entire nucleotide sequence of DNA40628-1216 is shown in Figure 45 (SEQ ID NO:118). Clone DNA40628-1216 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 52-54 and ending at the stop codon at nucleotide positions 949-951 (Figure 45). The predicted polypeptide precursor is 299 amino acids long with a predicted molecular weight of approximately 32,583 daltons and a pI of about 8.29 (Figure 46; SEQ ID NO:119).

Analysis of the full-length PRO301 sequence shown in Figure 46 (SEQ ID NO:119) evidences the presence of important polypeptide domains as shown in Figure 46, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO301 sequence (Figure 46; SEQ ID NO:119) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 27; a transmembrane domain from about amino acid 235 to about amino acid 256; an N-glycosylation site from about amino acid 185 to about amino acid 189; a cAMP- and cGMP-dependent protein kinase phosphorylation site from about amino acid 270 to about amino acid 274; and N-myristoylation sites from about amino acid 105 to about amino acid 111, from about amino acid 116 to about amino acid 122, from about amino acid 158 to about amino acid 164, from about amino acid 219 to about amino acid 225, from about amino acid 237 to about amino acid 243, and from about amino acid 256 to about amino acid 262.

Clone DNA40628-1216 has been deposited with ATCC on November 7, 1997 and is assigned ATCC deposit no. 209432.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 46 (SEQ ID NO:119), evidenced sequence identity of the PRO301 amino acid sequence to A33 antigen precursor (30%) and coxsackie and adenovirus receptor protein (29%).

## EXAMPLE 27

Isolation of cDNA clones encoding PRO322

A consensus DNA sequence was assembled relative to two EST sequences using phrap as described in Example 1 above. The two EST sequences used were "2604309" and "1968970", both from Incyte Pharmaceuticals. The assembled consensus sequence is herein designated "<consen01>". Based on the "<consen01>" consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO322. PCR primers (forward and reverse) were synthesized based upon the "<consen01>" consensus sequence. Additionally, a synthetic oligonucleotide hybridization probe was constructed from the "<consen01>" consensus sequence.

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO322 gene using the probe oligonucleotide and one of the PCR primers.

The oligonucleotide sequences used in the above procedure were the following:

15 forward PCR primer (f.1)

5'-CAGCCTACAGAATAAAGATGGCCCC-3' (SEQ ID NO:128)

reverse PCR primer (r.1)

5'-GGTGCAATGATCTGCCAGGCTGAT-3' (SEQ ID NO:129)

hybridization probe

20 5'-AGAAATACCTGTGGTTCAGTCCATCCCAAACCCCTGCTACAACAGCAG-3'  
(SEQ ID NO:130)

RNA for construction of the cDNA libraries was isolated from human fetal kidney tissue. (LIB227). The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes *et al.*, Science, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO322 [herein designated as DNA48336-1309] (Figure 47, SEQ ID NO:126) and the derived protein sequence for PRO322.

The entire nucleotide sequence of DNA48336-1309 is shown in Figure 47 (SEQ ID NO:126). Clone DNA48336-1309 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 166-168 and ending at the stop codon at nucleotide positions 946-948 (Figure 47). The predicted polypeptide precursor is 260 amino acids long (Figure 48; SEQ ID NO:127). The full-length PRO322 protein shown in Figure 48 has a calculated molecular weight of approximately 28,028 daltons and an estimated pI of approximately 7.87.

Analysis of the full-length PRO322 sequence shown in Figure 48 (SEQ ID NO:127) evidences the presence of important polypeptide domains as shown in Figure 48, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO322 sequence (Figure 48; SEQ ID NO:127), evidences the presence of the following: a putative signal peptide from about amino acid 1 to about amino acid 23; a transmembrane domain from about amino acid 51 to about amino acid 71; a glycosaminoglycan attachment site from about amino acid 155 to about amino acid 159; a tyrosine kinase phosphorylation site from about amino acid 182 to about amino acid 189; N-myristoylation sites from about amino acid 19 to about amino acid 25, from about amino acid 20 to about amino acid 26, from about amino acid 35 to about amino acid 41, from about amino acid 60 to about amino acid 66, from about amino acid 149 to about amino acid 155, from about amino acid 195 to about amino acid 201, from about amino acid 200 to about amino acid 206, from about amino acid 204 to about amino acid 210, from about amino acid 220 to about amino acid 226, from about amino acid 229 to about amino acid 235, and from about amino acid 240 to about amino acid 246; a potential N-glycosylation site from about amino acid 110 to about amino acid 114; a serine proteases, trypsin family and histidine active site from about amino acid 69 to about amino acid 75; a consensus sequence from about amino acid 207 to about amino acid 217; and a kringle domain protein motif from about amino acid 205 to about amino acid 217.

Clone DNA48336-1309 has been deposited with ATCC on March 11, 1998 and is assigned ATCC deposit no. 209669.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 48 (SEQ ID NO:127), evidenced sequence identity of the PRO322 amino acid sequence to neuropsin and other serine proteases, thereby indicating that PRO322 is a novel serine protease related to neuropsin.

#### EXAMPLE 28

##### Isolation of cDNA clones encoding PRO328

A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA35615. Based on the DNA35615 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO328. PCR primers (forward and reverse) were synthesized based upon the DNA35615 sequence. Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA35615 sequence.

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO328 gene using the probe oligonucleotide and one of the PCR primers.

The oligonucleotide sequences used in the above procedure were the following:

forward PCR primer

5'-TCCTGCAGTTTCCTGATGC-3' (SEQ ID NO:133)

reverse PCR primer

5'-CTCATATTGCACACCAGTAATTCG-3' (SEQ ID NO:134)

hybridization probe

5'-ATGAGGAGAAACGTTTGATGGTGGAGCTGCACAACCTCTACCGGG-3'

5 (SEQ ID NO:135)

RNA for construction of the cDNA libraries was isolated from human fetal kidney tissue. The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes *et al.*, Science, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO328 [herein designated as DNA40587-1231] (Figure 49, SEQ ID NO:131) and the derived protein sequence for PRO328.

The entire nucleotide sequence of DNA40587-1231 is shown in Figure 49 (SEQ ID NO:131). Clone DNA40587-1231 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 15-17 and ending at the stop codon at nucleotide positions 1404-1406 (Figure 49). The predicted polypeptide precursor is 463 amino acids long (Figure 50; SEQ ID NO:132).

Analysis of the full-length PRO328 sequence shown in Figure 50 (SEQ ID NO:132) evidences the presence of important polypeptide domains as shown in Figure 50, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO328 sequence (Figure 50; SEQ ID NO:132) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 22; N-glycosylation sites from about amino acid 114 to about amino acid 118, from about amino acid 403 to about amino acid 407 and from about amino acid 409 to about amino acid 413; a glycosaminoglycan attachment site from about amino acid 439 to about amino acid 443; N-myristoylation sites from about amino acid 123 to about amino acid 129, from about amino acid 143 to about amino acid 149, from about amino acid 152 to about amino acid 158, from about amino acid 169 to about amino acid 175, from about amino acid 180 to about amino acid 186, from about amino acid 231 to about amino acid 237, and from about amino acid 250 to about amino acid 256; amidation sites from about amino acid 82 to about amino acid 86 and from about amino acid 172 to about amino acid 176; a peroxidase proximal heme-ligand signature from about amino acid 287 to about amino acid 298; an extracellular protein SCP/Tpx-1/Ag5/PR-1/Sc7 signature 1 from about amino acid 127 to about amino acid 138; and an extracellular protein SCP/Tpx-1/Ag5/PR-1/Sc7 signature 2 from about amino acid 160 to about amino acid 172.

Clone DNA40587-1231 has been deposited with ATCC on November 7, 1997 and is assigned ATCC deposit no. 209438.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 50 (SEQ ID NO:132), evidenced sequence identity of the

PRO328 amino acid sequence to the human glioblastoma protein and to the cysteine rich secretory protein, thereby indicating that PRO328 may be a novel glioblastoma protein or cysteine rich secretory protein.

#### EXAMPLE 29

##### Isolation of cDNA clones encoding PRO331

5 A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. Based on the consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO331. PCR primers (forward and reverse) were synthesized based upon the consensus sequence. Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus  
10 sequence.

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO331 gene using the probe oligonucleotide and one of the PCR primers.

The oligonucleotide sequences used in the above procedure were the following:

##### 15 forward PCR primer

5'-GCCTTTGACAACCTTCAGTCACTAGTGG-3' (SEQ ID NO:138)

##### reverse PCR primer

5'-CCCCATGTGTCCATGACTGTTCCC-3' (SEQ ID NO:139)

##### hybridization probe

20 5'-TACTGCCTCATGACCTCTTCACTCCCTTGCATCATCTTAGAGCGG-3'  
(SEQ ID NO:140)

RNA for construction of the cDNA libraries was isolated from human fetal brain tissue. The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with NotI site, linked with blunt to SalI  
25 hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes *et al.*, Science, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO331 [herein designated as DNA40981-1234] (Figures 51A-51B, SEQ ID NO:136) and the derived protein sequence for  
30 PRO331.

The entire nucleotide sequence of DNA40981-1234 is shown in Figures 51A-51B (SEQ ID NO:136). Clone DNA40981-1234 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 812-814 and ending at the stop codon at nucleotide positions 2732-2734 (Figures 51A-51B). The predicted polypeptide precursor is 640 amino acids long (Figure 52; SEQ ID NO:137).

35 Analysis of the full-length PRO331 sequence shown in Figure 52 (SEQ ID NO:137) evidences the presence of important polypeptide domains as shown in Figure 52, wherein the locations given for those important

polypeptide domains are approximate as described above. Analysis of the full-length PRO331 sequence (Figure 52; SEQ ID NO:137) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 44; a transmembrane domain from about amino acid 525 to about amino acid 545; N-glycosylation sites from about amino acid 278 to about amino acid 282, from about amino acid 364 to about amino acid 368, from about amino acid 390 to about amino acid 394, from about amino acid 412 to about amino acid 416, from about amino acid 415 to about amino acid 419, from about amino acid 434 to about amino acid 438, from about amino acid 442 to about amino acid 446, from about amino acid 488 to about amino acid 492, and from about amino acid 606 to about amino acid 610; a cAMP- and cGMP-dependent protein kinase phosphorylation site from about amino acid 183 to about amino acid 187; and N-myristoylation sites from about amino acid 40 to about amino acid 46, from about amino acid 73 to about amino acid 79, from about amino acid 118 to about amino acid 124, from about amino acid 191 to about amino acid 197, from about amino acid 228 to about amino acid 234, from about amino acid 237 to about amino acid 243, from about amino acid 391 to about amino acid 397, from about amino acid 422 to about amino acid 428, from about amino acid 433 to about amino acid 439, and from about amino acid 531 to about amino acid 537.

Clone DNA40981-1234 has been deposited with ATCC on November 21, 1997 and is assigned ATCC deposit no. 209489.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 52 (SEQ ID NO:137), evidenced sequence identity of the PRO331 amino acid sequence to the LIG-1 protein, thereby indicating that PRO331 may be a novel LIG-1 related protein.

### EXAMPLE 30

#### Isolation of cDNA clones encoding PRO364

An expressed sequence tag (EST) DNA database (LIFESEQ®, Incyte Pharmaceuticals, Palo Alto, CA) was searched and an Incyte EST sequence (Incyte EST no. 3003460) was identified that had homology to members of the tumor necrosis factor receptor (TNFR) family of polypeptides. A consensus DNA sequence was then assembled relative to Incyte EST no. 3003460 and other EST sequences, and is herein designated as DNA44825.

Oligonucleotide probes based upon the above described EST sequence were then synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO364. Forward and reverse PCR primers generally range from 20 to 30 nucleotides and are often designed to give a PCR product of about 100-1000 bp in length. The probe sequences are typically 40-50 bp in length. In order to screen libraries for a full-length clone, DNA from the libraries were screened by PCR amplification, as per Ausubel *et al.*, Current Protocols in Molecular Biology, *supra*, with the PCR primer pair. A positive library was then used to isolate clones encoding the gene of interest using the probe oligonucleotide and one of the primer pairs.

Pairs of PCR primers (forward and reverse) were synthesized:



forward PCR primer (44825.f1)

5'-CACAGCACGGGGCGATGGG-3'

(SEQ ID NO:143)

forward PCR primer (44825.f2)

5'-GCTCTGCGTTCTGCTCTG-3'

(SEQ ID NO:144)

5 forward PCR primer (44825.GITR.f)

5'-GGCACAGCACGGGGCGATGGGCGCGTTT-3'

(SEQ ID NO:145)

reverse PCR primer (44825.r1)

5'-CTGGTCACTGCCACCTTCCTGCAC-3'

(SEQ ID NO:146)

reverse PCR primer (44825.r2)

10 5'-CGCTGACCCAGGCTGAG-3'

(SEQ ID NO:147)

reverse PCR primer (44825.GITR.r)

5'-GAAGGTCCCCGAGGCACAGTCGATACA-3'

(SEQ ID NO:148)

Additionally, synthetic oligonucleotide hybridization probes were constructed from the DNA44825 consensus sequence.

15 hybridization probe (44825.p1)

5'-GAGGAGTGCTGTTCCGAGTGGGACTGCATGTGTGTCCAGC-3'

(SEQ ID NO:149)

hybridization probe (44825.GITR.p)

5'-AGCCTGGGTCAGCGCCCCACCGGGGTCCCGGGTGCGGCC-3'

(SEQ ID NO:150)

RNA for construction of the cDNA libraries was isolated from human small intestine tissue (LIB231). The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes *et al.*, *Science*, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

25 A cDNA clone was identified and sequenced in entirety. The entire nucleotide sequence of DNA47365-1206 is shown in Figure 53 (SEQ ID NO:141). Clone DNA47365-1206 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 121-123, and a stop codon at nucleotide positions 844-846 (Figure 53; SEQ ID NO:141). The predicted polypeptide precursor is 241 amino acids long. The full-length PRO364 protein shown in Figure 54 has an estimated molecular weight of 26,000 daltons and a pI of about 6.34.

30 Analysis of the full-length PRO364 sequence shown in Figure 54 (SEQ ID NO:142) evidences the presence of a variety of important polypeptide domains as shown in Figure 54, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO364 sequence (Figure 54; SEQ ID NO:142) evidences the presence of the following: a putative signal peptide from about amino acid 1 to about amino acid 25; a potential transmembrane domain from about amino acid 162 to about amino acid 180; a potential N-linked glycosylation site from about amino acid 146 to about amino acid 150; N-myristoylation sites from about amino acid 5 to about amino acid 11, from about amino acid 8 to about amino acid 14, from about

amino acid 25 to about amino acid 31, from about amino acid 30 to about amino acid 36, from about amino acid 33 to about amino acid 39, from about amino acid 118 to about amino acid 124, from about amino acid 122 to about amino acid 128, and from about amino acid 156 to about amino acid 162; a prokaryotic membrane lipoprotein lipid attachment site from about amino acid 166 to about amino acid 177; and a leucine zipper pattern from about amino acid 171 to about amino acid 193.

Clone DNA47365-1206 has been deposited with ATCC on November 7, 1997 and is assigned ATCC deposit no. 209436. It is understood that the deposited clone has the actual correct sequence rather than the representations provided herein.

Analysis of the amino acid sequence of the full-length PRO364 polypeptide suggests that portions of it possess significant homology to members of the tumor necrosis factor receptor family, thereby indicating that PRO364 may be a novel member of the tumor necrosis factor receptor family. Based on an ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 54 (SEQ ID NO:142), PRO364 evidences amino acid sequence homology with the mouse GITR protein reported by Nocentini *et al.*, Proc. Natl. Acad. Sci. USA, 94:6216-6221 (1997). It is possible, therefore, that PRO364 represents the human counterpart to the mouse GITR protein reported by Nocentini *et al.*

#### EXAMPLE 31

##### Isolation of cDNA clones encoding PRO366

A clone designated herein as DNA21705 was isolated as described in Example 2 and was found to have certain amino acid sequence similarity or homology with human TNFR1. Based on the DNA21705 sequence, probes were generated and used to screen a human fetal lung library ("LIB25"). Oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO366. Forward and reverse PCR primers generally range from 20 to 30 nucleotides and are often designed to give a PCR product of about 100-1000 bp in length. The probe sequences are typically 40-55 bp in length. In order to screen several libraries for a full-length clone, DNA from the libraries was screened by PCR amplification, as per Ausubel *et al.*, Current Protocols in Molecular Biology, *supra*, with the PCR primer pair. A positive library was then used to isolate clones encoding the gene of interest using the probe oligonucleotide and one of the primer pairs.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO366 and the derived protein sequence for PRO366.

The entire nucleotide sequence of DNA33085-1110 is shown in Figure 55 (SEQ ID NO:151). Clone DNA33085-1110 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 193-195 and ending at the stop codon at nucleotide positions 970-972 (Figure 55). The predicted polypeptide precursor is 259 amino acids long (Figure 56, SEQ ID NO:152) and has a calculated molecular weight of approximately 27,400 daltons. Applicants have shown that the apparent translational initiation site may alternatively be assigned at nucleotide positions 73-75 (identified as amino acid residue -40). The nucleic acid sequences and encoding amino acid sequences comprising the apparent translational initiation site at nucleotide

positions 73-75 and ending at the stop codon at nucleotide positions 970-972 as well as variants thereof as defined above, are also contemplated to be within the scope of this invention.

Sequence analysis indicated an N-terminal signal peptide from about amino acid 1 to about amino acid 29; a transmembrane domain from about amino acid 240 to about amino acid 259; N-glycosylation sites from about amino acid 77 to about amino acid 81, from about amino acid 140 to about amino acid 144, and from about amino acid 156 to about amino acid 160; a cAMP- and cGMP-dependent protein kinase phosphorylation site from about amino acid 126 to about amino acid 130; N-myristoylation sites from about amino acid 56 to about amino acid 62, from about amino acid 72 to about amino acid 78, from about amino acid 114 to about amino acid 120, from about amino acid 154 to about amino acid 160, and from about amino acid 233 to about amino acid 239; two cysteine-rich domains; a sequence that contains four nearly identical 15 amino acid tandem repeats; and a hydrophobic C-terminal region. The hydrophobic C-terminal region is preceded by a pair of small amino acids (Ala223 and Ala224); this structure and the absence of an apparent cytoplasmic domain suggests that PRO366 may be a glycosylphosphatidylinositol (GPI) anchored protein [see, Moran, *J. Biol. Chem.*, 266:1250-1257 (1991)].

TNF receptor family proteins are typically characterized by the presence of multiple (usually four) cysteine-rich domains in their extracellular regions – each cysteine-rich domain being approximately 45 amino acids long and containing approximately 6, regularly spaced, cysteine residues. Based on the crystal structure of the type I TNF receptor, the cysteines in each domain typically form three disulfide bonds in which usually cysteines 1 and 2, 3 and 5, and 4 and 6 are paired together. Like DR4 and Apo-2 (described below), PRO366 contains two extracellular cysteine-rich pseudorepeats, whereas other identified mammalian TNFR family members contain three or more such domains [Smith *et al.*, *Cell*, 76:959 (1994)].

Clone DNA33085-1110 has been deposited with the ATCC on May 30, 1997 and is assigned an ATCC deposit no. 209087. Regarding the sequence, it is understood that the deposited clone contains the correct sequence, and the sequences provided herein are based on known sequencing techniques.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 56 (SEQ ID NO:152), evidenced sequence identity between the PRO366 amino acid sequence and the following: PRO366 shows more sequence identity to DR4 (60%) and Apo-2 (50%) than to other apoptosis-linked receptors, such as Apo-3, TNFR1, or Fas/Apo-1.

### EXAMPLE 32

#### Isolation of cDNA clones encoding PRO535

Use of the signal sequence algorithm described in Example 3 above allowed identification of an EST cluster sequence from the Incyte database, designated herein as ss.clu12694.init. This ss.clu12694.init. EST cluster sequence was then compared to a variety of expressed sequence tag (EST) databases which included public EST databases (*e.g.*, GenBank) and a proprietary EST DNA database (LIFESEQ®, Incyte Pharmaceuticals, Palo Alto, CA) to identify existing homologies. The homology search was performed using the computer program BLAST or BLAST2 (Altschul *et al.*, *Methods in Enzymology*, 266:460-480 (1996)). Those comparisons resulting in a BLAST score of 70 (or in some cases, 90) or greater that did not encode known proteins were clustered and

assembled into a consensus DNA sequence with the program "phrap" (Phil Green, University of Washington, Seattle, Washington). The consensus sequence obtained therefrom is herein designated DNA48352.

In light of an observed sequence homology between the DNA48352 consensus sequence and an EST sequence encompassed within clone no. H86994, from the Merck database, clone H86994 was purchased and the cDNA insert was obtained and sequenced. It was found herein that that insert encoded a full-length protein. The sequence of this cDNA insert is shown in Figure 57 (SEQ ID NO:153) and is herein designated DNA49143-1429.

Clone DNA49143-1429 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 78-80 and ending at the stop codon at nucleotide positions 681-683 (Figure 57). The predicted polypeptide precursor is 201 amino acids long (Figure 58, SEQ ID NO:154). The full-length PRO535 protein shown in Figure 58 has an estimated molecular weight of about 22,180 daltons and a pI of about 9.68.

Analysis of the full-length PRO535 sequence shown in Figure 58 (SEQ ID NO:154) evidences the presence of a variety of important polypeptide domains as shown in Figure 58, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO535 sequence evidences the presence of the following features: a signal peptide from about amino acid 1 to about amino acid 25; a transmembrane domain from about amino acid 155 to about amino acid 174; a potential N-glycosylation site from about amino acid 196 to about amino acid 200; an N-myristoylation site from about amino acid 95 to about amino acid 100; an amidation site from about amino acid 119 to about amino acid 123; and FKBP-type peptidyl-prolyl cis-trans isomer signature sequences from about amino acid 62 to about amino acid 78, from about amino acid 87 to about amino acid 123 and from about amino acid 128 to about amino acid 141. Clone DNA49143-1429 has been deposited with ATCC on June 23, 1998 and is assigned ATCC deposit no. 203013.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 58 (SEQ ID NO:154), evidenced sequence identity between the PRO535 amino acid sequence and the following Dayhoff sequences: S71237, P\_R93551, P\_R28980, S71238, FKB2\_HUMAN, CELC05C8\_1, S55383, S72485, CELC50F2\_6 and S75144.

25

### EXAMPLE 33

#### Isolation of cDNA clones encoding PRO819

Use of the signal sequence algorithm described in Example 3 above allowed identification of an EST cluster sequence from the Incyte database, designated herein as 49605. This EST 49605 cluster sequence was then compared to a variety of expressed sequence tag (EST) databases which included public EST databases (e.g., GenBank) and a proprietary EST DNA database (LIFESEQ®, Incyte Pharmaceuticals, Palo Alto, CA) to identify existing homologies. The homology search was performed using the computer program BLAST or BLAST2 (Altschul *et al.*, Methods in Enzymology, 266:460-480 (1996)). Those comparisons resulting in a BLAST score of 70 (or in some cases, 90) or greater that did not encode known proteins were clustered and assembled into a consensus DNA sequence with the program "phrap" (Phil Green, University of Washington, Seattle, Washington). The consensus sequence obtained therefrom is herein designated DNA56015.

In light of an observed sequence homology between the DNA56015 consensus sequence and an EST

sequence encompassed within clone no. H65785 from the Merck database, clone H65785 was purchased and the cDNA insert was obtained and sequenced. It was found herein that that insert encoded a full-length protein. The sequence of this cDNA insert is shown in Figure 59 (SEQ ID NO:155) and is herein designated DNA57695-1340.

5 Clone DNA57695-1340 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 46-48 and ending at the stop codon at nucleotide positions 202-204 (Figure 59). The predicted polypeptide precursor is 52 amino acids long (Figure 60; SEQ ID NO:156). The full-length PRO819 protein shown in Figure 60 has an estimated molecular weight of about 5,216 daltons and a pI of about 4.67.

Analysis of the full-length PRO819 sequence shown in Figure 60 (SEQ ID NO:156) evidences the presence of a variety of important polypeptide domains as shown in Figure 60, wherein the locations given for those  
10 important polypeptide domains are approximate as described above. Analysis of the full-length PRO819 sequence evidences the presence of the following features: a signal peptide from about amino acid 1 to about amino acid 24; a potential N-myristoylation site from about amino acid 2 to about amino acid 8; and a region having homology to immunoglobulin light chain from about amino acid 5 to about amino acid 33. Clone DNA57695-1340 has been deposited with ATCC on June 23, 1998 and is assigned ATCC deposit no. 203006.

15 An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 60 (SEQ ID NO:156), evidenced sequence identity between the PRO819 amino acid sequence and the following Dayhoff sequences: HSU03899\_1, HUMIGLITEB\_1, VG28\_HSVSA, AF031522\_1, PAD1\_YEAST and AF045484\_1.

#### EXAMPLE 34

##### 20 Isolation of cDNA clones encoding PRO826

Use of the signal sequence algorithm described in Example 3 above allowed identification of an EST cluster sequence from the Incyte database designated herein as 47283. This EST 47283 cluster sequence was then compared to a variety of expressed sequence tag (EST) databases which included public EST databases (e.g., GenBank) and a proprietary EST DNA database (LIFESEQ®, Incyte Pharmaceuticals, Palo Alto, CA) to identify  
25 existing homologies. The homology search was performed using the computer program BLAST or BLAST2 (Altschul *et al.*, Methods in Enzymology, 266:460-480 (1996)). Those comparisons resulting in a BLAST score of 70 (or in some cases, 90) or greater that did not encode known proteins were clustered and assembled into a consensus DNA sequence with the program "phrap" (Phil Green, University of Washington, Seattle, Washington). The consensus sequence obtained therefrom is herein designated DNA56000.

30 In light of an observed sequence homology between the DNA56000 consensus sequence and an EST sequence encompassed within clone no. W69233, from the Merck database, clone no. W69233 was purchased and the cDNA insert was obtained and sequenced. It was found herein that that insert encoded a full-length protein. The sequence of this cDNA insert is shown in Figure 61 (SEQ ID NO:157) and is herein designated DNA57694-1341.

35 Clone DNA57694-1341 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 13-15 and ending at the stop codon at nucleotide positions 310-312 (Figure 61). The predicted polypeptide precursor is 99 amino acids long (Figure 62, SEQ ID NO:158). The full-length PRO826 protein shown

in Figure 62 has an estimated molecular weight of about 11,052 daltons and a pI of about 7.47.

Analysis of the full-length PRO826 sequence shown in Figure 62 (SEQ ID NO:158) evidences the presence of a variety of important polypeptide domains as shown in Figure 62, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO826 sequence evidences the presence of the following features: a signal peptide from about amino acid 1 to about amino acid 22; potential N-myristoylation sites from about amino acid 22 to about amino acid 28 and from about amino acid 90 to about amino acid 96; and an amino acid sequence block having homology to peroxidase from about amino acid 16 to about amino acid 48.

Clone DNA57694-1341 was deposited with the ATCC on June 23, 1998, and is assigned ATCC deposit no. 203017.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 62 (SEQ ID NO:158), evidenced sequence identity between the PRO826 amino acid sequence and the following Dayhoff sequences: CCUI2315\_1, SCU96108\_6, CELF39F10\_4 and HELT\_HELHO.

### EXAMPLE 35

#### Isolation of cDNA clones encoding PRO1160

A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. The consensus sequence is herein designated DNA40650. Based on this DNA40650 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO1160. PCR primers (forward and reverse) were synthesized based upon the DNA40650 consensus sequence. Additionally, a synthetic oligonucleotide hybridization probe was constructed from the DNA40659 consensus sequence.

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification as per Ausubel, *et al.*, Current Protocols in Molecular Biology, *supra*, with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO1160 gene using the probe oligonucleotide and one of the PCR primers.

The oligonucleotide sequences used in the above procedure were the following:

forward PCR primer (40650.f1)

5'-GCTCCCTGATCTTCATGTCACCACC-3' (SEQ ID NO:161)

reverse PCR primer (40650.r1)

5'-CAGGGACACACTCTACCATTCTGGGAG-3' (SEQ ID NO:162)

hybridization probe (40650.p1)

5'-CCATCTTTCTGGTCTCTGCCAGAAATCCGACAACAGCTGCTC-3'  
(SEQ ID NO:163)

RNA for construction of the cDNA libraries was isolated from human breast tissue. The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as

those from Invitrogen, San Diego, CA. The cDNA was primed with NotI site, linked with blunt to Sall hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes *et al.*, Science, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

- 5 DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO1160 [herein designated as DNA62872-1509] (Figure 63, SEQ ID NO:159) and the derived protein sequence for PRO1160.

The entire nucleotide sequence of DNA62872-1509 is shown in Figure 63 (SEQ ID NO:159). Clone DNA62872-1509 contains a single open reading frame with an apparent translational initiation site at nucleotide  
10 positions 40-42 and ending at the stop codon at nucleotide positions 310-312 (Figure 63). The predicted polypeptide precursor is 90 amino acids long (Figure 64; SEQ ID NO:160). The full-length PRO1160 protein shown in Figure 64 has an estimated molecular weight of about 9,039 daltons and a pI of about 4.37.

Analysis of the full-length PRO1160 sequence shown in Figure 64 (SEQ ID NO:160) evidences the presence of a variety of important polypeptide domains as shown in Figure 64, wherein the locations given for those  
15 important polypeptide domains are approximate as described above. Analysis of the full-length PRO1160 sequence evidences the presence of the following features: a signal peptide from about amino acid 1 to about amino acid 19; and a protein kinase C phosphorylation site from about amino acid 68 to about amino acid 70. Clone DNA62872-1509 has been deposited with ATCC on August 4, 1998 and is assigned ATCC deposit no. 203100.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment  
20 analysis of the full-length sequence shown in Figure 64 (SEQ ID NO:160) evidenced sequence identity of the PRO1160 amino acid sequence to the following Dayhoff sequences: B30305, GEN13490, I53641, S53363, HA34\_BRELC, SP96\_DICDI, S36326, SSU51197\_10, MUC1\_XENLA, TCU32448\_1 and AF000409\_1.

### EXAMPLE 36

#### Isolation of cDNA clones encoding PRO1186

25 Use of the signal sequence algorithm described in Example 3 above allowed identification of an EST cluster sequence from the Incyte database. This Incyte EST cluster sequence was then compared to a variety of expressed sequence tag (EST) databases which included public EST databases (*e.g.*, GenBank) and a proprietary EST DNA database (LIFESEQ®, Incyte Pharmaceuticals, Palo Alto, CA) to identify existing homologies. The homology search was performed using the computer program BLAST or BLAST2 (Altschul *et al.*, Methods in Enzymology,  
30 266:460-480 (1996)). Those comparisons resulting in a BLAST score of 70 (or in some cases, 90) or greater that did not encode known proteins were clustered and assembled into a consensus DNA sequence with the program "phrap" (Phil Green, University of Washington, Seattle, Washington). The consensus sequence obtained therefrom is herein designated DNA56748.

In light of an observed sequence homology between the DNA56748 consensus sequence and an EST sequence  
35 no. 3476792 encompassed within a clone (from a library constructed from ovarian tissue) including this Incyte EST, identified from the Incyte database, the Incyte clone including Incyte EST no. 3476792 was purchased and

the cDNA insert was obtained and sequenced. It was found herein that that insert encoded a full-length protein. The sequence of this cDNA insert is shown in Figure 65 (SEQ ID NO:164) and is herein designated DNA60621-1516.

5 Clone DNA60621-1516 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 91-93 and ending at the stop codon at nucleotide positions 406-408 (Figure 65). The predicted polypeptide precursor is 105 amino acids long (Figure 66; SEQ ID NO:165). The full-length PRO1186 protein shown in Figure 66 has an estimated molecular weight of about 11,715 daltons and a pI of about 9.05.

10 Analysis of the full-length PRO1186 sequence shown in Figure 66 (SEQ ID NO:165) evidences the presence of important polypeptide domains as shown in Figure 66, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO1186 sequence evidences the presence of a signal peptide from about amino acid 1 to about amino acid 19; a tyrosine kinase phosphorylation site from about amino acid 88 to about amino acid 95; and N-myristoylation sites from about amino acid 33 to about amino acid 39, from about amino acid 35 to about amino acid 41, and from about amino acid 46 to about amino acid 52. Clone DNA60621-1516 was deposited with the ATCC on August 4, 1998, and is assigned ATCC deposit no. 15 203091.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 66 (SEQ ID NO:165), evidenced sequence identity between the PRO1186 amino acid sequence and the following Dayhoff sequences: VPRA\_DENPO, LFE4\_CHICK, AF034208\_1, AF030433\_1, A55035, COL\_RABIT, CELB0507\_9, S67826\_1, S34665 and CRU73817\_1.

20

#### EXAMPLE 37

##### Isolation of cDNA clones encoding PRO1246

Use of the signal sequence algorithm described in Example 3 above allowed identification of an EST cluster sequence from the Incyte database, designated Incyte EST cluster sequence no. 56853. This Incyte EST cluster sequence no. 56853 was then compared to a variety of expressed sequence tag (EST) databases which included 25 public EST databases (e.g., GenBank) and a proprietary EST DNA database (LIFESEQ<sup>®</sup>, Incyte Pharmaceuticals, Palo Alto, CA) to identify existing homologies. The homology search was performed using the computer program BLAST or BLAST2 (Altshul *et al.*, Methods in Enzymology, 266:460-480 (1996)). Those comparisons resulting in a BLAST score of 70 (or in some cases, 90) or greater that did not encode known proteins were clustered and assembled into a consensus DNA sequence with the program "phrap" (Phil Green, University of Washington, 30 Seattle, Washington). The consensus sequence obtained therefrom is herein designated DNA56021.

In light of an observed sequence homology between the DNA56021 consensus sequence and an EST sequence encompassed within clone no. 2481345, from the Incyte database, clone no. 2481345 was purchased and the cDNA insert was obtained and sequenced. It was found herein that that insert encoded a full-length protein. The sequence of this cDNA insert is shown in Figures 67A-67B (SEQ ID NO:166) and is herein designated DNA64885-1529.

35 Clone DNA64885-1529 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 119-121 and ending at the stop codon at nucleotide positions 1727-1729 (Figures 67A-67B).



The predicted polypeptide precursor is 536 amino acids long (Figure 68; SEQ ID NO:167). The full-length PRO1246 protein shown in Figure 68 has an estimated molecular weight of about 61,450 daltons and a pI of about 9.17.

Analysis of the full-length PRO1246 sequence shown in Figure 68 (SEQ ID NO:167) evidences the presence of a variety of important polypeptide domains as shown in Figure 68, wherein the locations given for those important polypeptide domains are approximate as described above. Analysis of the full-length PRO1246 sequence evidences the presence of the following features: a signal peptide from amino acid 1 to about amino acid 15; a transmembrane domain from about amino acid 347 to about amino acid 365; potential N-glycosylation sites from about amino acid 108 to about amino acid 112, from about amino acid 166 to about amino acid 170, from about amino acid 193 to about amino acid 197, from about amino acid 262 to about amino acid 266, from about amino acid 375 to about amino acid 379, from about amino acid 413 to about amino acid 417, and from about amino acid 498 to about amino acid 502; tyrosine kinase phosphorylation sites from about amino acid 103 to about amino acid 110 and from about amino acid 455 to about amino acid 462; N-myristoylation sites from about amino acid 104 to about amino acid 110, from about amino acid 344 to about amino acid 350, from about amino acid 348 to about amino acid 354, from about amino acid 409 to about amino acid 415, and from about amino acid 496 to about amino acid 502; and amino acid sequence blocks having homology to sulfatase proteins from about amino acid 286 to about amino acid 315, from about amino acid 359 to about amino acid 369 and from amino acid 78 to about amino acid 97. Clone DNA64885-1529 was deposited with the ATCC on November 3, 1998, and is assigned ATCC deposit no. 203457.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in Figure 68 (SEQ ID NO:167) evidenced sequence identity between the PRO1246 amino acid sequence and the following Dayhoff sequences: P\_R51355, CELK09C4\_1, BCU44852\_1, IDS\_HUMAN, G65169, E64903, ARSA\_HUMAN, GL6S\_HUMAN, HSARSF\_1 and GEN12648.

#### EXAMPLE 38

##### Stimulation of Endothelial Cell Proliferation Assay #8

This assay is designed to determine whether PRO polypeptides of the present invention show the ability to stimulate adrenal cortical capillary endothelial cell (ACE) growth. PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of conditions or disorders where angiogenesis would be beneficial including, for example, wound healing, and the like (as would agonists of these PRO polypeptides). Antagonists of the PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of cancerous tumors. PRO1186 showed positive results in this assay.

Bovine adrenal cortical capillary endothelial cells (ACE) (from primary culture, maximum of 12-14 passages) were plated in 96-well plates at 500 cells/well per 100 microliter. Assay media included low glucose DMEM, 10% calf serum, 2 mM glutamine, and 1X penicillin/streptomycin/fungizone. Control wells included the following: (1) no ACE cells added; (2) ACE cells alone; (3) ACE cells plus VEGF (5 ng/ml); and (4) ACE cells plus FGF (5ng/ml). The control or test sample, (in 100 microliter volumes), was then added to the wells (at dilutions of 1%,

0.1% and 0.01%, respectively). The cell cultures were incubated for 6-7 days at 37°C/5% CO<sub>2</sub>. After the incubation, the media in the wells was aspirated, and the cells were washed 1X with PBS. An acid phosphatase reaction mixture (100 microliter; 0.1M sodium acetate, pH 5.5, 0.1% Triton X-100, 10 mM p-nitrophenyl phosphate) was then added to each well. After a 2 hour incubation at 37°C, the reaction was stopped by addition  
 5 of 10 microliters 1N NaOH. Optical density (OD) was measured on a microplate reader at 405 nm.

The activity of PRO1186 was calculated as the fold increase in proliferation (as determined by the acid phosphatase activity, OD 405 nm) relative to (1) cell only background, and (2) relative to maximum stimulation by VEGF. VEGF (at 3-10 ng/ml) and FGF (at 1-5 ng/ml) were employed as an activity reference for maximum stimulation. Results of the assay were considered "positive" if the observed stimulation was ≥ 50% increase over  
 10 background.

PRO1186 assayed "positive" as follows:

1% dilution = 1.75 fold stimulation

0.1% dilution = 1.39 fold stimulation

0.01% dilution = 1.28 fold stimulation

15 Compared to VEGF (5 ng/ml) control:

1% dilution = 1.24 fold stimulation

Compared to FGB (5 ng/ml) control:

1% dilution = 1.46 fold stimulation

#### EXAMPLE 39

#### 20 Inhibition of Vascular Endothelial Growth Factor (VEGF) Stimulated Proliferation of Endothelial Cell Growth

##### Assay #9

The ability of various PRO polypeptides to inhibit VEGF stimulated proliferation of endothelial cells was tested. Polypeptides testing positive in this assay are useful for inhibiting endothelial cell growth in mammals where such an effect would be beneficial, *e.g.*, for inhibiting tumor growth.

25 Specifically, bovine adrenal cortical capillary endothelial cells (ACE) (from primary culture, maximum of 12-14 passages) were plated in 96-well plates at 500 cells/well per 100 microliter. Assay media included low glucose DMEM, 10% calf serum, 2 mM glutamine, and 1X penicillin/streptomycin/fungizone. Control wells included the following: (1) no ACE cells added; (2) ACE cells alone; (3) ACE cells plus 5 ng/ml FGF; (4) ACE cells plus 3 ng/ml VEGF; (5) ACE cells plus 3 ng/ml VEGF plus 1 ng/ml TGF-beta; and (6) ACE cells plus 3 ng/ml  
 30 VEGF plus 5 ng/ml LIF. The test samples, poly-his tagged PRO polypeptides (in 100 microliter volumes), were then added to the wells (at dilutions of 1%, 0.1% and 0.01%, respectively). The cell cultures were incubated for 6-7 days at 37°C/5% CO<sub>2</sub>. After the incubation, the media in the wells was aspirated, and the cells were washed 1X with PBS. An acid phosphatase reaction mixture (100 microliter; 0.1M sodium acetate, pH 5.5, 0.1% Triton X-100, 10 mM p-nitrophenyl phosphate) was then added to each well. After a 2 hour incubation at 37°C, the  
 35 reaction was stopped by addition of 10 microliters 1N NaOH. Optical density (OD) was measured on a microplate reader at 405 nm.

The activity of PRO polypeptides was calculated as the percent inhibition of VEGF (3 ng/ml) stimulated proliferation (as determined by measuring acid phosphatase activity at OD 405 nm) relative to the cells without stimulation. TGF-beta was employed as an activity reference at 1 ng/ml, since TGF-beta blocks 70-90% of VEGF-stimulated ACE cell proliferation. The results, as shown in TABLE 4 below, are indicative of the utility of the PRO polypeptides in cancer therapy and specifically in inhibiting tumor angiogenesis. The numerical values (relative inhibition) shown in TABLE 4 are determined by calculating the percent inhibition of VEGF stimulated proliferation by the PRO polypeptides relative to cells without stimulation and then dividing that percentage into the percent inhibition obtained by TGF- $\beta$  at 1 ng/ml which is known to block 70-90% of VEGF stimulated cell proliferation. The results are considered positive if the PRO polypeptide exhibits 30% or greater inhibition of VEGF stimulation of endothelial cell growth (relative inhibition 30% or greater).

TABLE 4  
Inhibition of VEGF Stimulated Endothelial Cell Growth

	<u>PRO Name</u>	<u>PRO Concentration</u>	<u>Relative % Inhibition</u>
5	PRO172	0.01%	90.0
	PRO172	0.10%	95.0
	PRO172	1.00%	67.0
	PRO175	0.01%	100.0
	PRO175	0.10%	82.0
	PRO175	1.00%	31.0
10	PRO175	0.01%	73.0
	PRO175	0.10%	37.0
	PRO175	1.00%	16.0
15	PRO178	0.01%	101.0
	PRO178	0.10%	101.0
	PRO178	1.00%	54.0
	PRO188	0.01%	35.0
	PRO188	0.10%	34.0
	PRO188	1.00%	82.0
20	PRO188	0.01%	119.0
	PRO188	0.10%	113.0
	PRO188	1.00%	65.0
	PRO179	0.01%	119.0
	PRO179	0.10%	107.0
	PRO179	1.00%	45.0
25	PRO179	0.01%	96.0
	PRO179	0.10%	83.0
	PRO179	1.00%	26.0
30	PRO197	0.01%	90.0
	PRO197	0.10%	79.0
	PRO197	1.00%	68.0
	PRO198	0.01%	48.0
	PRO198	0.10%	50.0
	PRO198	1.00%	64.0
35	PRO198	0.01%	119.0
	PRO198	0.10%	119.0
	PRO198	1.00%	60.0
	PRO182	0.01%	52.0
	PRO182	0.10%	26.0
	PRO182	1.00%	13.0

TABLE 4 Continued  
Inhibition of VEGF Stimulated Endothelial Cell Growth

	<u>PRO Name</u>	<u>PRO Concentration</u>	<u>Relative % Inhibition</u>
5	PRO182	0.099 nM	107.0
	PRO182	0.99 nM	100.0
	PRO182	9.90 nM	66.0
	PRO200	0.01%	96.0
	PRO200	0.10%	90.0
	PRO200	1.00%	52.0
10	PRO200	0.01%	91.0
	PRO200	0.10%	93.0
	PRO200	1.00%	57.0
15	PRO200	0.072 nM	115.0
	PRO200	0.72 nM	113.0
	PRO200	7.2 nM	60.0
	PRO211	0.01%	99.0
	PRO211	0.10%	67.0
	PRO211	1.00%	20.0
20	PRO211	0.01%	109.0
	PRO211	0.10%	95.0
	PRO211	1.00%	27.0
	PRO217	0.01%	106.0
	PRO217	0.10%	84.0
	PRO217	1.00%	39.0
25	PRO217	2.5 nM	20.0
	PRO217	25 nM	88.0
	PRO217	250 nM	58.0
30	PRO219	5.7 nM	61.0
	PRO219	57 nM	109.0
	PRO219	570 nM	97.0
	PRO221	0.01%	103.0
	PRO221	0.10%	82.0
	PRO221	1.00%	31.0
35	PRO221	0.01%	106.0
	PRO221	0.10%	93.0
	PRO221	1.00%	43.0
	PRO224	0.01%	101.0
	PRO224	0.10%	65.0
	PRO224	1.00%	23.0
40	PRO228	0.01%	99.0
	PRO228	0.10%	93.0
	PRO228	1.00%	57.0

TABLE 4 Continued  
Inhibition of VEGF Stimulated Endothelial Cell Growth

	<u>PRO Name</u>	<u>PRO Concentration</u>	<u>Relative % Inhibition</u>
5	PRO228	0.01%	95.0
	PRO228	0.10%	77.0
	PRO228	1.00%	16.0
	PRO228	0.01%	98.0
	PRO228	0.10%	88.0
	PRO228	1.00%	48.0
10	PRO245	0.01%	76.0
	PRO245	0.10%	35.0
	PRO245	1.00%	11.0
15	PRO245	0.48 nM	103.0
	PRO245	4.80 nM	95.0
	PRO245	48 nM	49.0
	PRO246	0.01%	104.0
	PRO246	0.10%	100.0
	PRO246	1.00%	49.0
20	PRO258	0.01%	98.0
	PRO258	0.10%	95.0
	PRO258	1.00%	60.0
	PRO258	0.01%	106.0
	PRO258	0.10%	102.0
	PRO258	1.00%	69.0
25	PRO261	0.01 nM	95.0
	PRO261	0.10 nM	62.0
	PRO261	1.00 nM	50.0
30	PRO261	0.01 nM	110.0
	PRO261	0.10 nM	103.0
	PRO261	1.00 nM	60.0
	PRO272	0.01%	95.0
	PRO272	0.10%	57.0
	PRO272	1.00%	18.0
35	PRO301	0.01%	92.0
	PRO301	0.10%	85.0
	PRO301	1.00%	68.0
	PRO301	7 $\mu$ M	102.0
	PRO301	70 $\mu$ M	88.0
	PRO301	700 $\mu$ M	44.0
40	PRO322	0.01%	92.0
	PRO322	0.10%	91.0
	PRO322	1.00%	49.0

TABLE 4 Continued  
Inhibition of VEGF Stimulated Endothelial Cell Growth

	<u>PRO Name</u>	<u>PRO Concentration</u>	<u>Relative % Inhibition</u>
5	PRO322	0.045 nM	100.0
	PRO322	0.45 nM	111.0
	PRO322	4.50 nM	53.0
	PRO328	0.01%	98.0
	PRO328	0.10%	96.0
	PRO328	1.00%	60.0
10	PRO331	0.01%	88.0
	PRO331	0.10%	82.0
	PRO331	1.00%	56.0
15	PRO366	0.01%	107.0
	PRO366	0.10%	112.0
	PRO366	1.00%	60.0
	PRO366	0.01%	116.0
	PRO366	0.10%	112.0
	PRO366	1.00%	70.0
20	PRO366	0.009 nM	108.0
	PRO366	0.090 nM	106.0
	PRO366	0.900 nM	43.0

#### EXAMPLE 40

##### Induction of c-fos in Endothelial Cells Assay #34

25 This assay is designed to determine whether PRO polypeptides show the ability to induce c-fos in endothelial cells. PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of conditions or disorders where angiogenesis would be beneficial including, for example, wound healing, and the like (as would agonists of these PRO polypeptides). Antagonists of the PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of cancerous tumors.

30 Human venous umbilical vein endothelial cells (HUVEC, Cell Systems) in growth media (50% Ham's F12 w/o GHT: low glucose, and 50% DMEM without glycine: with NaHCO<sub>3</sub>, 1% glutamine, 10 mM HEPES, 10% FBS, 10 ng/ml bFGF) were plated on 96-well microtiter plates at a cell density of  $1 \times 10^4$  cells/well. The day after plating, the cells were starved by removing the growth media and treating the cells with 100  $\mu$ l/well test samples and controls (positive control: growth media; negative control: 10 mM HEPES, 140 mM NaCl, 4% (w/v) mannitol.  
 35 pH 6.8). The cells were incubated for 30 minutes at 37°C, in 5% CO<sub>2</sub>. The samples were removed, and the first part of the bDNA kit protocol (Chiron Diagnostics, cat. #6005-037) was followed, where each capitalized reagent/buffer listed below was available from the kit.

Briefly, the amounts of the TM Lysis Buffer and Probes needed for the tests were calculated based on

information provided by the manufacturer. The appropriate amounts of thawed Probes were added to the TM Lysis Buffer. The Capture Hybridization Buffer was warmed to room temperature. The bDNA strips were set up in the metal strip holders, and 100  $\mu$ l of Capture Hybridization Buffer was added to each b-DNA well needed, followed by incubation for at least 30 minutes. The test plates with the cells were removed from the incubator, and the media  
5 was gently removed using the vacuum manifold. 100  $\mu$ l of Lysis Hybridization Buffer with Probes were quickly pipetted into each well of the microtiter plates. The plates were then incubated at 55°C for 15 minutes. Upon removal from the incubator, the plates were placed on the vortex mixer with the microtiter adapter head and vortexed on the #2 setting for one minute. 80  $\mu$ l of the lysate was removed and added to the bDNA wells containing the Capture Hybridization Buffer, and pipetted up and down to mix. The plates were incubated at 53°C for at least  
10 16 hours.

On the next day, the second part of the bDNA kit protocol was followed. Specifically, the plates were removed from the incubator and placed on the bench to cool for 10 minutes. The volumes of additions needed were calculated based upon information provided by the manufacturer. An Amplifier Working Solution was prepared by making a 1:100 dilution of the Amplifier Concentrate (20 fm/ $\mu$ l) in AL Hybridization Buffer. The hybridization  
15 mixture was removed from the plates and washed twice with Wash A. 50  $\mu$ l of Amplifier Working Solution was added to each well and the wells were incubated at 53°C for 30 minutes. The plates were then removed from the incubator and allowed to cool for 10 minutes. The Label Probe Working Solution was prepared by making a 1:100 dilution of Label Concentrate (40 pmoles/ $\mu$ l) in AL Hybridization Buffer. After the 10-minute cool-down period, the amplifier hybridization mixture was removed and the plates were washed twice with Wash A. 50  $\mu$ l of Label  
20 Probe Working Solution was added to each well and the wells were incubated at 53°C for 15 minutes. After cooling for 10 minutes, the Substrate was warmed to room temperature. Upon addition of 3  $\mu$ l of Substrate Enhancer to each ml of Substrate needed for the assay, the plates were allowed to cool for 10 minutes, the label hybridization mixture was removed, and the plates were washed twice with Wash A and three times with Wash D. 50  $\mu$ l of the Substrate Solution with Enhancer was added to each well. The plates were incubated for 30 minutes  
25 at 37°C and RLU was read in an appropriate luminometer.

The replicates were averaged and the coefficient of variation was determined. The measure of activity of the fold increase over the negative control (HEPES buffer described above) value was indicated by chemiluminescence units (RLU). The results are shown in TABLE 5 below, and are considered positive if the PRO polypeptide exhibits at least a two-fold value over the negative control. Negative control = 1.00 RLU at 1.00%  
30 dilution. Positive control = 8.39 RLU at 1.00% dilution.



TABLE 5  
Induction of c-fos in Endothelial Cells

	<u>PRO Name</u>	<u>PRO Concentration</u>	<u>RLU values</u>
5	PRO356	1.8 nM	2.12
	PRO356	18.0 nM	2.45
	PRO356	180.0 nM	1.40
	PRO356	1.8 nM	2.33
	PRO356	18.0 nM	2.77
	PRO356	180.0 nM	2.05
10	PRO179	0.01%	1.78
	PRO179	0.10%	1.73
	PRO179	1.00%	3.11
15	PRO179	0.10%	1.23
	PRO179	1.00%	1.69
	PRO179	10.00%	2.43
	PRO364	0.086 nM	1.10
	PRO364	0.860 nM	1.43
	PRO364	8.600 nM	1.34
20	PRO364	0.086 nM	3.35
	PRO364	0.860 nM	3.76
	PRO364	8.600 nM	2.19
	PRO535	1.35 nM	1.26
	PRO535	13.50 nM	1.59
	PRO535	135 nM	2.17
25	PRO535	1.35 nM	3.16
	PRO535	13.50 nM	4.03
	PRO535	135 nM	2.26
30	PRO819	0.0232 nM	2.18
	PRO819	0.232 nM	2.12
	PRO819	2.320 nM	1.77
	PRO826	0.11 nM	2.16
	PRO826	1.10 nM	1.87
	PRO826	11.00 nM	2.39
35	PRO1160	0.01%	2.46
	PRO1160	0.10%	2.19
	PRO1160	1.00%	1.40
	PRO1160	0.01%	2.64
	PRO1160	0.10%	2.08
	PRO1160	1.00%	1.44

## EXAMPLE 41

Human Venous Endothelial Cell Ca Flux Assay ASY#68

This assay is designed to determine whether PRO polypeptides show the ability to stimulate calcium flux in human umbilical vein endothelial cells (HUVEC, Cell Systems). Ca influx is a well documented response upon  
 5 binding of certain ligands to their receptors. A test compound that results in a positive response in the present Ca influx assay can be said to bind to a specific receptor and activate a biological signaling pathway in human endothelial cells. This could ultimately lead, for example to cell division, inhibition of cell proliferation, endothelial tube formation, cell migration, apoptosis, etc.

Human venous umbilical vein endothelial cells (HUVEC, Cell Systems) in growth media (50:50 without  
 10 glycine, 1% glutamine, 10mM Hepes, 10% FBS, 10 ng/ml Bfgf), were plated on 96-well microtiter ViewPlates-96 (Packard Instrument Company Part #6005182) microtiter plates at a cell density of  $2 \times 10^4$  cells/well. The day after plating, the cells were washed three times with buffer (HBSS plus 10 mM Hepes), leaving 100  $\mu$ l/well. Then 100  $\mu$ l/well of 8  $\mu$ M Fluo-3 (2x) was added. The cells were incubated for 1.5 hours at 37°C/5% CO<sub>2</sub>. After incubation, the cells were then washed 3x with buffer (described above) leaving 100  $\mu$ l/well. Test samples of the PRO  
 15 polypeptides were prepared on different 96-well plates at 5x concentration in buffer. The positive control corresponded to 50  $\mu$ M ionomycin (5x); the negative control corresponded to Protein 32. Cell plate and sample plates were run on a FLIPR (Molecular Devices) machine. The FLIPR machine added 25  $\mu$ l of test sample to the cells, and readings were taken every second for one minute, then every 3 seconds for the next three minutes.

The fluorescence change from baseline to the maximum rise of the curve ( $\Delta$  change) was calculated, and  
 20 replicates averaged. The rate of fluorescence increase was monitored, and only those samples which had a  $\Delta$  change greater than 1000 and a rise within 60 seconds, were considered positive. In the following TABLE 6, the results are expressed relative to the positive control.

TABLE 6

Human Venous Endothelial Cell Ca Flux Assay

25	<u>PRO Name</u>	<u>PRO Concentration</u>	<u>Relative <math>\Delta</math> in Fluorescence</u>
	PRO179	0.01%	1
	PRO179	0.10%	1
	PRO179	1.00%	3
30	PRO179	0.01%	1
	PRO179	0.10%	1
	PRO179	1.00%	3
	PRO1246	0.007 nM	1
	PRO1246	0.070 nM	1
	PRO1246	0.700 nM	3
35	PRO1246	0.007 nM	1
	PRO1246	0.070 nM	1
	PRO1246	0.700 nM	3

## EXAMPLE 42

Induction of Endothelial Cell Apoptosis ASY#73

The ability of PRO polypeptides to induce apoptosis in endothelial cells was tested in human venous umbilical vein endothelial cells (HUVEC, Cell Systems). A positive test in the assay is indicative of the usefulness of the polypeptide in therapeutically treating tumors as well as vascular disorders where inducing apoptosis of endothelial cells would be beneficial.

The ability of PRO polypeptides to induce apoptosis in endothelial cells was tested in human venous umbilical vein endothelial cells (HUVEC, Cell Systems), using a 96-well format, in 0% serum media supplemented with 100 ng/ml VEGF. (As HUVEC cells are easily dislodged from the plating surface, all pipetting in the wells must be done as gently as practicable.)

The medium was aspirated and the cells washed once with PBS. 5 ml of 1 x trypsin was added to the cells in a T-175 flask, and the cells were allowed to stand until they were released from the plate (about 5-10 minutes). Trypsinization was stopped by adding 5 ml of growth media. The cells were spun at 1000 rpm for 5 minutes at 4°C. The media was aspirated and the cells were resuspended in 10 ml of 10% serum complemented medium (Cell Systems), 1 x penicillin/streptomycin.

The cells were plated on 96-well microtiter plates (Amersham Life Science, cytostar-T scintillating microplate, RPNQ160, sterile, tissue-culture treated, individually wrapped), in 10% serum (CSG-medium, Cell Systems), at a density of  $2 \times 10^4$  cells per well in a total volume of 100  $\mu$ l. PRO228 polypeptide was added in triplicate at dilutions of 1%, 0.33% and 0.11%. Wells without cells were used as a blank and wells with cells only were used as a negative control. As a positive control 1:3 serial dilutions of 50  $\mu$ l of a 3x stock of staurosporine were used. The ability of the PRO228 polypeptide to induce apoptosis was determined using Annexin V, a member of the calcium and phospholipid binding proteins, to detect apoptosis.

0.2 ml Annexin V - Biotin stock solution (100  $\mu$ g/ml) were diluted in 4.6 ml 2 x  $\text{Ca}^{2+}$  binding buffer and 2.5% BSA (1:25 dilution). 50  $\mu$ l of the diluted Annexin V - Biotin solution were added to each well (except controls) to a final concentration of 1.0  $\mu$ g/ml. The samples were incubated for 10-15 minutes with Annexin-Biotin prior to direct addition of  $^{35}\text{S}$ -Streptavidin.  $^{35}\text{S}$ -Streptavidin was diluted in 2x  $\text{Ca}^{2+}$  Binding buffer, 2.5% BSA and was added to all wells at a final concentration of  $3 \times 10^4$  cpm/well. The plates were then sealed, centrifuged at 1000 rpm for 15 minutes and placed on orbital shaker for 2 hours. The analysis was performed on 1450 Microbeta Trilux (Wallac). The results are shown in TABLE 7 below where percent above background represents the percentage amount of counts per minute above the negative controls. Percents greater than or equal to 30% above background are considered positive.

TABLE 7

Induction of Endothelial Cell Apoptosis

	<u>PRO Name</u>	<u>PRO Concentration</u>	<u>Percent Above Background</u>
5	PRO228	0.11%	0.7%
	PRO228	0.11%	47.6%
	PRO228	0.33%	92.2%
	PRO228	0.33%	123.7%
	PRO228	1.00%	51.4%
	PRO228	1.00%	95.3%

10

## EXAMPLE 43

Enhancement of Heart Neonatal Hypertrophy Assay #1

This assay is designed to measure the ability of PRO polypeptides to stimulate hypertrophy of neonatal heart. PRO polypeptides testing positive in this assay are expected to be useful for the therapeutic treatment of various cardiac insufficiency disorders.

- 15 Cardiac myocytes from 1-day old Harlan Sprague Dawley rats were obtained. Cells ( $180 \mu\text{l}$  at  $7.5 \times 10^4/\text{ml}$ , serum <0.1%, freshly isolated) are added on day 1 to 96-well plates previously coated with DMEM/F12 + 4% FCS. Test samples containing the test PRO polypeptide or growth medium only (negative control) ( $20 \mu\text{l}/\text{well}$ ) are added directly to the wells on day 1. PGF ( $20 \mu\text{l}/\text{well}$ ) is then added on day 2 at final concentration of  $10^{-6}$  M. The cells are then stained on day 4 and visually scored on day 5, wherein cells showing no increase in size as compared to negative controls are scored 0.0, cells showing a small to moderate increase in size as compared to negative controls are scored 1.0 and cells showing a large increase in size as compared to negative controls are scored 2.0. The results are shown in TABLE 8 below.

TABLE 8

Enhancement of Heart Neonatal Hypertrophy

	<u>PRO Name</u>	<u>PRO Concentration</u>	<u>Growth Enhancement Score</u>
25	PRO179	0.01%	0
	PRO179	0.10%	0
	PRO179	1.00%	1
30	PRO179	0.01%	0
	PRO179	0.10%	0
	PRO179	1.00%	1
	PRO195	0.01%	0
	PRO195	0.10%	1
	PRO195	1.00%	1
35	PRO224	0.01%	0
	PRO224	0.10%	0
	PRO224	1.00%	1

## EXAMPLE 44

In situ Hybridization

*In situ* hybridization is a powerful and versatile technique for the detection and localization of nucleic acid sequences within cell or tissue preparations. It may be useful, for example, to identify sites of gene expression, analyze the tissue distribution of transcription, identify and localize viral infection, follow changes in specific mRNA synthesis, and aid in chromosome mapping.

*In situ* hybridization was performed following an optimized version of the protocol by Lu and Gillett, Cell Vision, 1: 169-176 (1994), using PCR-generated <sup>33</sup>P-labeled riboprobes. Briefly, formalin-fixed, paraffin-embedded human tissues were sectioned, deparaffinized, deproteinized in proteinase K (20 g/ml) for 15 minutes at 37°C, and further processed for *in situ* hybridization as described by Lu and Gillett, *supra*. A (<sup>33</sup>-P)UTP-labeled antisense riboprobe was generated from a PCR product and hybridized at 55°C overnight. The slides were dipped in Kodak NTB2™ nuclear track emulsion and exposed for 4 weeks.

<sup>33</sup>P-Riboprobe synthesis

6.0 μl (125 mCi) of <sup>33</sup>P-UTP (Amersham BF 1002, SA<2000 Ci/mmol) were speed-vacuum dried. To each tube containing dried <sup>33</sup>P-UTP, the following ingredients were added:

- 2.0 μl 5x transcription buffer
- 1.0 μl DTT (100 mM)
- 2.0 μl NTP mix (2.5 mM: 10 μl each of 10 mM GTP, CTP & ATP + 10 μl H<sub>2</sub>O)
- 1.0 μl UTP (50 μM)
- 1.0 μl RNAsin
- 1.0 μl DNA template (1 μg)
- 1.0 μl H<sub>2</sub>O
- 1.0 μl RNA polymerase (for PCR products T3 = AS, T7 = S, usually)

The tubes were incubated at 37°C for one hour. A total of 1.0 μl RQ1 DNase was added, followed by incubation at 37°C for 15 minutes. A total of 90 μl TE (10 mM Tris pH 7.6/1 mM EDTA pH 8.0) was added, and the mixture was pipetted onto DE81 paper. The remaining solution was loaded in a MICROCON-50™ ultrafiltration unit, and spun using program 10 (6 minutes). The filtration unit was inverted over a second tube and spun using program 2 (3 minutes). After the final recovery spin, a total of 100 μl TE was added, then 1 μl of the final product was pipetted on DE81 paper and counted in 6 ml of BIOFLUOR II™.

The probe was run on a TBE/urea gel. A total of 1-3 μl of the probe or 5 μl of RNA Mrk III was added to 3 μl of loading buffer. After heating on a 95°C heat block for three minutes, the gel was immediately placed on ice. The wells of gel were flushed, and the sample was loaded and run at 180-250 volts for 45 minutes. The gel was wrapped in plastic wrap (SARAN™ brand) and exposed to XAR film with an intensifying screen in a -70°C freezer one hour to overnight.

<sup>33</sup>P-HybridizationA. *Pretreatment of frozen sections*

The slides were removed from the freezer, placed on aluminum trays, and thawed at room temperature for 5

minutes. The trays were placed in a 55°C incubator for five minutes to reduce condensation. The slides were fixed for 10 minutes in 4% paraformaldehyde on ice in the fume hood, and washed in 0.5 x SSC for 5 minutes, at room temperature (25 ml 20 x SSC + 975 ml SQ H<sub>2</sub>O). After deproteinization in 0.5 µg/ml proteinase K for 10 minutes at 37°C (12.5 µl of 10 mg/ml stock in 250 ml prewarmed RNase-free RNase buffer), the sections were washed in 0.5 x SSC for 10 minutes at room temperature. The sections were dehydrated in 70%, 95%, and 100% ethanol, 2 minutes each.

B. *Pretreatment of paraffin-embedded sections*

The slides were deparaffinized, placed in SQ H<sub>2</sub>O, and rinsed twice in 2 x SSC at room temperature, for 5 minutes each time. The sections were deproteinized in 20 µg/ml proteinase K (500 µl of 10 mg/ml in 250 ml RNase-free RNase buffer; 37°C, 15 minutes) for human embryo tissue, or 8 x proteinase K (100 µl in 250 ml RNase buffer, 37°C, 30 minutes) for formalin tissues. Subsequent rinsing in 0.5 x SSC and dehydration were performed as described above.

C. *Prehybridization*

The slides were laid out in a plastic box lined with Box buffer (4 x SSC, 50% formamide) - saturated filter paper. The tissue was covered with 50 µl of hybridization buffer (3.75 g dextran sulfate + 6 ml SQ H<sub>2</sub>O), vortexed, and heated in the microwave for 2 minutes with the cap loosened. After cooling on ice, 18.75 ml formamide, 3.75 ml 20 x SSC, and 9 ml SQ H<sub>2</sub>O were added, and the tissue was vortexed well and incubated at 42°C for 1-4 hours.

D. *Hybridization*

1.0 x 10<sup>6</sup> cpm probe and 1.0 µl tRNA (50 mg/ml stock) per slide were heated at 95°C for 3 minutes. The slides were cooled on ice, and 48 µl hybridization buffer was added per slide. After vortexing, 50 µl <sup>33</sup>P mix was added to 50 µl prehybridization on the slide. The slides were incubated overnight at 55°C.

E. *Washes*

Washing was done for 2x10 minutes with 2xSSC, EDTA at room temperature (400 ml 20 x SSC + 16 ml 0.25 M EDTA, V<sub>f</sub>=4L), followed by RNaseA treatment at 37°C for 30 minutes (500 µl of 10 mg/ml in 250 ml RNase buffer = 20 µg/ml). The slides were washed 2 x10 minutes with 2 x SSC, EDTA at room temperature. The stringency wash conditions were as follows: 2 hours at 55°C, 0.1 x SSC, EDTA (20 ml 20 x SSC + 16 ml EDTA, V<sub>f</sub>=4L).

F. *Oligonucleotides*

*In situ* analysis was performed on twelve of the DNA sequences disclosed herein. The oligonucleotides employed for these analyses are as follows:

(1) DNA19355-1150 (PRO175) (TNF motif homolog)

19355 p1:

5'-TCTAATACGACTCACTATAGCTCAGGGGAAGAGCCAAAGA-3' (SEQ ID NO:168)

19355 p2:

35 5'-TGAATTAACCCTCACTAAAGCAGTGCAATGCAGGGGACTA-3' (SEQ ID NO:169)

(2) DNA23339-1130 (PRO178) (NL8 - TIE ligand homolog)

NL8 p1:

5'-GGATTCTAATACGACTCACTATAGGGCCACGGGCGCTGTGTGCTGGAG-3' (SEQ ID NO:170)

NL8 p2:

5 5'-CTATGAAATTAACCCTCACTAAAGGGATGGTGGGGACCGCAGGGTGAC-3' (SEQ ID NO:171)

NL8 p3:

5'-GGATTCTAATACGACTCACTATAGGGCCCGCCACGAGGAGCTGTTACG-3' (SEQ ID NO:172)

NL8 p4:

5'-CTATGAAATTAACCCTCACTAAAGGGATGACCTGCAGGCATGGGAGAA-3' (SEQ ID NO:173)

10 NL8 p5:

5'-GGATTCTAATACGACTCACTATAGGGCGGCCGCCACGAGGAGCTGTTA-3' (SEQ ID NO:174)

NL8 p6:

5'-CTATGAAATTAACCCTCACTAAAGGGAGGGGCTCTGGGGCTGGGTC-3' (SEQ ID NO:175)

(3) DNA28497-1130 (PRO188) (NL5 - TIE ligand homolog)

15 NL5 p1:

5'-GGATTCTAATACGACTCACTATAGGGCCAACACCAAGGGGCAAGATG-3' (SEQ ID NO:176)

NL5 p2:

5'-CTATGAAATTAACCCTCACTAAAGGGAGGGCTTTTGGTGGGAGAAGTT-3' (SEQ ID NO:177)

(4) DNA29101-1122 (PRO200) (VEGF homolog)

20 29101 p1:

5'-GGATTCTAATACGACTCACTATAGGGCGGCGGAATCCAACCTGAGTAG-3' (SEQ ID NO:178)

29101p2:

5'-CTATGAAATTAACCCTCACTAAAGGGAGCGGCTATCCTCCTGTGCTC-3' (SEQ ID NO:179)

(5) DNA33094-1131 (PRO217) (EGF homolog)

25 33094 p1:

5'-GGATTCTAATACGACTCACTATAGGGCTCAGAAAAGCGCAACAGAGAA-3' (SEQ ID NO:180)

33094 p2:

5'-CTATGAAATTAACCCTCACTAAAGGGATGTCTTCCATGCCAACCTTC-3' (SEQ ID NO:181)

(6) DNA33089-1132 (PRO221) (1 TM receptor homolog)

30 33089 p1:

5'-GGATTCTAATACGACTCACTATAGGGCTGTGCTTTCAATTCTGCCAGTA-3' (SEQ ID NO:182)

33089 p2:

5'-CTATGAAATTAACCCTCACTAAAGGGAGGGTACAATTAAGGGGTGGAT-3' (SEQ ID NO:183)

(7) DNA33221-1133 (PRO224) (LDLR homolog - 1 TM)

33221 p1:

5'-GGATTCTAATACGACTCACTATAGGGCGCAGCGATGGCAGCGATGAGG-3' (SEQ ID NO:184)

33221 p2:

5 5'-CTATGAAATTAACCCTCACTAAAGGGACAGACGGGGCAGCAGGGAGTG-3' (SEQ ID NO:185)

(8) DNA35638-1141 (PRO245) (1 TM receptor homolog)

35638 p1:

5'-GGATTCTAATACGACTCACTATAGGGCGGGAAGATGGCGAGGAGGAG-3' (SEQ ID NO:186)

35638 p2:

10 5'-CTATGAAATTAACCCTCACTAAAGGGACCAAGGCCACAAACGGAAATC-3' (SEQ ID NO:187)

(9) DNA35639-1172 (PRO246) (adenovirus R homolog)

35639 p1:

5'-GGATTCTAATACGACTCACTATAGGGCTTGCTGCGGTTTTTGTTCCTG-3' (SEQ ID NO:188)

35639 p2:

15 5'-CTATGAAATTAACCCTCACTAAAGGGAGCTGCCGATCCCACTGGTATT-3' (SEQ ID NO:189)

(10) DNA35918-1174 (PRO258) (CRTAM homolog)

35918 p1:

5'-GGATTCTAATACGACTCACTATAGGGCCCGCCTCGCTCCTGCTCCTG-3' (SEQ ID NO:190)

35918 p2:

20 5'-CTATGAAATTAACCCTCACTAAAGGGAGGATTGCCGCGACCCTCACAG-3' (SEQ ID NO:191)

(11) DNA33473-1176 (PRO261) (WISP2 - CTGF homolog)

33473 p1:

5'-GGATTCTAATACGACTCACTATAGGGCGCGAGGACGGCGGCTTCA-3' (SEQ ID NO:192)

33473 p2:

25 5'-CTATGAAATTAACCCTCACTAAAGGGAAGAGTCGCGGCCGCCCTTTTT-3' (SEQ ID NO:193)

(12) DNA47365-1206 (PRO364) (novel TNF receptor)

47365 p1:

5'-GGATTCTAATACGACTCACTATAGGGCAACCCGAGCATGGCACAGCAC-3' (SEQ ID NO:194)

47365 p2:

30 5'-CTATGAAATTAACCCTCACTAAAGGGATCTCCAGCCGCCCTTCTC-3' (SEQ ID NO:195)



### G. Results

*In situ* analysis was performed on the above twelve DNA sequences disclosed herein. The results from these analyses are as follows:

(1) DNA19355-1150 (PRO175) (TNF motif homolog)

- 5 In human fetal tissues, there was a specific positive signal over the population of adrenal endothelial cells. Endothelial cells elsewhere were negative.

(2) DNA23339-1130 (PRO178) (NL8 TIE ligand homolog)

- 10 In human fetal tissues, there was a distinctive expression pattern in the lower limb at the connective tissue interface between skeletal muscle and bone (primitive periosteum). Expression adjacent to vascular tissue suggests a possible link with angiogenesis. In thymus tissue, there was an apparent increase in expression in the medulla. Expression in the smooth muscle of the trachea was also observed. In the brain cerebral cortex, an increase in expression appeared to occur in cortical neurons. In the small intestine, expression was observed in smooth muscle and connective tissue of the lamina propria. Expression was also observed in the smooth muscle and connective tissue of the stomach (lamina propria). No expression was observed in the spinal cord, thyroid, adrenals or liver. No expression was observed in the placenta but the vascular smooth muscle of the cord showed positive results. Expression in the body wall showed a similar pattern to lower limb expression.

The highly organized expression patterns in the developing limb, intestine and body wall suggests a distinctive role at these sites indicating a possible role in angiogenesis and patterning.

- 20 Fetal multiblock tissues included: liver, kidney, adrenals, thyroid, lungs, heart, great vessels, small intestine, spleen, thymus, pancreas, brain, spinal cord, body wall, pelvis and lower limb.

Adult multiblock tissues included: liver, kidney, adrenals, myocardium, aorta, spleen, lymph node, pancreas, lung and skin. All tissues examined were negative.

(3) DNA28497-1130 (PRO188) (NL5 TIE ligand homolog)

- 25 In fetal tissues, the lower limb showed a high degree of expression at sites of enchondral bone formation, in osteocytes and in the periosteum/perichondrium of developing bones. In addition, high expression was observed in osteocytes and in the periosteum/perichondrium of developing bones. The expression distribution suggests a role in bone formation and/or differentiation. A faint increase in expression in thyroid epithelial cells was also observed. In addition, the body wall showed a high degree of expression in osteocytes and in the periosteum/perichondrium of developing bones, likewise suggesting a role in bone formation/differentiation. Thymus, trachea, brain (cerebral cortex), spinal cord, small intestine, adrenals, liver, stomach, placenta and cord all showed negative results.

In adult tissues, expression was observed over benign breast epithelium in areas of apocrine metaplasia and sclerosing adenosis. Expression was also seen over infiltrating breast ductal carcinoma cells. Liver, heart and hepatocellular carcinoma showed negative results.

- 35 Possible expression occurred in adult squamous epithelium of skin and in the adult adrenal cortex. All other tissues were negative.

Fetal multiblock tissues included: liver, kidney, adrenal, thyroid, lungs, heart, great vessels, small intestine, spleen, thymus, pancreas, brain, spinal cord, body wall, pelvis and lower limb. Adult multiblock tissues included: liver, kidney, adrenal, myocardium, aorta, spleen, lymph node, pancreas, lung and skin.

(4) DNA29101-1122 (PRO200) (VEGF homolog)

5 *Expression in Human Tissues*

In fetal tissues, expression was observed in developing lower limb bones at the edge of the cartilaginous anlage (*i.e.*, around the outside edge). Also, expression was seen in developing tendons, in vascular smooth muscle and in cells embracing developing skeletal muscle myocytes and myotubes. Expression was also observed at the epiphyseal growth plate. Expression was also seen in the marginal sinus of developing lymph nodes. In addition, expression in the thymus was seen in the subcapsular region of the thymic cortex, possibly representing either the subcapsular epithelial cells or the proliferating thymocytes that are found in this region. Expression in the smooth muscle of the trachea was also reported. In the brain cerebral cortex, focal expression occurred in cortical neurons. Expression was also observed in the following tissues: smooth muscle of the small intestine, generalized expression over the thyroid epithelium, liver expression in ductal plate cells, stomach expression in mural smooth muscle, fetal skin expression in the basal layer of squamous epithelium, placenta expression in interstitial cells (in trophoblastic villi) and cord expression in the wall of arteries and vein. The spleen, spinal cord, and the adrenals showed negative results.

Expression patterns suggest that this VEGF homolog may be involved in cell differentiation and/or proliferation. The expression pattern in developing skeletal muscle suggests a possible role in myoblast differentiation and/or proliferation.

*Expression in Tumors and Fetal Tissues*

High expression was observed at the following sites:

Chimp ovary - granulosa cells of maturing follicles, lower intensity signal observed over thecal cells

25 Chimp parathyroid - high expression over chief cells

Human fetal testis - moderate expression over stromal cells surrounding developing tubules

Human fetal lung - high expression over chondrocytes in developing bronchial tree, and low level expression over branching bronchial epithelium

Specific expression was not observed over the renal cell, gastric and colonic carcinomas.

30 Fetal tissues examined (E12-E16 weeks) included: placenta, umbilical cord, liver, kidney, adrenals, thyroid, lungs, heart, great vessels, esophagus, stomach, small intestine, spleen, thymus, pancreas, brain, eye, spinal cord, body wall, pelvis and lower limb.

Adult tissues examined included: liver, kidney, adrenals, myocardium, aorta, spleen, lymph node, pancreas, lung, skin, cerebral cortex (rm), hippocampus (rm), cerebellum (rm), penis, eye, bladder, gastric carcinoma, colon, colonic carcinoma, chondrosarcoma, acetaminophen induced liver injury and hepatic cirrhosis.

(5) DNA33094-1131 (PRO217) (EGF homolog)

A highly distinctive expression pattern was observed in the human embryo in the outer smooth muscle layer of the GI tract, respiratory cartilage, branching respiratory epithelium, osteoblasts, tendons, gonads, in the optic nerve head and the developing dermis.

5 In adult tissues, expression was seen in the epidermal pegs of the chimp tongue, and the basal epithelial/myoepithelial cells of the prostate and urinary bladder. Also, expression was observed in the alveolar lining cells of the adult lung, mesenchymal cells juxtaposed to erectile tissue in the penis and the cerebral cortex (possibly in glial cells). In the kidney, expression was only seen in disease tissue, specifically in cells surrounding thyroidized renal tubules.

10 Human fetal tissues examined (E12-E16 weeks) included: placenta, umbilical cord, liver, kidney, adrenals, thyroid, lungs, heart, great vessels, esophagus, stomach, small intestine, spleen, thymus, pancreas, brain, eye, spinal cord, body wall, pelvis and lower limb.

Adult human tissues examined included: kidney (normal and end-stage), adrenals, myocardium, aorta, spleen, lymph node, gall bladder, pancreas, lung, skin, eye (including retina), prostate, bladder, and liver (normal, cirrhotic, 15 acute failure).

Non-human primate tissues examined included:

Chimp tissues: salivary gland, stomach, thyroid, parathyroid, skin, thymus, ovary, lymph node

Rhesus Monkey tissues: cerebral cortex, hippocampus, cerebellum and penis

20 (6) DNA33089-1132 (PRO221) (1 TM receptor homolog)

Specific expression was observed over fetal cerebral white and grey matter, as well as over neurons in the spinal cord. The probe appears to cross react with rat. In addition, low level expression occurred over cerebellar neurons in adult rhesus brain. All other tissues were negative.

25 Fetal tissues examined (E12-E16 weeks) included: placenta, umbilical cord, liver, kidney, adrenals, thyroid, lungs, heart, great vessels, esophagus, stomach, small intestine, spleen, thymus, pancreas, brain, eye, spinal cord, body wall, pelvis and lower limb.

Adult tissues examined included: liver, kidney, adrenals, myocardium, aorta, spleen, lymph node, pancreas, lung, skin, cerebral cortex (rm), hippocampus (rm), cerebellum (rm), penis, eye, bladder, stomach, gastric carcinoma, colon, colonic carcinoma, chondrosarcoma, acetaminophen induced liver injury and hepatic cirrhosis.

30 (7) DNA33221-1133 (PRO224) (LDLR homolog - 1 TM)

Expression was limited to the vascular endothelium in fetal spleen, adult spleen, fetal liver, adult thyroid and adult lymph node (in chimp). Additional site of expression was observed in the developing spinal ganglia. All other tissues were negative.

35 Human fetal tissues examined (E12-E16 weeks) included: placenta, umbilical cord, liver, kidney, adrenals, thyroid, lungs, heart, great vessels, esophagus, stomach, small intestine, spleen, thymus, pancreas, brain, eye, spinal

cord, body wall, pelvis and lower limb.

Adult human tissues examined included: kidney (normal and end-stage), adrenals, myocardium, aorta, spleen, lymph node, pancreas, lung skin, eye (including retina), bladder, and liver (normal, cirrhotic, acute failure).

Non-human primate tissues examined included:

- 5       Chimp Tissues: salivary gland, stomach, thyroid, parathyroid, skin, thymus, ovary, lymph node

Rhesus Monkey Tissues: cerebral cortex, hippocampus, cerebellum, penis

(8) DNA35638-1141 (PRO245) (1 TM receptor homolog)

*Expression in human adult and fetal tissues*

- 10       Expression was observed in the endothelium lining of fetal and placental vessels. Endothelial expression was confined to these tissue blocks. Expression was also observed over intermediate trophoblast cells of the placenta. All other tissues were negative.

- 15       Fetal tissues examined (E12-E16 weeks) included: placenta, umbilical cord, liver, kidney, adrenals, thyroid, lungs, heart, great vessels, esophagus, stomach, small intestine, spleen, thymus, pancreas, brain, eye, spinal cord, body wall, pelvis and lower limb.

Adult tissues examined included: liver, kidney, adrenals, myocardium, aorta, spleen, lymph node, pancreas, lung, skin, cerebral cortex (rm), hippocampus (rm), cerebellum (rm), penis, eye, bladder, stomach, gastric carcinoma, colon, colonic carcinoma, thyroid (chimp), parathyroid (chimp), ovary (chimp), chondrosarcoma, acetaminophen induced liver injury and hepatic cirrhosis.

- 20       *Expression in inflamed human tissues (psoriasis, IBD, inflamed kidney, inflamed lung, hepatitis (liver block), normal tonsil, adult and chimp multiblocks)*

- 25       Expression was observed in a subset of fetal vessels in the endothelium and in the placenta but no expression was seen in a variety of normal adult human tissues or vessels. Evaluation of expression of this molecule in vessels of inflamed human tissues were compared to non-inflamed tissues. In this regard, expression was observed in the endothelium/intima of large vessels in the lung afflicted with chronic inflammation, in superficial dermal vessels of psoriatic skin, in arterioles in a specimen of chronic sclerosing nephritis, and in capillaries including the perfollicular sinuses of tonsil. No expression was seen in normal skin (human foreskin specimens), normal lung, inflamed (eight IBD specimens) or normal large bowel, chronically inflamed or cirrhotic liver, normal adult cardiac tissue or adrenal gland.

- 30       (9) DNA35639-1172 (PRO246) (adenovirus R homolog)

Strong expression was observed in fetal vascular endothelium, including tissues of the CNS. A lower level of expression was seen in adult vasculature, including the CNS. Higher levels of expression were not obvious in tumor vascular endothelium. A signal was also seen over bone matrix and adult spleen but not obviously cell associated.

- 35       Human fetal tissues examined (E12-E16 weeks) included: placenta, umbilical cord, liver, kidney, adrenals,

thyroid, lungs, heart, great vessels, esophagus, stomach, small intestine, spleen, thymus, pancreas, brain, eye, spinal cord, body wall, pelvis, testis and lower limb.

Adult human tissues examined included: kidney (normal and end-stage), adrenals, spleen, lymph node, pancreas, lung, eye (including retina), bladder, and liver (normal, cirrhotic, acute failure).

- 5 Non-human primate tissues examined included: Chimp adrenal tissues and Rhesus monkey tissues (cerebral cortex, hippocampus).

(10) DNA35918-1174 (PRO258) (CRTAM homolog)

*Expression in human adult and fetal tissues*

- 10 Strong expression was observed in the nervous system. In the rhesus monkey brain, expression was seen in cortical, cerebellar and hippocampus neurons. Expression was also seen over spinal neurons in the fetal spinal cord, the developing brain and the inner aspects of the fetal retina. In addition, expression occurred over developing dorsal root and autonomic ganglia as well as enteric nerves. In the adult prostate, expression occurred over ganglion cells. In the rat, there was strong expression over the developing hind brain and spinal cord. Strong expression was also seen over interstitial cells in the placental villi. All other tissues were negative.

- 15 Fetal tissues examined (E12-E16 weeks) included: placenta, umbilical cord, liver, kidney, adrenals, thyroid, lungs, heart, great vessels, esophagus, stomach, small intestine, spleen, thymus, pancreas, brain, eye, spinal cord, body wall, pelvis and lower limb.

- 20 Adult tissues examined included: liver, kidney, renal cell carcinoma, adrenals, aorta, spleen, lymph node, pancreas, lung, myocardium, skin, cerebral cortex (rm), hippocampus (rm), cerebellum (rm), bladder, prostate, stomach, gastric carcinoma, colon, colonic carcinoma, thyroid (chimp), parathyroid (chimp), ovary (chimp), chondrosarcoma, acetaminophen induced liver injury and hepatic cirrhosis.

*Expression in normal and damaged Rhesus, Chimp and human brain sections*

- 25 In normal cerebral cortex, expression was observed in all layers in a neuronal pattern. Expression was also observed in pyramidal, granule and polymorphic cell layers of the hippocampus. In cerebellum, mRNA was localized to the granule cell layer and cells in the molecular layer, but not in Purkinje neurons. In infarcted cerebellum, the pattern was slightly different: the granule layer signal was reduced and therefore the Golgi II signal stood out as being strongly expressed. Also, the signal was dispersed around the exterior of areas where Purkinje cells were missing.

*Expression in mouse brains*

- 30 High expression was observed throughout the developing and adult mouse brain but does not appear to show any regional specificity.

(11) DNA33473-1176 (PRO261) (WISP 2 - CTGF homolog)

- 35 Strong expression was observed in dermal fibroblasts in normal adult skin. Strong expression was also seen in two cirrhotic livers at sites of active hepatic fibrosis. In addition, moderate expression was seen over fasciculated cells of the adrenal cortex. Localization of expression supports a role for this factor in extracellular matrix

formation and/or turnover.

*Expression in human breast carcinoma and normal breast tissue and in lung carcinoma*

Expression was negative in benign and malignant epithelial cells in two breast tumors examined, but specific hybridization occurred in mesenchymal cells, particularly in areas of tissue repair including dystrophic ossification.

- 5 The signal seemed to be localized to the same cell population in both breast tumors. Most positive cells had the morphology of fibroblasts, yet smooth muscle cells appeared to be negative. The signal is less intense in lung tumor tissue, however the section examined showed less tissue repair compared with the breast tumor slides. Normal lung and kidney tissue were essentially negative. In summary, expression of mRNA occurred in mesenchymal cells involved in tissue repair and/or collagen deposition. The signal was particularly strong in benign fibroblast-like cells adjacent to either infiltrating breast carcinoma cells or tissue destruction due to benign, inflammatory conditions (duct rupture). Of note, was the fact that deposition of benign osteoid seemed to correlate with strong expression of RNA.

*Expression in normal human colon and colon carcinoma*

- 15 None of the sections examined showed a positive hybridization signal in tumor cells. Positive signals of variable intensity were observed in mesenchymal cells of either fibroblast or smooth muscle differentiation. Fibroblasts with a positive signal were observed adjacent to invasive tumor, if this tumor elicited a so-called desmoplastic response (fibroblastic proliferation with deposition of collagenous fibrosis). Positive smooth muscle cells were seen in mostly arterial vessels of medium size. The positive signals were focally distributed within individual sections but showed variability from section to section.

20 (12) DNA47365-1206 (PRO364) (novel TNF receptor)

In fetal tissues, expression was observed in the fascia lining the anterior surface of the vertebral body. Expression was also seen over the fetal retina. Low expression occurred over fetal neurons. All other tissues were negative.

EXAMPLE 45

- 25 Use of PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 as a Hybridization Probe

- 30 The following method describes use of a nucleotide sequence encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 as a hybridization probe.

- 35 DNA comprising the coding sequence of full-length or mature PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331.

PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 (as shown in Figures 1A-1B, 3A-3B, 5, 7A-7B, 9A-9B, 11A-11B, 13, 15, 17, 19, 21A-21B, 23, 25A-25B, 27A-27B, 29, 31, 33A-33B, 35, 37, 39, 41, 43, 45, 47, 49, 51A-51B, 53, 55, 57, 59, 61, 63, 65 and 67A-67B, respectively, SEQ ID NOS: 1, 8, 10, 15, 20, 25, 30, 35, 40, 45, 50, 56, 61, 66, 71, 76, 81, 90, 95, 100, 107, 112, 118, 126, 131, 136, 141, 151, 153, 155, 157, 159, 164 and 166, respectively) or a fragment thereof is employed as a probe to screen for homologous DNAs (such as those encoding naturally-occurring variants of PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246) in human tissue cDNA libraries or human tissue genomic libraries.

Hybridization and washing of filters containing either library DNAs is performed under the following high-stringency conditions. Hybridization of radiolabeled probe derived from the gene encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide to the filters is performed in a solution of 50% formamide, 5x SSC, 0.1% SDS, 0.1% sodium pyrophosphate, 50 mM sodium phosphate, pH 6.8, 2x Denhardt's solution, and 10% dextran sulfate at 42°C for 20 hours. Washing of the filters is performed in an aqueous solution of 0.1x SSC and 0.1% SDS at 42°C.

DNAs having a desired sequence identity with the DNA encoding full-length native sequence can then be identified using standard techniques known in the art.

#### EXAMPLE 46

Expression of Nucleic Acid Encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 in *E. coli*

This Example illustrates preparation of an unglycosylated form of PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 by recombinant expression in *E. coli*.

The DNA sequence encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 (SEQ ID NOS: 1, 8, 10, 15, 20, 25, 30, 35, 40, 45, 50, 56, 61, 66, 71, 76, 81, 90, 95, 100, 107, 112, 118, 126, 131, 136, 141, 151, 153, 155, 157, 159, 164, or 166, respectively) is initially amplified using selected PCR primers. The primers should contain restriction enzyme sites that correspond to the

restriction enzyme sites on the selected expression vector. A variety of expression vectors may be employed. An example of a suitable vector is pBR322 (derived from *E. coli*; see Bolivar *et al.*, Gene, 2: 95 (1977)), which contains genes for ampicillin and tetracycline resistance. The vector is digested with restriction enzyme and dephosphorylated. The PCR-amplified sequences are then ligated into the vector. The vector will preferably  
 5 include sequences that encode an antibiotic-resistance gene, a trp promoter, a poly-His leader (including the first six STII codons, poly-His sequence, and enterokinase cleavage site), the region encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246, lambda  
 10 transcriptional terminator, and an argU gene.

The ligation mixture is then used to transform a selected *E. coli* strain using the methods described in Sambrook *et al.*, *supra*. Transformants are identified by their ability to grow on LB plates and antibiotic-resistant colonies are then selected. Plasmid DNA can be isolated and confirmed by restriction analysis and DNA sequencing.

15 Selected clones can be grown overnight in liquid culture medium such as LB broth supplemented with antibiotics. The overnight culture may subsequently be used to inoculate a larger-scale culture. The cells are then grown to a desired optical density, during which the expression promoter is turned on.

After culturing the cells for several more hours, the cells can be harvested by centrifugation. The cell pellet obtained by the centrifugation can be solubilized using various agents known in the art, and the solubilized PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide can then be purified using a metal-chelating column under conditions that allow tight binding of the polypeptide.

25 PRO197, PRO195, PRO200, PRO224, PRO301, PRO328, PRO364, PRO366 and PRO1186 were successfully expressed in *E. coli* in a poly-His tagged form by the above procedure.

#### EXAMPLE 47

Expression of Nucleic Acid Encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 in Mammalian Cells

This Example illustrates preparation of a potentially glycosylated form of PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 by recombinant  
 35 expression in mammalian cells.



The vector, pRK5 (*see*, EP 307,247, published March 15, 1989), is employed as the expression vector. Optionally, the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 DNA is ligated into pRK5 with selected restriction enzymes to allow insertion of the DNA encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 using ligation methods such as described in Sambrook *et al.*, *supra*. The resulting vector is called pRK5-(DNA encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246).

In one embodiment, the selected host cells are 293 cells. Human 293 cells (ATCC CCL 1573) are grown to confluence in tissue culture plates in medium such as DMEM supplemented with fetal calf serum and optionally, nutrient components and/or antibiotics. About 10  $\mu$ g DNA of pRK5-(DNA encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246) is mixed with about 1  $\mu$ g DNA encoding the VA RNA gene (Thimmappaya *et al.*, *Cell*, 31: 543 (1982)) and dissolved in 500  $\mu$ l of 1 mM Tris-HCl, 0.1 mM EDTA, 0.227 M CaCl<sub>2</sub>. To this mixture is added, dropwise, 500  $\mu$ l of 50 mM HEPES (pH 7.35), 280 mM NaCl, 1.5 mM NaPO<sub>4</sub>, and a precipitate is allowed to form for 10 minutes at 25°C. The precipitate is suspended and added to the 293 cells and allowed to settle for about four hours at 37°C. The culture medium is aspirated off and 2 ml of 20% glycerol in PBS is added for 30 seconds. The 293 cells are then washed with serum-free medium, fresh medium is added, and the cells are incubated for about 5 days.

Approximately 24 hours after the transfections, the culture medium is removed and replaced with culture medium (alone) or culture medium containing 200  $\mu$ Ci/ml <sup>35</sup>S-cysteine and 200  $\mu$ Ci/ml <sup>35</sup>S-methionine. After a 12-hour incubation, the conditioned medium is collected, concentrated on a spin filter, and loaded onto a 15% SDS gel. The processed gel may be dried and exposed to film for a selected period of time to reveal the presence of the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide. The cultures containing transfected cells may undergo further incubation (in serum-free medium) and the medium is tested in selected bioassays.

In an alternative technique, the gene encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366,

PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 may be introduced into 293 cells transiently using the dextran sulfate method described by Somparyrac *et al.*, Proc. Natl. Acad. Sci., 12: 7575 (1981). 293 cells are grown to maximal density in a spinner flask and 700  $\mu$ g pRK5-(DNA encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246) is added. The cells are first concentrated from the spinner flask by centrifugation and washed with PBS. The DNA-dextran precipitate is incubated on the cell pellet for four hours. The cells are treated with 20% glycerol for 90 seconds, washed with tissue culture medium, and re-introduced into the spinner flask containing tissue culture medium, 5  $\mu$ g/ml bovine insulin, and 0.1  $\mu$ g/ml bovine transferrin. After about four days, the conditioned media is centrifuged and filtered to remove cells and debris. The sample containing the expressed gene encoding the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide can then be concentrated and purified by any selected method, such as dialysis and/or column chromatography.

In another embodiment, the gene encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 can be expressed in CHO cells. The pRK5-(DNA encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246) nucleic acid can be transfected into CHO cells using known reagents such as  $\text{CaPO}_4$  or DEAE-dextran. As described above, the cell cultures can be incubated, and the medium replaced with culture medium (alone) or medium containing a radiolabel such as  $^{35}\text{S}$ -methionine. After determining the presence of PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide, the culture medium may be replaced with serum-free medium. Preferably, the cultures are incubated for about 6 days, and then the conditioned medium is harvested. The medium containing the expressed PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 can then be concentrated and purified by any selected method.

Epitope-tagged gene encoding the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535,

PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide may also be expressed in host CHO cells. The gene encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 may be subcloned out of the pRK5 vector. The subclone insert can undergo PCR amplification to fuse in frame with a selected epitope tag such as a poly-His tag into a baculovirus expression vector. The gene insert encoding the poly-His-tagged-[PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246] can then be subcloned into a SV40- driven vector containing a selection marker such as DHFR for selection of stable clones. Finally, the CHO cells can be transfected (as described above) with the SV40-driven vector. Labeling may be performed, as described above, to verify expression. The culture medium containing the expressed gene encoding the poly-His-tagged-[PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246] can then be concentrated and purified by any selected method, such as by Ni<sup>2+</sup>-chelate affinity chromatography.

PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO245, PRO258, PRO261, PRO301, PRO322, PRO328, PRO364, PRO366, PRO535, PRO826, PRO1160, and PRO1246 were stably expressed in CHO cells by the above described method. In addition, PRO172, PRO178, PRO356, PRO182, PRO245, PRO258, PRO301, PRO322, PRO328, PRO364, PRO366, PRO535 and PRO826 were expressed in CHO cells by a transient procedure.

#### EXAMPLE 48

Expression of Nucleic Acid Encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 in Yeast

The following method describes recombinant expression of the gene encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 in yeast.

First, yeast expression vectors are constructed for intracellular production or secretion of PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 from the ADH2/GAPDH promoter. DNA encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197,

- PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 and the promoter is inserted into suitable restriction enzyme sites in the selected plasmid to direct intracellular expression of the gene encoding PRO172, PRO175, PRO178,
- 5 PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246. For secretion, DNA encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261,
- 10 PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 can be cloned into the selected plasmid, together with DNA encoding the ADH2/GAPDH promoter, a native PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826,
- 15 PRO1160, PRO1186 or PRO1246 signal peptide or other mammalian signal peptide, or, for example, a yeast alpha-factor or invertase secretory signal/leader sequence, and linker sequences (if needed) for expression of the gene encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160,
- 20 PRO1186 or PRO1246.

Yeast cells, such as yeast strain AB110, can then be transformed with the expression plasmids described above and cultured in selected fermentation media. The transformed yeast supernatants can be analyzed by precipitation with 10% trichloroacetic acid and separation by SDS-PAGE, followed by staining of the gels with Coomassie Blue stain.

- 25 Recombinant PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 can subsequently be isolated and purified by removing the yeast cells from the fermentation medium by centrifugation and then concentrating the medium using selected cartridge filters. The
- 30 concentrate containing PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 may further be purified using selected column-chromatography resins.

## EXAMPLE 49

Expression of Nucleic Acid Encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 in Baculovirus-Infected Insect Cells

The following method describes recombinant expression in Baculovirus-infected insect cells.

The sequence coding for PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 is fused upstream of an epitope tag contained within a baculovirus expression vector. Such epitope tags include poly-His tags and immunoglobulin tags (like Fc regions of IgG). A variety of plasmids may be employed, including plasmids derived from commercially available plasmids such as pVL1393 (Novagen). Briefly, the sequence encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 or the desired portion of the coding sequence of PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 [such as the sequence encoding the extracellular domain of a transmembrane protein or the sequence encoding the mature protein if the protein is extracellular] is amplified by PCR with primers complementary to the 5' and 3' regions. The 5' primer may incorporate flanking (selected) restriction enzyme sites. The product is then digested with those selected restriction enzymes and subcloned into the expression vector.

Recombinant baculovirus is generated by co-transfecting the above plasmid and BaculoGold™ virus DNA (Pharmingen) into *Spodoptera frugiperda* ("Sf9") cells (ATCC CRL 1711) using lipofectin (commercially available from GIBCO-BRL). After 4 - 5 days of incubation at 28°C, the released viruses are harvested and used for further amplifications. Viral infection and protein expression are performed as described by O'Reilley *et al.*, Baculovirus Expression Vectors: A Laboratory Manual (Oxford: Oxford University Press (1994)).

Expressed poly-His tagged-[PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246] can then be purified, for example, by Ni<sup>2+</sup>-chelate affinity chromatography as follows. Extracts are prepared from recombinant virus-infected Sf9 cells as described by Rupert *et al.*, Nature, **362**:175-179 (1993). Briefly, Sf9 cells are washed, resuspended in sonication buffer (25 ml Hepes, pH 7.9; 12.5 mM MgCl<sub>2</sub>; 0.1 mM EDTA; 10% glycerol; 0.1% NP-40; 0.4 M KCl), and sonicated twice for 20 seconds on ice. The sonicates are cleared by centrifugation, and the supernatant is diluted 50-fold in loading buffer (50 mM phosphate, 300 mM NaCl, 10% glycerol, pH 7.8) and filtered through a 0.45 µm filter. A Ni<sup>2+</sup>-NTA

agarose column (commercially available from Qiagen) is prepared with a bed volume of 5 ml, washed with 25 ml of water and equilibrated with 25 ml of loading buffer. The filtered cell extract is loaded onto the column at 0.5 ml per minute. The column is washed to baseline  $A_{280}$  with loading buffer, at which point fraction collection is started. Next, the column is washed with a secondary wash buffer (50 mM phosphate; 300 mM NaCl, 10% glycerol, pH 6.0), which elutes non-specifically-bound protein. After reaching  $A_{280}$  baseline again, the column is developed with a 0 to 500 mM imidazole gradient in the secondary wash buffer. One ml fractions are collected and analyzed by SDS-PAGE and silver staining or Western blot with  $Ni^{2+}$ -NTA-conjugated to alkaline phosphatase (Qiagen). Fractions containing the eluted His<sub>10</sub>-tagged-[PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246], respectively, are pooled and dialyzed against loading buffer.

Alternatively, purification of the IgG-tagged (or Fc tagged)-[PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246] can be performed using known chromatography techniques, including for instance, Protein A or protein G column chromatography.

While expression was actually performed in a 0.5-2 L scale, it can be readily scaled up for larger (e.g., 8 L) preparations. The proteins were expressed as an IgG construct (immunoadhesin), in which the protein extracellular region was fused to an IgG1 constant region sequence containing the hinge, CH2 and CH3 domains and/or in poly-His tagged forms.

Following PCR amplification, the respective coding sequences were subcloned into a baculovirus expression vector (pb.PH.IgG for IgG fusions and pb.PH.His.c for poly-His tagged proteins), and the vector and Baculogold® baculovirus DNA (Pharmingen) were co-transfected into 105 *Spodoptera frugiperda* ("Sf9") cells (ATCC CRL 1711), using Lipofectin (Gibco BRL). pb.PH.IgG and pb.PH.His are modifications of the commercially available baculovirus expression vector pVL1393 (Pharmingen), with modified polylinker regions to include the His or Fc tag sequences. The cells were grown in Hink's TNM-FH medium supplemented with 10% FBS (Hyclone). Cells were incubated for 5 days at 28°C. The supernatant was harvested and subsequently used for the first viral amplification by infecting Sf9 cells in Hink's TNM-FH medium supplemented with 10% FBS at an approximate multiplicity of infection (MOI) of 10. Cells were incubated for 3 days at 28°C. The supernatant was harvested and the expression of the constructs in the baculovirus expression vector was determined by batch binding of 1 ml of supernatant to 25 ml of  $Ni^{2+}$ -NTA beads (QIAGEN) for histidine tagged proteins or Protein-A Sepharose CL-4B beads (Pharmacia) for IgG tagged proteins followed by SDS-PAGE analysis comparing to a known concentration of protein standard by Coomassie blue staining.

The first viral amplification supernatant was used to infect a spinner culture (500 ml) of Sf9 cells grown in ESF-921 medium (Expression Systems LLC) at an approximate MOI of 0.1. Cells were incubated for 3 days at 28°C. The supernatant was harvested and filtered. Batch binding and SDS-PAGE analysis was repeated, as

necessary, until expression of the spinner culture was confirmed.

The conditioned medium from the transfected cells (0.5 to 3 L) was harvested by centrifugation to remove the cells and filtered through 0.22 micron filters. For the poly-His tagged constructs, the protein construct were purified using a Ni<sup>2+</sup>-NTA column (Qiagen). Before purification, imidazole was added to the conditioned media to a concentration of 5 mM. The conditioned media were pumped onto a 6 ml Ni<sup>2+</sup>-NTA column equilibrated in 20 mM Hepes, pH 7.4, buffer containing 0.3 M NaCl and 5 mM imidazole at a flow rate of 4-5 ml/min. at 4°C. After loading, the column was washed with additional equilibration buffer and the protein eluted with equilibration buffer containing 0.25 M imidazole. The highly purified protein was subsequently desalted into a storage buffer containing 10 mM Hepes, 0.14 M NaCl and 4% mannitol, pH 6.8, with a 25 ml G25 Superfine (Pharmacia) column and stored at -80°C.

Immunoaderhin (Fc containing) constructs of proteins were purified from the conditioned media as follows. The conditioned media were pumped onto a 5 ml Protein A column (Pharmacia) which had been equilibrated in 20 mM Na phosphate buffer, pH 6.8. After loading, the column was washed extensively with equilibration buffer before elution with 100 mM citric acid, pH 3.5. The eluted protein was immediately neutralized by collecting 1 ml fractions into tubes containing 275 ml of 1 M Tris buffer, pH 9. The highly purified protein was subsequently desalted into storage buffer as described above for the poly-His tagged proteins. The homogeneity of the proteins was verified by SDS polyacrylamide gel (PEG) electrophoresis and N-terminal amino acid sequencing by Edman degradation.

PRO172, PRO175, PRO178, PRO356, PRO228, PRO258, PRO272, PRO301, PRO322, PRO819, and PRO826 were successfully expressed in Baculovirus- infected insect Sf9 cells by the above procedure.

Alternatively, a modified baculovirus procedure may be used incorporating high-5 cells. In this procedure, the DNA encoding the desired sequence was amplified with suitable systems, such as Pfu (Stratagene), or fused upstream (5'-of) of an epitope tag contained with a baculovirus expression vector. Such epitope tags include poly-His tags and immunoglobulin tags (like Fc regions of IgG). A variety of plasmids may be employed, including plasmids derived from commercially available plasmids such as pIE1-1 (Novagen). The pIE1-1 and pIE1-2 vectors are designed for constitutive expression of recombinant proteins from the baculovirus iel promoter in stably-transformed insect cells. The plasmids differ only in the orientation of the multiple cloning sites and contain all promoter sequences known to be important for iel-mediated gene expression in uninfected insect cells as well as the hr5 enhancer element. pIE1-1 and pIE1-2 include the translation initiation site and can be used to produce fusion proteins. Briefly, the desired sequence or the desired portion of the sequence (such as the sequence encoding the extracellular domain of a transmembrane protein) was amplified by PCR with primers complementary to the 5' and 3' regions. The 5' primer may incorporate flanking (selected) restriction enzyme sites. The product was then digested with those selected restriction enzymes and subcloned into the expression vector. For example, derivatives of pIE1-1 can include the Fc region of human IgG (pb.PH.IgG) or an 8 histidine (pb.PH.His) tag downstream (3'-of) the desired sequence. Preferably, the vector construct is sequenced for confirmation.

High-5 cells are grown to a confluency of 50% under the conditions of, 27°C, no CO<sub>2</sub>, NO pen/strep. For each 150 mm plate, 30 µg of pIE based vector containing the sequence was mixed with 1 ml Ex-Cell medium (Media:

Ex-Cell 401 + 1/100 L-Glu JRH Biosciences #14401-78P (note: this media is light sensitive)), and in a separate tube, 100  $\mu$ l of CellFectin (CellFECTIN (GibcoBRL #10362-010) (vortexed to mix)) was mixed with 1 ml of Ex-Cell medium. The two solutions were combined and allowed to incubate at room temperature for 15 minutes. 8 ml of Ex-Cell media was added to the 2 ml of DNA/CellFECTIN mix and this is layered on high-5 cells that have been washed once with Ex-Cell media. The plate is then incubated in darkness for 1 hour at room temperature. The DNA/CellFECTIN mix is then aspirated, and the cells are washed once with Ex-Cell to remove excess CellFECTIN, 30 ml of fresh Ex-Cell media was added and the cells are incubated for 3 days at 28°C. The supernatant was harvested and the expression of the sequence in the baculovirus expression vector was determined by batch binding of 1 ml of supernatant to 25 ml of Ni<sup>2+</sup>-NTA beads (QIAGEN) for histidine tagged proteins or Protein-A Sepharose CL-4B beads (Pharmacia) for IgG tagged proteins followed by SDS-PAGE analysis comparing to a known concentration of protein standard by Coomassie blue staining.

The conditioned media from the transfected cells (0.5 to 3 L) was harvested by centrifugation to remove the cells and filtered through 0.22 micron filters. For the poly-His tagged constructs, the protein comprising the sequence is purified using a Ni<sup>2+</sup>-NTA column (Qiagen). Before purification, imidazole is added to the conditioned media to a concentration of 5 mM. The conditioned media was pumped onto a 6 ml Ni<sup>2+</sup>-NTA column equilibrated in 20 mM Hepes, pH 7.4, buffer containing 0.3 M NaCl and 5 mM imidazole at a flow rate of 4-5 ml/min. at 48°C. After loading, the column was washed with additional equilibration buffer and the protein eluted with equilibration buffer containing 0.25 M imidazole. The highly purified protein was then subsequently desalted into a storage buffer containing 10 mM Hepes, 0.14 M NaCl and 4% mannitol, pH 6.8, with a 25 ml G25 Superfine (Pharmacia) column and stored at -80°C.

Immunoadhesin (Fc containing) constructs of proteins were purified from the conditioned media as follows. The conditioned media was pumped onto a 5 ml Protein A column (Pharmacia) which had been equilibrated in 20 mM Na phosphate buffer, pH 6.8. After loading, the column was washed extensively with equilibration buffer before elution with 100 mM citric acid, pH 3.5. The eluted protein was immediately neutralized by collecting 1 ml fractions into tubes containing 275 ml of 1 M Tris buffer, pH 9. The highly purified protein was subsequently desalted into storage buffer as described above for the poly-His tagged proteins. The homogeneity of the sequence was assessed by SDS polyacrylamide gels and by N-terminal amino acid sequencing by Edman degradation and other analytical procedures as desired or necessary.

PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, and PRO1186 were expressed in high 5 cells by the above described method.



## EXAMPLE 50

Preparation of Antibodies that Bind PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246

This Example illustrates preparation of monoclonal antibodies that can specifically bind PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246.

Techniques for producing the monoclonal antibodies are known in the art and are described, for instance, in Goding, *supra*. Immunogens that may be employed include purified PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 fusion proteins containing PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246, and cells expressing the gene encoding PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 on the cell surface. Selection of the immunogen can be made by the skilled artisan without undue experimentation.

Mice, such as Balb/c, are immunized with the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 immunogen emulsified in complete Freund's adjuvant and injected subcutaneously or intraperitoneally in an amount from 1 to 100 micrograms. Alternatively, the immunogen is emulsified in MPL-TDM adjuvant (Ribi Immunochemical Research, Hamilton, MT) and injected into the animal's hind foot pads. The immunized mice are then boosted 10 to 12 days later with additional immunogen emulsified in the selected adjuvant. Thereafter, for several weeks, the mice may also be boosted with additional immunization injections. Serum samples may be periodically obtained from the mice by retro-orbital bleeding for testing in ELISA assays to detect anti-PRO172, anti-PRO175, anti-PRO178, anti-PRO188, anti-PRO356, anti-PRO179, anti-PRO197, anti-PRO198, anti-PRO182, anti-PRO195, anti-PRO200, anti-PRO211, anti-PRO217, anti-PRO219, anti-PRO221, anti-PRO224, anti-PRO228, anti-PRO245, anti-PRO246, anti-PRO258, anti-PRO261, anti-PRO272, anti-PRO301, anti-PRO322, anti-PRO328, anti-PRO331, anti-PRO364, anti-PRO366, anti-PRO535, anti-PRO819, anti-PRO826, anti-PRO1160, anti-PRO1186 or anti-PRO1246 antibodies.

After a suitable antibody titer has been detected, the animals "positive" for antibodies can be injected with a

final intravenous injection of PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246. Three to four days later, the mice are sacrificed and the spleen cells are harvested. The spleen cells are then fused (using 35% polyethylene glycol) to a selected murine myeloma cell line such as P3X63AgU.1, available from ATCC, No. CRL 1597. The fusions generate hybridoma cells that can then be plated in 96-well tissue culture plates containing HAT medium to inhibit proliferation of non-fused cells, myeloma hybrids, and spleen cell hybrids.

The hybridoma cells will be screened in an ELISA for reactivity against PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246. Determination of "positive" hybridoma cells secreting the desired monoclonal antibodies against PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 is within the skill in the art.

The positive hybridoma cells can be injected intraperitoneally into syngeneic Balb/c mice to produce ascites containing the anti-PRO172, anti-PRO175, anti-PRO178, anti-PRO188, anti-PRO356, anti-PRO179, anti-PRO197, anti-PRO198, anti-PRO182, anti-PRO195, anti-PRO200, anti-PRO211, anti-PRO217, anti-PRO219, anti-PRO221, anti-PRO224, anti-PRO228, anti-PRO245, anti-PRO246, anti-PRO258, anti-PRO261, anti-PRO272, anti-PRO301, anti-PRO322, anti-PRO328, anti-PRO331, anti-PRO364, anti-PRO366, anti-PRO535, anti-PRO819, anti-PRO826, anti-PRO1160, anti-PRO1186 or anti-PRO1246 monoclonal antibodies. Alternatively, the hybridoma cells can be grown in tissue-culture flasks or roller bottles. Purification of the monoclonal antibodies produced in the ascites can be accomplished using ammonium-sulfate precipitation, followed by gel-exclusion chromatography. Alternatively, affinity chromatography based upon binding of antibody to protein A or protein G can be employed.

#### Deposit of Material

The following material(s) has/have been deposited with the American Type Culture Collection, 10801 University Blvd., Manassas, VA 20110-2209, USA (ATCC):

Material	ATCC Dep. No.	Deposit Date
DNA35916-1161	209419	October 28, 1997
DNA19355-1150	209466	November 18, 1997
DNA23339-1130	209282	September 18, 1997
DNA28497-1130	209279	September 18, 1997
DNA47470-1130-P1	209422	October 28, 1997
DNA16451-1078	209281	September 18, 1997

	DNA22780-1078	209284	September 18, 1997
	DNA33457-1078	209283	September 18, 1997
	DNA27865-1091	209296	September 23, 1997
	DNA26847-1395	209772	April 14, 1998
5	DNA29101-1122	209653	March 5, 1998
	DNA32292-1131	209258	September 16, 1997
	DNA33094-1131	209256	September 16, 1997
	DNA32290-1164	209384	October 16, 1997
	DNA33089-1132	209262	September 16, 1997
10	DNA33221-1133	209263	September 16, 1997
	DNA33092-1202	209420	October 28, 1997
	DNA35638-1141	209265	September 16, 1997
	DNA35639-1172	209396	October 17, 1997
	DNA35918-1174	209402	October 17, 1997
15	DNA33473-1176	209391	October 17, 1997
	DNA40620-1183	209388	October 17, 1997
	DNA40628-1216	209432	November 7, 1997
	DNA48336-1309	209669	March 11, 1998
	DNA40587-1231	209438	November 7, 1997
20	DNA40981-1234	209489	November 21, 1997
	DNA47365-1206	209436	November 7, 1997
	DNA33085-1110	209087	May 30, 1997
	DNA49143-1429	203013	June 23, 1998
	DNA57695-1340	203006	June 23, 1998
25	DNA57694-1341	203017	June 23, 1998
	DNA62872-1509	203100	August 4, 1998
	DNA60621-1516	203091	August 4, 1998
	DNA64885-1529	203457	November 3, 1998

This deposit was made under the provisions of the Budapest Treaty on the International Recognition of the  
 30 Deposit of Microorganisms for the Purpose of Patent Procedure and the Regulations thereunder (Budapest Treaty).  
 This assures maintenance of a viable culture of the deposit for 30 years from the date of deposit. The deposit will  
 be made available by ATCC under the terms of the Budapest Treaty, and subject to an agreement between  
 Genentech, Inc., and ATCC, which assures permanent and unrestricted availability of the progeny of the culture  
 of the deposit to the public upon issuance of the pertinent U.S. patent or upon laying open to the public of any U.S.  
 35 or foreign patent application, whichever comes first, and assures availability of the progeny to one determined by  
 the U.S. Commissioner of Patents and Trademarks to be entitled thereto according to 35 USC §122 and the

Commissioner's rules pursuant thereto (including 37 CFR §1.14 with particular reference to 886 OG 638).

5 The assignee of the present application has agreed that if a culture of the material(s) on deposit should die or be lost or destroyed when cultivated under suitable conditions, the material(s) will be promptly replaced on notification with another of the same. Availability of the deposited material(s) is not to be construed as a license to practice the invention in contravention of the rights granted under the authority of any government in accordance with its patent laws.

10 The foregoing written specification is considered to be sufficient to enable one skilled in the art to practice the invention. The present invention is not to be limited in scope by the construct(s) deposited, since the deposited embodiment(s) is/are intended as single illustration(s) of certain aspects of the invention and any constructs that are functionally equivalent are within the scope of this invention. The deposit of material(s) herein does not constitute an admission that the written description herein contained is inadequate to enable the practice of any aspect of the invention, including the best mode thereof, nor is it to be construed as limiting the scope of the claims to the specific illustrations that it represents. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and fall within the scope of the appended claims.

15

WHAT IS CLAIMED IS:

1. A composition comprising a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide or agonist or antagonist thereof, in admixture with a pharmaceutically acceptable carrier.
2. The composition of Claim 1 comprising a therapeutically effective amount of said polypeptide or said agonist or antagonist thereof.
3. The composition of Claim 1, wherein the agonist is an anti-PRO172, anti-PRO175, anti-PRO178, anti-PRO188, anti-PRO356, anti-PRO179, anti-PRO197, anti-PRO198, anti-PRO182, anti-PRO195, anti-PRO200, anti-PRO211, anti-PRO217, anti-PRO219, anti-PRO221, anti-PRO224, anti-PRO228, anti-PRO245, anti-PRO246, anti-PRO258, anti-PRO261, anti-PRO272, anti-PRO301, anti-PRO322, anti-PRO328, anti-PRO331, anti-PRO364, anti-PRO366, anti-PRO535, anti-PRO819, anti-PRO826, anti-PRO1160, anti-PRO1186 or anti-PRO1246 antibody.
4. The composition of Claim 1, wherein the antagonist is an anti-PRO172, anti-PRO175, anti-PRO178, anti-PRO188, anti-PRO356, anti-PRO179, anti-PRO197, anti-PRO198, anti-PRO182, anti-PRO195, anti-PRO200, anti-PRO211, anti-PRO217, anti-PRO219, anti-PRO221, anti-PRO224, anti-PRO228, anti-PRO245, anti-PRO246, anti-PRO258, anti-PRO261, anti-PRO272, anti-PRO301, anti-PRO322, anti-PRO328, anti-PRO331, anti-PRO364, anti-PRO366, anti-PRO535, anti-PRO819, anti-PRO826, anti-PRO1160, anti-PRO1186 or anti-PRO1246 antibody.
5. The composition of Claim 1 further comprising a cardiovascular, endothelial, angiogenic or angiostatic agent.
6. A method of preparing the composition of Claim 1 comprising admixing a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide or agonist or antagonist thereof, with a pharmaceutically acceptable carrier.
7. An article of manufacture comprising:
  - (1) a composition comprising (a) a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide, (b) an agonist of a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219,

PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide, or (c) an antagonist of a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide, in admixture with a pharmaceutically acceptable carrier;

- (2) a container containing said composition; and
- (3) a label affixed to said container, or a package insert included in said container, referring to the use of said composition, in the treatment of a cardiovascular, endothelial, and angiogenic disorder.

8. The article of manufacture of Claim 7, wherein said agonist is an anti-PRO172, anti-PRO175, anti-PRO178, anti-PRO188, anti-PRO356, anti-PRO179, anti-PRO197, anti-PRO198, anti-PRO182, anti-PRO195, anti-PRO200, anti-PRO211, anti-PRO217, anti-PRO219, anti-PRO221, anti-PRO224, anti-PRO228, anti-PRO245, anti-PRO246, anti-PRO258, anti-PRO261, anti-PRO272, anti-PRO301, anti-PRO322, anti-PRO328, anti-PRO331, anti-PRO364, anti-PRO366, anti-PRO535, anti-PRO819, anti-PRO826, anti-PRO1160, anti-PRO1186 or anti-PRO1246 antibody.

9. The article of manufacture of Claim 7, wherein said antagonist is an anti-PRO172, anti-PRO175, anti-PRO178, anti-PRO188, anti-PRO356, anti-PRO179, anti-PRO197, anti-PRO198, anti-PRO182, anti-PRO195, anti-PRO200, anti-PRO211, anti-PRO217, anti-PRO219, anti-PRO221, anti-PRO224, anti-PRO228, anti-PRO245, anti-PRO246, anti-PRO258, anti-PRO261, anti-PRO272, anti-PRO301, anti-PRO322, anti-PRO328, anti-PRO331, anti-PRO364, anti-PRO366, anti-PRO535, anti-PRO819, anti-PRO826, anti-PRO1160, anti-PRO1186 or anti-PRO1246 antibody.

10. The article of manufacture of Claim 7, wherein said composition comprises a therapeutically effective amount of said polypeptide or agonist or antagonist thereof, in admixture with said pharmaceutically acceptable carrier.

11. A method for identifying an agonist of a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide comprising:

- (a) contacting cells and a test compound to be screened under conditions suitable for the induction of a cellular response normally induced by a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide; and

(b) determining the induction of said cellular response to determine if the test compound is an effective agonist, wherein the induction of said cellular response is indicative of said test compound being an effective agonist.

12. The method of Claim 11, wherein the cellular response normally induced by said polypeptide is stimulation of cell proliferation.

13. A method for identifying a compound that inhibits an activity of a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide comprising contacting a test compound with said polypeptide under conditions and for a time sufficient to allow the test compound and polypeptide to interact and determining whether the activity of said polypeptide is inhibited.

14. A method for identifying a compound the inhibits an activity of a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide comprising the steps of:

- (a) contacting cells and a test compound to be screened in the presence of said polypeptide under conditions suitable for the induction of a cellular response normally induced by said polypeptide; and
- (b) determining the induction of said cellular response to determine if the test compound is an effective antagonist.

15. The method of Claim 14, wherein the cellular response normally induced by said polypeptide is stimulation of cell proliferation.

16. A method for identifying a compound that inhibits the expression of a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide in cells that normally expresses the polypeptide, wherein the method comprises contacting the cells with a test compound under conditions suitable for allowing expression of said polypeptide and determining whether the expression of said polypeptide is inhibited.

17. An agonist of a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258,

PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide.

18. An antagonist of a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide.

19. A compound that inhibits the expression of a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide in a mammalian cell which expresses said polypeptide.

20. The compound of Claim 19, wherein said compound is an antisense oligonucleotide.

21. An isolated antibody that binds to a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide.

22. The antibody of Claim 21 which is a monoclonal antibody.

23. The antibody of Claim 21 which is an antibody fragment.

24. The antibody of Claim 21 which is a single-chain antibody.

25. A method for diagnosing a disease or susceptibility to a disease which is related to a mutation in a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide-encoding nucleic acid sequence comprising determining the presence or absence of said mutation in said polypeptide-encoding nucleic acid sequence, wherein the presence or absence of said mutation is indicative of the presence of said disease or susceptibility to said disease.

26. A method of diagnosing a cardiovascular, endothelial or angiogenic disorder in a mammal which comprises analyzing the level of expression of a gene encoding a PRO172, PRO175, PRO178, PRO188, PRO356,



PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide (a) in a test sample of tissue cells obtained from said mammal, and (b) in a control sample of known normal tissue cells of the same cell type, wherein a higher or lower expression level in the test sample as compared to the control sample is indicative of the presence of a cardiovascular, endothelial or angiogenic disorder in said mammal.

27. A method of diagnosing a cardiovascular, endothelial or angiogenic disorder in a mammal which comprises detecting the presence or absence of a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide in a test sample of tissue cells obtained from said mammal, wherein the presence or absence of said polypeptide in said test sample is indicative of the presence of a cardiovascular, endothelial or angiogenic disorder in said mammal.

28. A method of diagnosing a cardiovascular, endothelial or angiogenic disorder in a mammal comprising (a) contacting an anti-PRO172, anti-PRO175, anti-PRO178, anti-PRO188, anti-PRO356, anti-PRO179, anti-PRO197, anti-PRO198, anti-PRO182, anti-PRO195, anti-PRO200, anti-PRO211, anti-PRO217, anti-PRO219, anti-PRO221, anti-PRO224, anti-PRO228, anti-PRO245, anti-PRO246, anti-PRO258, anti-PRO261, anti-PRO272, anti-PRO301, anti-PRO322, anti-PRO328, anti-PRO331, anti-PRO364, anti-PRO366, anti-PRO535, anti-PRO819, anti-PRO826, anti-PRO1160, anti-PRO1186 or anti-PRO1246 antibody with a test sample of tissue cells obtained from the mammal, and (b) detecting the formation of a complex between said antibody and a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide in the test sample, wherein the formation of said complex is indicative of the presence of a cardiovascular, endothelial or angiogenic disorder in the mammal.

29. A method for determining the presence of a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide in a sample comprising contacting a sample suspected of containing said polypeptide with an anti-PRO172, anti-PRO175, anti-PRO178, anti-PRO188, anti-PRO356, anti-PRO179, anti-PRO197, anti-PRO198, anti-PRO182, anti-PRO195, anti-PRO200, anti-PRO211, anti-PRO217, anti-PRO219, anti-PRO221, anti-PRO224, anti-PRO228, anti-PRO245, anti-PRO246, anti-PRO258, anti-PRO261, anti-PRO272, anti-PRO301, anti-PRO322, anti-PRO328, anti-PRO331, anti-PRO364, anti-PRO366, anti-PRO535, anti-PRO819, anti-PRO826, anti-PRO1160, anti-PRO1186 or anti-PRO1246 antibody and

determining binding of said antibody to a component of said sample.

30. A cardiovascular, endothelial or angiogenic disorder diagnostic kit comprising an anti-PRO172, anti-PRO175, anti-PRO178, anti-PRO188, anti-PRO356, anti-PRO179, anti-PRO197, anti-PRO198, anti-PRO182, anti-PRO195, anti-PRO200, anti-PRO211, anti-PRO217, anti-PRO219, anti-PRO221, anti-PRO224, anti-PRO228, anti-PRO245, anti-PRO246, anti-PRO258, anti-PRO261, anti-PRO272, anti-PRO301, anti-PRO322, anti-PRO328, anti-PRO331, anti-PRO364, anti-PRO366, anti-PRO535, anti-PRO819, anti-PRO826, anti-PRO1160, anti-PRO1186 or anti-PRO1246 antibody and a carrier in suitable packaging.

31. A method for treating a cardiovascular, endothelial or angiogenic disorder in a mammal comprising administering to the mammal a therapeutically effective amount of a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide or agonist or antagonist thereof.

32. The method according to Claim 31, wherein the mammal is human.

33. The method of Claim 32, wherein the human has suffered myocardial infarction.

34. The method of Claim 32, wherein the human has cardiac hypertrophy, trauma, a cancer, or age-related macular degeneration.

35. The method of Claim 34, wherein the cardiac hypertrophy is characterized by the presence of an elevated level of  $\text{PGF}_{2\alpha}$ .

36. The method of Claim 31, wherein the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide is administered together with a cardiovascular, endothelial or angiogenic agent.

37. The method of Claim 34, wherein the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide is administered following primary angioplasty.

38. The method of Claim 31, wherein the cardiovascular, endothelial or angiogenic disorder is cancer.

39. The method of Claim 38, wherein the PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide is administered in combination with a chemotherapeutic agent, a growth inhibitory agent or a cytotoxic agent.

40. The method of Claim 31 wherein said agonist is an anti-PRO172, anti-PRO175, anti-PRO178, anti-PRO188, anti-PRO356, anti-PRO179, anti-PRO197, anti-PRO198, anti-PRO182, anti-PRO195, anti-PRO200, anti-PRO211, anti-PRO217, anti-PRO219, anti-PRO221, anti-PRO224, anti-PRO228, anti-PRO245, anti-PRO246, anti-PRO258, anti-PRO261, anti-PRO272, anti-PRO301, anti-PRO322, anti-PRO328, anti-PRO331, anti-PRO364, anti-PRO366, anti-PRO535, anti-PRO819, anti-PRO826, anti-PRO1160, anti-PRO1186 or anti-PRO1246 antibody.

41. The method of Claim 31 wherein said antagonist is an anti-PRO172, anti-PRO175, anti-PRO178, anti-PRO188, anti-PRO356, anti-PRO179, anti-PRO197, anti-PRO198, anti-PRO182, anti-PRO195, anti-PRO200, anti-PRO211, anti-PRO217, anti-PRO219, anti-PRO221, anti-PRO224, anti-PRO228, anti-PRO245, anti-PRO246, anti-PRO258, anti-PRO261, anti-PRO272, anti-PRO301, anti-PRO322, anti-PRO328, anti-PRO331, anti-PRO364, anti-PRO366, anti-PRO535, anti-PRO819, anti-PRO826, anti-PRO1160, anti-PRO1186 or anti-PRO1246 antibody.

42. A method for treating a cardiovascular, endothelial or angiogenic disorder in a mammal comprising administering to the mammal a nucleic acid molecule that encodes a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide or agonist or antagonist thereof.

43. The method of Claim 42 wherein said agonist is an anti-PRO172, anti-PRO175, anti-PRO178, anti-PRO188, anti-PRO356, anti-PRO179, anti-PRO197, anti-PRO198, anti-PRO182, anti-PRO195, anti-PRO200, anti-PRO211, anti-PRO217, anti-PRO219, anti-PRO221, anti-PRO224, anti-PRO228, anti-PRO245, anti-PRO246, anti-PRO258, anti-PRO261, anti-PRO272, anti-PRO301, anti-PRO322, anti-PRO328, anti-PRO331, anti-PRO364, anti-PRO366, anti-PRO535, anti-PRO819, anti-PRO826, anti-PRO1160, anti-PRO1186 or anti-PRO1246 antibody.

44. The method of Claim 42 wherein said antagonist is an anti-PRO172, anti-PRO175, anti-PRO178, anti-PRO188, anti-PRO356, anti-PRO179, anti-PRO197, anti-PRO198, anti-PRO182, anti-PRO195, anti-PRO200, anti-PRO211, anti-PRO217, anti-PRO219, anti-PRO221, anti-PRO224, anti-PRO228, anti-PRO245, anti-PRO246, anti-PRO258, anti-PRO261, anti-PRO272, anti-PRO301, anti-PRO322, anti-PRO328, anti-PRO331, anti-PRO364, anti-

PRO366, anti-PRO535, anti-PRO819, anti-PRO826, anti-PRO1160, anti-PRO1186 or anti-PRO1246 antibody.

45. The method of Claim 42, wherein the mammal is human.
46. The method of Claim 42, wherein the nucleic acid molecule is administered via *ex vivo* gene therapy.
47. A recombinant retroviral particle comprising a retroviral vector consisting essentially of (1) a promoter, (2) nucleic acid encoding a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide or agonist or antagonist thereof, and (3) a signal sequence for cellular secretion of the polypeptide, wherein the retroviral vector is in association with retroviral structural proteins.
48. An *ex vivo* producer cell comprising a nucleic acid construct that expresses retroviral structural proteins and also comprises a retroviral vector consisting essentially of a (1) promoter, (2) nucleic acid encoding a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide or agonist or antagonist thereof, and (3) a signal sequence for cellular secretion of the polypeptide, wherein said producer cell packages the retroviral vector in association with the structural proteins to produce recombinant retroviral particles.
49. A method for inhibiting endothelial cell growth in a mammal comprising administering to the mammal a PRO172, PRO175, PRO178, PRO188, PRO179, PRO197, PRO198, PRO182, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331 or PRO366 polypeptide or agonist thereof, wherein endothelial cell growth in said mammal is inhibited.
50. A method for stimulating endothelial cell growth in a mammal comprising administering to the mammal a PRO356, PRO179, PRO364, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide or agonist thereof, wherein endothelial cell growth in said mammal is stimulated.
51. A method of inhibiting endothelial cell growth in a mammal comprising administering to the mammal an antagonist of a PRO356, PRO179, PRO364, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide, wherein endothelial cell growth in said mammal is inhibited.
52. A method of stimulating endothelial cell growth in a mammal comprising administering to the mammal

an antagonist of a PRO172, PRO175, PRO178, PRO188, PRO179, PRO197, PRO198, PRO182, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331 or PRO366 polypeptide, wherein endothelial cell growth in said mammal is stimulated.

53. A method for inducing cardiac hypertrophy in a mammal comprising administering to the mammal a PRO179, PRO195 or PRO224 polypeptide or agonist thereof, wherein cardiac hypertrophy in said mammal is induced.

54. A method of reducing cardiac hypertrophy in a mammal comprising administering to the mammal an antagonist of a PRO179, PRO195 or PRO224 polypeptide, wherein cardiac hypertrophy in said mammal is reduced.

55. A method for inhibiting angiogenesis induced by a PRO356, PRO179, PRO364, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide in a mammal comprising administering a therapeutically effective amount of an anti-PRO356, anti-PRO179, anti-PRO364, anti-PRO535, anti-PRO819, anti-PRO826, anti-PRO1160, anti-PRO1186 or anti-PRO1246 antibody to the mammal, wherein said angiogenesis is inhibited

56. A method for stimulating angiogenesis induced by a PRO356, PRO179, PRO364, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide in a mammal comprising administering a therapeutically effective amount of said polypeptide to the mammal, wherein said angiogenesis is stimulated.

57. Isolated nucleic acid having at least 80% nucleic acid sequence identity to a nucleotide sequence that encodes an amino acid sequence selected from the group consisting of the amino acid sequence shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:9), Figure 6 (SEQ ID NO:11), Figure 8 (SEQ ID NO:16), Figure 10 (SEQ ID NO:21), Figure 12 (SEQ ID NO:26), Figure 14 (SEQ ID NO:31), Figure 16 (SEQ ID NO:36), Figure 18 (SEQ ID NO:41), Figure 20 (SEQ ID NO:46), Figure 22 (SEQ ID NO:51), Figure 24 (SEQ ID NO:57), Figure 26 (SEQ ID NO:62), Figure 28 (SEQ ID NO:67), Figure 30 (SEQ ID NO:72), Figure 32 (SEQ ID NO:77), Figure 34 (SEQ ID NO:82), Figure 36 (SEQ ID NO:91), Figure 38 (SEQ ID NO:96), Figure 40 (SEQ ID NO:101), Figure 42 (SEQ ID NO:108), Figure 44 (SEQ ID NO:113), Figure 46 (SEQ ID NO:119), Figure 48 (SEQ ID NO:127), Figure 50 (SEQ ID NO:132), Figure 52 (SEQ ID NO:137), Figure 54 (SEQ ID NO:142), Figure 56 (SEQ ID NO:152), Figure 58 (SEQ ID NO:154), Figure 60 (SEQ ID NO:156), Figure 62 (SEQ ID NO:158), Figure 64 (SEQ ID NO:160), Figure 66 (SEQ ID NO:165), and Figure 68 (SEQ ID NO:167).

58. Isolated nucleic acid having at least 80% nucleic acid sequence identity to a nucleotide sequence selected from the group consisting of the nucleotide sequence shown in Figures 1A-1B (SEQ ID NO:1), Figures 3A-3B (SEQ ID NO:8), Figure 5 (SEQ ID NO:10), Figures 7A-7B (SEQ ID NO:15), Figures 9A-9B (SEQ ID NO:20), Figures 11A-11B (SEQ ID NO:25), Figure 13 (SEQ ID NO:30), Figure 15 (SEQ ID NO:35), Figure 17

(SEQ ID NO:40), Figure 19 (SEQ ID NO:45), Figures 21A-21B (SEQ ID NO:50), Figure 23 (SEQ ID NO:56), Figures 25A-25B (SEQ ID NO:61), Figures 27A-27B (SEQ ID NO:66), Figure 29 (SEQ ID NO:71), Figure 31 (SEQ ID NO:76), Figures 33A-33B (SEQ ID NO:81), Figure 35 (SEQ ID NO:90), Figure 37 (SEQ ID NO:95), Figure 39 (SEQ ID NO:100), Figure 41 (SEQ ID NO:107), Figure 43 (SEQ ID NO:112), Figure 45 (SEQ ID NO:118), Figure 47 (SEQ ID NO:126), Figure 49 (SEQ ID NO:131), Figures 51A-51B (SEQ ID NO:136), Figure 53 (SEQ ID NO:141), Figure 55 (SEQ ID NO:151), Figure 57 (SEQ ID NO:153), Figure 59 (SEQ ID NO:155), Figure 61 (SEQ ID NO:157), Figure 63 (SEQ ID NO:159), Figure 65 (SEQ ID NO:164), and Figures 67A-67B (SEQ ID NO:166).

59. Isolated nucleic acid having at least 80% nucleic acid sequence identity to a nucleotide sequence selected from the group consisting of the full-length coding sequence of the nucleotide sequence shown in Figures 1A-1B (SEQ ID NO:1), Figures 3A-3B (SEQ ID NO:8), Figure 5 (SEQ ID NO:10), Figures 7A-7B (SEQ ID NO:15), Figures 9A-9B (SEQ ID NO:20), Figures 11A-11B (SEQ ID NO:25), Figure 13 (SEQ ID NO:30), Figure 15 (SEQ ID NO:35), Figure 17 (SEQ ID NO:40), Figure 19 (SEQ ID NO:45), Figures 21A-21B (SEQ ID NO:50), Figure 23 (SEQ ID NO:56), Figures 25A-25B (SEQ ID NO:61), Figures 27A-27B (SEQ ID NO:66), Figure 29 (SEQ ID NO:71), Figure 31 (SEQ ID NO:76), Figures 33A-33B (SEQ ID NO:81), Figure 35 (SEQ ID NO:90), Figure 37 (SEQ ID NO:95), Figure 39 (SEQ ID NO:100), Figure 41 (SEQ ID NO:107), Figure 43 (SEQ ID NO:112), Figure 45 (SEQ ID NO:118), Figure 47 (SEQ ID NO:126), Figure 49 (SEQ ID NO:131), Figures 51A-51B (SEQ ID NO:136), Figure 53 (SEQ ID NO:141), Figure 55 (SEQ ID NO:151), Figure 57 (SEQ ID NO:153), Figure 59 (SEQ ID NO:155), Figure 61 (SEQ ID NO:157), Figure 63 (SEQ ID NO:159), Figure 65 (SEQ ID NO:164), and Figures 67A-67B (SEQ ID NO:166).

60. Isolated nucleic acid having at least 80% nucleic acid sequence identity to the full-length coding sequence of the DNA deposited under ATCC accession number 209419, 209466, 209282, 209279, 209422, 209281, 209284, 209283, 209296, 209772, 209653, 209258, 209256, 209384, 209262, 209263, 209420, 209265, 209396, 209402, 209391, 209388, 209432, 209669, 209438, 209489, 209436, 209087, 203013, 203006, 203017, 203100, 203091, or 203457.

61. A vector comprising the nucleic acid of any one of Claims 57 to 60.

62. The vector of Claim 61 operably linked to control sequences recognized by a host cell transformed with the vector.

63. A host cell comprising the vector of Claim 61.

64. The host cell of Claim 63, wherein said cell is a CHO cell.

65. The host cell of Claim 63, wherein said cell is an *E. coli*.
66. The host cell of Claim 63, wherein said cell is a yeast cell.
67. The host cell of Claim 63, wherein said cell is a Baculovirus infected insect cell.
68. A process for producing a PRO172, PRO175, PRO178, PRO188, PRO356, PRO179, PRO197, PRO198, PRO182, PRO195, PRO200, PRO211, PRO217, PRO219, PRO221, PRO224, PRO228, PRO245, PRO246, PRO258, PRO261, PRO272, PRO301, PRO322, PRO328, PRO331, PRO364, PRO366, PRO535, PRO819, PRO826, PRO1160, PRO1186 or PRO1246 polypeptide comprising culturing the host cell of Claim 63 under conditions suitable for expression of said polypeptide and recovering said polypeptide from the cell culture.
69. An isolated polypeptide having at least 80% amino acid sequence identity to an amino acid sequence selected from the group consisting of the amino acid sequence shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:9), Figure 6 (SEQ ID NO:11), Figure 8 (SEQ ID NO:16), Figure 10 (SEQ ID NO:21), Figure 12 (SEQ ID NO:26), Figure 14 (SEQ ID NO:31), Figure 16 (SEQ ID NO:36), Figure 18 (SEQ ID NO:41), Figure 20 (SEQ ID NO:46), Figure 22 (SEQ ID NO:51), Figure 24 (SEQ ID NO:57), Figure 26 (SEQ ID NO:62), Figure 28 (SEQ ID NO:67), Figure 30 (SEQ ID NO:72), Figure 32 (SEQ ID NO:77), Figure 34 (SEQ ID NO:82), Figure 36 (SEQ ID NO:91), Figure 38 (SEQ ID NO:96), Figure 40 (SEQ ID NO:101), Figure 42 (SEQ ID NO:108), Figure 44 (SEQ ID NO:113), Figure 46 (SEQ ID NO:119), Figure 48 (SEQ ID NO:127), Figure 50 (SEQ ID NO:132), Figure 52 (SEQ ID NO:137), Figure 54 (SEQ ID NO:142), Figure 56 (SEQ ID NO:152), Figure 58 (SEQ ID NO:154), Figure 60 (SEQ ID NO:156), Figure 62 (SEQ ID NO:158), Figure 64 (SEQ ID NO:160), Figure 66 (SEQ ID NO:165), and Figure 68 (SEQ ID NO:167).
70. An isolated polypeptide scoring at least 80% positives when compared to an amino acid sequence selected from the group consisting of the amino acid sequence shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:9), Figure 6 (SEQ ID NO:11), Figure 8 (SEQ ID NO:16), Figure 10 (SEQ ID NO:21), Figure 12 (SEQ ID NO:26), Figure 14 (SEQ ID NO:31), Figure 16 (SEQ ID NO:36), Figure 18 (SEQ ID NO:41), Figure 20 (SEQ ID NO:46), Figure 22 (SEQ ID NO:51), Figure 24 (SEQ ID NO:57), Figure 26 (SEQ ID NO:62), Figure 28 (SEQ ID NO:67), Figure 30 (SEQ ID NO:72), Figure 32 (SEQ ID NO:77), Figure 34 (SEQ ID NO:82), Figure 36 (SEQ ID NO:91), Figure 38 (SEQ ID NO:96), Figure 40 (SEQ ID NO:101), Figure 42 (SEQ ID NO:108), Figure 44 (SEQ ID NO:113), Figure 46 (SEQ ID NO:119), Figure 48 (SEQ ID NO:127), Figure 50 (SEQ ID NO:132), Figure 52 (SEQ ID NO:137), Figure 54 (SEQ ID NO:142), Figure 56 (SEQ ID NO:152), Figure 58 (SEQ ID NO:154), Figure 60 (SEQ ID NO:156), Figure 62 (SEQ ID NO:158), Figure 64 (SEQ ID NO:160), Figure 66 (SEQ ID NO:165), and Figure 68 (SEQ ID NO:167).
71. An isolated polypeptide having at least 80% amino acid sequence identity to an amino acid sequence

encoded by the full-length coding sequence of the DNA deposited under ATCC accession number 209419, 209466, 209282, 209279, 209422, 209281, 209284, 209283, 209296, 209772, 209653, 209258, 209256, 209384, 209262, 209263, 209420, 209265, 209396, 209402, 209391, 209388, 209432, 209669, 209438, 209489, 209436, 209087, 203013, 203006, 203017, 203100, 203091, or 203457.

72. A chimeric molecule comprising a polypeptide according to any one of Claims 69 to 71 fused to a heterologous amino acid sequence.

73. The chimeric molecule of Claim 72, wherein said heterologous amino acid sequence is an epitope tag sequence.

74. The chimeric molecule of Claim 72, wherein said heterologous amino acid sequence is a Fc region of an immunoglobulin.

75. An antibody which specifically binds to a polypeptide according to any one of Claims 69 to 71.

76. The antibody of Claim 75, wherein said antibody is a monoclonal antibody, a humanized antibody or a single-chain antibody.

77. Isolated nucleic acid having at least 80% nucleic acid sequence identity to:

(a) a nucleotide sequence encoding the polypeptide shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:9), Figure 6 (SEQ ID NO:11), Figure 8 (SEQ ID NO:16), Figure 10 (SEQ ID NO:21), Figure 12 (SEQ ID NO:26), Figure 14 (SEQ ID NO:31), Figure 16 (SEQ ID NO:36), Figure 18 (SEQ ID NO:41), Figure 20 (SEQ ID NO:46), Figure 22 (SEQ ID NO:51), Figure 24 (SEQ ID NO:57), Figure 26 (SEQ ID NO:62), Figure 28 (SEQ ID NO:67), Figure 30 (SEQ ID NO:72), Figure 32 (SEQ ID NO:77), Figure 34 (SEQ ID NO:82), Figure 36 (SEQ ID NO:91), Figure 38 (SEQ ID NO:96), Figure 40 (SEQ ID NO:101), Figure 42 (SEQ ID NO:108), Figure 44 (SEQ ID NO:113), Figure 46 (SEQ ID NO:119), Figure 48 (SEQ ID NO:127), Figure 50 (SEQ ID NO:132), Figure 52 (SEQ ID NO:137), Figure 54 (SEQ ID NO:142), Figure 56 (SEQ ID NO:152), Figure 58 (SEQ ID NO:154), Figure 60 (SEQ ID NO:156), Figure 62 (SEQ ID NO:158), Figure 64 (SEQ ID NO:160), Figure 66 (SEQ ID NO:165), or Figure 68 (SEQ ID NO:167), lacking its associated signal peptide;

(b) a nucleotide sequence encoding an extracellular domain of the polypeptide shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:9), Figure 6 (SEQ ID NO:11), Figure 8 (SEQ ID NO:16), Figure 10 (SEQ ID NO:21), Figure 12 (SEQ ID NO:26), Figure 14 (SEQ ID NO:31), Figure 16 (SEQ ID NO:36), Figure 18 (SEQ ID NO:41), Figure 20 (SEQ ID NO:46), Figure 22 (SEQ ID NO:51), Figure 24 (SEQ ID NO:57), Figure 26 (SEQ ID NO:62), Figure 28 (SEQ ID NO:67), Figure 30 (SEQ ID NO:72), Figure 32 (SEQ ID NO:77), Figure 34 (SEQ ID NO:82), Figure 36 (SEQ ID NO:91), Figure 38 (SEQ ID NO:96), Figure 40 (SEQ ID NO:101), Figure 42 (SEQ ID NO:108), Figure 44 (SEQ ID NO:113), Figure 46 (SEQ ID NO:119), Figure 48 (SEQ ID NO:127), Figure 50



(SEQ ID NO:132), Figure 52 (SEQ ID NO:137), Figure 54 (SEQ ID NO:142), Figure 56 (SEQ ID NO:152), Figure 58 (SEQ ID NO:154), Figure 60 (SEQ ID NO:156), Figure 62 (SEQ ID NO:158), Figure 64 (SEQ ID NO:160), Figure 66 (SEQ ID NO:165), or Figure 68 (SEQ ID NO:167), with its associated signal peptide; or

(c) a nucleotide sequence encoding an extracellular domain of the polypeptide shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:9), Figure 6 (SEQ ID NO:11), Figure 8 (SEQ ID NO:16), Figure 10 (SEQ ID NO:21), Figure 12 (SEQ ID NO:26), Figure 14 (SEQ ID NO:31), Figure 16 (SEQ ID NO:36), Figure 18 (SEQ ID NO:41), Figure 20 (SEQ ID NO:46), Figure 22 (SEQ ID NO:51), Figure 24 (SEQ ID NO:57), Figure 26 (SEQ ID NO:62), Figure 28 (SEQ ID NO:67), Figure 30 (SEQ ID NO:72), Figure 32 (SEQ ID NO:77), Figure 34 (SEQ ID NO:82), Figure 36 (SEQ ID NO:91), Figure 38 (SEQ ID NO:96), Figure 40 (SEQ ID NO:101), Figure 42 (SEQ ID NO:108), Figure 44 (SEQ ID NO:113), Figure 46 (SEQ ID NO:119), Figure 48 (SEQ ID NO:127), Figure 50 (SEQ ID NO:132), Figure 52 (SEQ ID NO:137), Figure 54 (SEQ ID NO:142), Figure 56 (SEQ ID NO:152), Figure 58 (SEQ ID NO:154), Figure 60 (SEQ ID NO:156), Figure 62 (SEQ ID NO:158), Figure 64 (SEQ ID NO:160), Figure 66 (SEQ ID NO:165), or Figure 68 (SEQ ID NO:167), lacking its associated signal peptide.

78. An isolated polypeptide having at least 80% amino acid sequence identity to:

(a) the polypeptide shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:9), Figure 6 (SEQ ID NO:11), Figure 8 (SEQ ID NO:16), Figure 10 (SEQ ID NO:21), Figure 12 (SEQ ID NO:26), Figure 14 (SEQ ID NO:31), Figure 16 (SEQ ID NO:36), Figure 18 (SEQ ID NO:41), Figure 20 (SEQ ID NO:46), Figure 22 (SEQ ID NO:51), Figure 24 (SEQ ID NO:57), Figure 26 (SEQ ID NO:62), Figure 28 (SEQ ID NO:67), Figure 30 (SEQ ID NO:72), Figure 32 (SEQ ID NO:77), Figure 34 (SEQ ID NO:82), Figure 36 (SEQ ID NO:91), Figure 38 (SEQ ID NO:96), Figure 40 (SEQ ID NO:101), Figure 42 (SEQ ID NO:108), Figure 44 (SEQ ID NO:113), Figure 46 (SEQ ID NO:119), Figure 48 (SEQ ID NO:127), Figure 50 (SEQ ID NO:132), Figure 52 (SEQ ID NO:137), Figure 54 (SEQ ID NO:142), Figure 56 (SEQ ID NO:152), Figure 58 (SEQ ID NO:154), Figure 60 (SEQ ID NO:156), Figure 62 (SEQ ID NO:158), Figure 64 (SEQ ID NO:160), Figure 66 (SEQ ID NO:165), or Figure 68 (SEQ ID NO:167), lacking its associated signal peptide;

(b) an extracellular domain of the polypeptide shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:9), Figure 6 (SEQ ID NO:11), Figure 8 (SEQ ID NO:16), Figure 10 (SEQ ID NO:21), Figure 12 (SEQ ID NO:26), Figure 14 (SEQ ID NO:31), Figure 16 (SEQ ID NO:36), Figure 18 (SEQ ID NO:41), Figure 20 (SEQ ID NO:46), Figure 22 (SEQ ID NO:51), Figure 24 (SEQ ID NO:57), Figure 26 (SEQ ID NO:62), Figure 28 (SEQ ID NO:67), Figure 30 (SEQ ID NO:72), Figure 32 (SEQ ID NO:77), Figure 34 (SEQ ID NO:82), Figure 36 (SEQ ID NO:91), Figure 38 (SEQ ID NO:96), Figure 40 (SEQ ID NO:101), Figure 42 (SEQ ID NO:108), Figure 44 (SEQ ID NO:113), Figure 46 (SEQ ID NO:119), Figure 48 (SEQ ID NO:127), Figure 50 (SEQ ID NO:132), Figure 52 (SEQ ID NO:137), Figure 54 (SEQ ID NO:142), Figure 56 (SEQ ID NO:152), Figure 58 (SEQ ID NO:154), Figure 60 (SEQ ID NO:156), Figure 62 (SEQ ID NO:158), Figure 64 (SEQ ID NO:160), Figure 66 (SEQ ID NO:165), or Figure 68 (SEQ ID NO:167), with its associated signal peptide; or

(c) an extracellular domain of the polypeptide shown in Figure 2 (SEQ ID NO:2), Figure 4 (SEQ ID NO:9), Figure 6 (SEQ ID NO:11), Figure 8 (SEQ ID NO:16), Figure 10 (SEQ ID NO:21), Figure 12 (SEQ ID NO:26),

Figure 14 (SEQ ID NO:31), Figure 16 (SEQ ID NO:36), Figure 18 (SEQ ID NO:41), Figure 20 (SEQ ID NO:46), Figure 22 (SEQ ID NO:51), Figure 24 (SEQ ID NO:57), Figure 26 (SEQ ID NO:62), Figure 28 (SEQ ID NO:67), Figure 30 (SEQ ID NO:72), Figure 32 (SEQ ID NO:77), Figure 34 (SEQ ID NO:82), Figure 36 (SEQ ID NO:91), Figure 38 (SEQ ID NO:96), Figure 40 (SEQ ID NO:101), Figure 42 (SEQ ID NO:108), Figure 44 (SEQ ID NO:113), Figure 46 (SEQ ID NO:119), Figure 48 (SEQ ID NO:127), Figure 50 (SEQ ID NO:132), Figure 52 (SEQ ID NO:137), Figure 54 (SEQ ID NO:142), Figure 56 (SEQ ID NO:152), Figure 58 (SEQ ID NO:154), Figure 60 (SEQ ID NO:156), Figure 62 (SEQ ID NO:158), Figure 64 (SEQ ID NO:160), Figure 66 (SEQ ID NO:165), or Figure 68 (SEQ ID NO:167), lacking its associated signal peptide.

FIGURE 1A

TGGGGGCCCCCAGGCTCGCGCGTGGAGCGAAGCAGCATGGGCAGTCGGTGCGCGCTGGCCC  
TGGCGGTGCTCTCGGCCTTGCTGTGTCAAGTCTGGAGCTCTGGGGTGTTTGAAGTGAAGCTG  
CAGGAGTTCGTCAACAAGAAGGGGCTGCTGGGGAACCGCAATTGCTGCCGCGGGGGCGCGGG  
GCCACCGCCGTGCGCCTGCCGGACCTTCTTCCGCGTGTGCCTCAAGCACTACCAGGCCAGCG  
TGTCCCCGAGCCGCCCTGCACCTACGGCAGCGCCGTACCCCCGTGCTGGGCGTCGACTCC  
TTCAGTCTGCCCCGACGGCGGGGGCGCCGACTCCGCGTTTCAAGCAACCCCATCCGCTTCCCCTT  
CGGCTTACCTGGCCGGGCACCTTCTCTCTGATTATTGAAGCTCTCCACACAGATTCTCCTG  
ATGACCTCGCAACAGAAAACCCAGAAAGACTCATCAGCCGCCTGGCCACCCAGAGGCACCTG  
ACGGTGGGCGAGGAGTGGTCCCAGGACCTGCACAGCAGCGGCCGCACGGACCTCAAGTACTC  
CTACCGCTTCGTGTGTGACGAACACTACTACGGAGAGGGCTGCTCCGTTTTCTGCCGTCCCC  
GGGACGATGCCTTCGGCCACTTACCTGTGGGGAGCGTGGGGAGAAAGTGTGAACCCCTGGC  
TGGAAGGGCCCTACTGCACAGAGCCGATCTGCCTGCCTGGATGTGATGAGCAGCATGGATT  
TTGTGACAAACCAGGGGAATGCAAGTGCAGAGTGGGCTGGCAGGGCCGGTACTGTGACGAGT  
GTATCCGCTATCCAGGCTGTCTCCATGGCACCTGCCAGCAGCCCTGGCAGTGCAACTGCCAG  
GAAGGCTGGGGGGGCGCTTTTCTGCAACCAGGACCTGAACTACTGCACACACCATAAGCCCTG  
CAAGAATGGAGCCACCTGCACCAACACGGGCCAGGGGAGCTACACTTGCTCTTGCCGGCCTG  
GGTACACAGGTGCCACCTGCGAGCTGGGGATTGACGAGTGTGACCCAGCCCTTGTAAGAAC  
GGAGGGAGCTGCACGGATCTCGAGAACAGCTACTCCTGTACCTGCCCACCCGGCTTCTACGG  
CAAAATCTGTGAATTGAGTGCCATGACCTGTGCGGACGGCCCTTGCTTTAACGGGGGTGGT  
GCTCAGACAGCCCCGATGGAGGGTACAGCTGCCGCTGCCCCGTGGGCTACTCCGGCTTCAAC  
TGTGAGAAGAAAATTGACTACTGCAGCTCTTACCCTGTTCTAATGGTGCCAAGTGTGTGGA  
CCTCGGTGATGCCTACCTGTGCCGCTGCCAGGCCGGCTTCTCGGGGAGGCACTGTGACGACA  
ACGTGGACGACTGCGCCTCCTCCCCGTGCGCCAACGGGGGCACCTGCCGGGATGGCGTGAAC  
GACTTCTCCTGCACCTGCCCCGCTGGCTACACGGGCAGGAAGTGCAGTGCCCCCGTCAGCAG  
GTGCGAGCACGCACCCTGCCACAATGGGGCCACCTGCCACGAGAGGGGGCCACCGCTATGTGT  
GCGAGTGTGCCCGAGGCTACGGGGGTCCCAACTGCCAGTTCCTGCTCCCCGAGCTGCCCCCG  
GGCCCAGCGGTGGTGGACCTCACTGAGAAGCTAGAGGGCCAGGGCGGGCCATTCCCCTGGGT  
GGCCGTGTGCGCCGGGTCATCCTTGTCTCATGCTGCTGCTGGGCTGTGCCGCTGTGGTGG  
TCTGCGTCCGGCTGAGGCTGCAGAAGCACCGGCCCCCAGCCGACCCCTGCCGGGGGGAGACG  
GAGACCATGAACAACCTGGCCAAGTCCAGCGTGAGAAGGACATCTCAGTCAGCATCATCGG  
GGCCACGCAGATCAAGAACACCAACAAGAAGGCGGACTTCCACGGGGACCACAGCGCCGACA  
AGAATGGCTTCAAGGCCCGCTACCCAGCGGTGGACTATAACCTCGTGCAGGACCTCAAGGGT  
GACGACACCGCCGTCAGGGACGCGCACAGCAAGCGTGACACCAAGTGCCAGCCCCAGGGCTC

FIGURE 1B

CTCAGGGGAGGAGAAGGGGACCCCGACCACACTCAGGGGTGGAGAAGCATCTGAAAGAAAAA  
GGCCGGACTCGGGCTGTTCAACTTCAAAGACACCAAGTACCAGTCGGTGTACGTCATATCC  
GAGGAGAAGGATGAGTGCCTCATAGCAACTGAGGTGTAAAATGGAAGTGAGATGGCAAGACT  
CCCGTTTCTCTTAAATAAGTAAATTCGAAGGATATATGCCCAACGAATGCTGCTGAAGA  
GGAGGGAGGCCTCGTGGACTGCTGCTGAGAAACCGAGTTCAGACCGAGCAGGTTCTCCTCCT  
GAGGTCCTCGACGCCTGCCGACAGCCTGTCGCGGCCCGGCCCTGCGGCACTGCCTTCCGT  
GACGTCGCGCTTGCACTATGGACAGTTGCTCTTAAGAGAATATATATTTAAATGGGTGAACT  
GAATTACGCATAAGAAGCATGCACTGCCTGAGTGTATATTTTGGATTCTTATGAGCCAGTCT  
TTTCTTGAATTAGAAACACAAACACTGCCTTTATTGTCCTTTTTTGATACGAAGATGTGCTTT  
TTCTAGATGGAAAAGATGTGTGTTATTTTTTGGATTGTAAAAATATTTTTCATGATATCTG  
TAAAGCTTGAGTATTTTGTGATGTTTCGTTTTTTATAATTTAAATTTTGGTAAATATGTACAA  
AGGCACTTCGGGTCTATGTGACTATATTTTTTTGTATATAAATGTATTTATGGAATATTGTG  
CAAATGTTATTTGAGTTTTTTACTGTTTTGTAAATGAAGAAATTCCTTTTTTAAATATTTTT  
CCAAAATAAATTTTATGAATGACAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA  
AAAAAAAAAAAAAAAAAAAAA

## FIGURE 2

Signal peptide: Amino acids 1-19

Transmembrane domain: Amino acids 544-565

N-glycosylation site: Amino acids 475-479

cAMP- and cGMP-dependent protein kinase phosphorylation site:  
Amino acids 658-662

Tyrosine kinase phosphorylation sites:  
Amino acids 174-183;250-259

N-myristoylation sites: Amino acids 35-41;38-44;  
96-102;97-103;260-266;  
279-285;280-286;299-305;  
308-314;326-332;338-344;  
376-382;385-391;510-516;  
674-680;681-687;693-699

Aspartic acid and asparagine hydroxylation sites:  
Amino acids 341-353;418-430;  
456-468

Prokaryotic membrane lipoprotein lipid attachment site:  
Amino acids 550-561

EGF-like domain cysteine pattern signatures:  
Amino acids 241-253;272-284;  
312-324;350-362;389-401;  
427-439;465-477;503-515

MGSRCALALAVLSALLCQVWSSGVFELKLQEFVNKKGLLGNRNCCRGGAGPPPCACRTFFRV  
CLKHYQASVSPEPPCTYGS AVTPVLGVDSFSLPDGGGADSAFSNPIRFPFGFTWPGTFSLI  
EALHTDSPDDLATENPERLISR LATQRHLTVGEEWSQDLHSSGRTDLKYSYRFVCDEHYIGE  
GCSVFCRPRDDAFGHFTCGERGEKVCNPGWKGPYCTEPICLPGCDEQHGFC DKPGECKCRVG  
WQGRYCDECIRYPGCLHGTCQQPWQCNCQEGWGGLFCNQDLNYCTHHKPKNGATCTNTGQG  
SYTCSCRPGYTGATCELGIDEC DPSPCKNGGSCTDLENSYSCTCPPGFGYKICELSAMTCAD  
GPCFNNGRCS DSPDGGYSCRCPVGYSGFNCEKKIDYCSSSPCSNGAKCVDLGDAYLCRCQAG  
FSGRHCDNVDDCASSPCANGGTCRDGVNDF SCTCPPGYTGRNCSAPVSRCEHAPCHNGATC  
HERGHRYVCECARGYGGPNCQFLLPELPPGPAVVDL TEKLEGQGGPFPWAVCAGVILVLML  
LLGCAAVVVCVRLRLQKHRPPADPCRGETETMNNLANCQREKDISVSIIGATQIKNTNKKAD  
FHGDHSADKNGFKARYPAVDYNLVQDLKGDDTAVRDAHSKRDTKCQPQGSSGEEKGTPTTLR  
GGEASERKRPDSCGSKDTKYQSVYVISEEKDECVIATEV

FIGURE 3A

CAGCTCTCATTCTCCAAAAATGTGTTTGAGCCACTTGGAATATGCCTTTAAGCCATTCA  
AGAACTCAAGGAGCTCAGAGATCATCCTGGAAGCTGTGGCTCTTTTGCTCAATAGTTATGTT  
GCTATTTCTTTGCTCCTTCAGTTGGCTAATCTTTATTTTTCTCCAATTAGAGACTGCTAAGG  
AGCCCTGTATGGCTAAGTTTGGACCATTACCCTCAAATGGCAAATGGCATCTTCTGAACCT  
CCTTGCGTGAATAAGGTGTCTGACTGGAAGCTGGAGATACTTCAGAATGGCTTATATTTAAT  
TTATGGCCAAGTGGCTCCCAATGCAAACTACAATGATGTAGCTCCTTTTGAGGTGCGGCTGT  
ATAAAAACAAAGACATGATACAACTCTAACAAACAAATCTAAAATCCAAAATGTAGGAGGG  
ACTTATGAATTGCATGTTGGGGACACCATAGACTTGATATTCAACTCTGAGCATCAGGTTCT  
AAAAATAATACATACTGGGGTATCATTTTACTAGCAAATCCCCAATTCATCTCCTAGAGAC  
TTGATTTGATCTCCTCATTCCTTCAGCACATGTAGAGGTGCCAGTGGGTGGATTGGAGGGA  
GAAGATATTCAATTTCTAGAGTTTGTCTGTCTACAAAAATCAACACAAACAGAACTCCTCTG  
CACGTGAATTTTCATCTATCATGCCTATCTGAAAGAGACTCAGGGGAAGAGCCAAAGACTTT  
TGTTTGGATCTGCAGAAATACTTCATTAATCCATGATAAAACAAATATGGATGACAGAGGAC  
ATGTGCTTTTCAAAGAATCTTTATCTAATTCTTGAATTCATGAGTGGAAAAATGGAGTTCTA  
TTCCCATGGAAGATTTACCTGGTATGCAAAAAGGATCTGGGGCAGTAGCCTGGCTTTGTTCT  
CATATTCTTGGGCTGCTGTAATTCATTCTTCTCATACTCCCATCTTCTGAGACCCTCCCAAT  
AAAAAGTAGACTGATAGGATGGCCACAGATATGCCTACCATAACCCTACTTTAGATATGGTGG  
TGTTAGAAGATAAAGAACAATCTGAGAACTATTGGAATAGAGGTACAAGTGGCATAAAATGG  
AATGTACGCTATCTGGAAATTTCTCTTGGTTTTATCTTCCTCAGGATGCAGGGTGCTTTAAA  
AAGCCTTATCAAAGGAGTCATTCCGAACCCTCACGTAGAGCTTTGTGAGACCTTACTGTTGG  
TGTGTGTGTCTAAACATTGCTAATTGTAAAGAAAGAGTAACCATTAGTAATCATTAGGTTTA  
ACCCAGAAATGGTATTATCATTACTGGATTATGTATGTAATGATTTAGTATTTTATAGCTAG  
CTTTCCACAGTTTGCAAAGTGCTTTTCGTAAACAGTTAGCAATTCTATGAAGTTAATTGGGC  
AGGCATTTGGGGGAAAATTTTAGTGATGAGAATGTGATAGCATAGCATAGCCAACCTTTCCTC  
AACTCATAGGACAAGTGAACAAGAGGCAATGGGTAGTCCCCTGCATTGCACTGTCTCAGC  
TTTAGAATTGTTATTTCTGCTATCGTGTTATAAGACTCTAAACTTAGCGAATTCACCTTTTC  
AGGAAGCATATTCCCCTTTAGCCCAAGGTGAGCAGAGTGAAGCTACAACAGATCTTTCCTTT  
ACCAGCACACTTTTTTTTTTTTTTCTGCTGAATCAGGGAGATCCAGGATGCTGTTTCAGGC  
CAAATCCCAACCAAATTTCCCCTTTTCACTTTGCAGGGCCCATCTTAGTCAAATGTGCTAACT  
TCTAAAATAATAATAGCACTAATTCAAAATTTTGAATCTTAAATTAGCTACTTGCNGGT  
TGCTTGTTGAAAGGNATATAATGATTACATTGTAAACAAATTTAAATATTTATGGATATT

FIGURE 3B

GTGAAAAGCTGCATTATGTTAAATAATATTACATGTAAAGCT

## FIGURE 4

Signal peptide:	Amino acids 1-25
Transmembrane Domain:	Amino acids 26-51
Extracellular Domain:	Amino acids 52-177
N-linked glycosylation sites:	Amino acids 129-133
	Amino acids 161-165
N-myristoylation site:	Amino acids 18-24

MCLSHLENMPLSHSRTQGAQRSSWKLWLFCSIVMLLFLCSFSWLIFIFLQLETAKEPCMAKF  
GPLPSKWQMASSEPPCVNKVSDWKLEILQNGLYLIYGQVAPNANYNDVAPFEVRLYKNKDMI  
QTLTNKSKIQNVGGTYELHVGDTIDLIFNSEHQVLKNNTYWGIIILLANPQFIS



FIGURE 5

GGCTCAGAGGCCCCACTGGACCCTCGGCTCTTCCTTGGACTTCTTGTGTGTTCTGTGAGCTT  
CGCTGGATTTCAGGGTCTTGGGCATCAGAGGTGAGAGGGTGGGAAGGTCCGCCGCGATGGGGA  
AGCCCTGGCTGCGTGCGCTACAGCTGCTGCTCCTGCTGGGCGCGTTCGTGGGCGCGGGCGGGC  
GCCCCGCGCTGCACCTACACCTTCGTGCTGCCCCCGCAGAAGTTCACGGGCGCTGTGTGCTG  
GAGCGGCCCCCGCATCCACGCGGGCGACGCCCCGAGGCCGCCAACGCCAGCGAGCTGGCGGGCGC  
TGCGCATGCGCGTTCGGCCGCCACGAGGAGCTGTTACGCGAGCTGCAGAGGCTGGCGGGCGGCC  
GACGGCGCCGTGGCCGGCGAGGTGCGCGCGCTGCGCAAGGAGAGCCGCGGCCTGAGCGCGCG  
CCTGGGCCAGTTGCGCGCGCAGCTGCAGCACGAGGCGGGGCCCGGGCGGGCCCCGGGGCGG  
ATCTGGGGGCGGAGCCTGCCGCGGCGCTGGCGCTGCTCGGGGAGCGCGTGCTCAACGCGTCC  
GCCGAGGCTCAGCGCGCAGCCGCCCGGTTCCACCAGCTGGACGTCAAGTTCGCGAGCTGGC  
GCAGCTCGTCACCCAGCAGAGCAGTCTCATCGCCCGCTGGAGCGCCTGTGCCCGGGAGGCG  
CGGGCGGGCAGCAGCAGGTCTGCCGCCACCCCCACTGGTGCCTGTGGTTCCGGTCCGTCTT  
GTGGGTAGCACCAAGTGACACCAAGTAGGATGCTGGACCCAGCCCCAGAGCCCCAGAGAGACCA  
GACCCAGAGACAGCAGGAGCCCATGGCTTCTCCCATGCCTGCAGGTACCCCTGCGGTCCCCA  
CCAAGCCTGTGGGCCCCGTGGCAGGATTGTGCAGAGGCCCGCCAGGCAGGCCATGAACAGAGT  
GGAGTGTATGAACTGCGAGTGGGCCGTACGTAGTGTGAGTATGGTGTGAGCAGCAACTGGA  
GGGTGGAGGCTGGACTGTGATCCAGCGGAGGCAAGATGGTTCAGTCAACTTCTTCACTACCT  
GGCAGCACTATAAGGCGGGCTTTGGGCGGCCAGACGGAGAATACTGGCTGGGCCTTGAACCC  
GTGTATCAGCTGACCAGCCGTGGGGACCATGAGCTGCTGGTTCTCCTGGAGGACTGGGGGGG  
CCGTGGAGCACGTGCCCACTATGATGGCTTCTCCCTGGAACCCGAGAGCGACCACTACCGCC  
TGCGGCTTGGCCAGTACCATGGTGATGCTGGAGACTCTCTTTCCTGGCACAATGACAAGCCC  
TTCAGCACCGTGGATAGGGACCGAGACTCCTATTCTGGTAACTGTGCCCTGTACCAGCGGGG  
AGGCTGGTGGTACCATGCCTGTGCCCACTCCAACCTCAACGGTGTGTGGCACCACGGCGGGC  
ACTACCGAAGCCGCTACCAGGATGGTGTCTACTGGGCTGAGTTTCGTGGTGGGGCATATTCT  
CTCAGGAAGGCCCGCATGCTCATTCGGCCCCCTGAAGCTGTGA<sup>1</sup>CTCTGTGTTCTCTGTCCCC  
TAGGCCCTAGAGGACATTGGTCAGCAGGAGCCCAAGTTGTTCTGGCCACACCTTCTTTGTGG  
CTCAGTGCCAATGTGTCCACAGAACTTCCCACTGTGGATCTGTGACCCCTGGGCGCTGAAAA  
TGGGACCCAGGAATCCCCCCCCGTCAATATCTTGGCCTCAGATGGCTCCCCAAGGTCATTAT  
ATCTCGGTTTGGAGCTCATATCTTATAATAACACAAAGTAGCCAC

## FIGURE 6

Signal peptide:	Amino acids 1-20
N-glycosylation sites:	Amino acids 58-62;145-149
cAMP- and cGMP-dependent protein kinase phosphorylation site:	Amino acids 97-101
Tyrosine kinase phosphorylation site:	
	Amino acids 441-448
N-myristoylation sites:	Amino acids 16-22;23-29;
	87-93;108-114;121-127;
	125-131;129-135;187-193;
	293-299;353-359;378-384;
	445-451;453-459
Cell attachment sequence:	Amino acids 340-343
Fibrinogen beta and gamma chains C-terminal domain signature:	
	Amino acids 418-431

MGKPWLRALQLLLLLLGASWARAGAPRCTYTFVLPPQKFTGAVCWSGPASTRATPEAANASEL  
 AALRMRVGRHEELLRELQRLAAADGAVAGEVRALRKESRGLSARLGQLRAQLQHEAGPGAGP  
 GADLGAEPAAALALLGERVLNASAEARAAARFHQLDVKFRELAQLVTQQSSLIARLERLCP  
 GGAGGQQQVLPPPPLVPVVPVRLVGSTSDTSRMLDPAPEPQRDQTQRQQEPMASPMAGHPA  
 VPTKPVGPWQDCAEARQAGHEQSGVYELRVGRHVSVWCEQQLEGGGWTVIQRRQDGSVNFF  
 TTWQHYKAGFRPDGEYWLGLEPVYQLTSRGDHELLVLLEDWGGRGARAHYDGFSLPEPSDH  
 YRLRLGQYHGDAGDSLWHNDKPFSTVDRDRDSYSGNCALYQRGGWWYHACAHSNLNGVWHH  
 GGHYRSRYQDGVYWAEFRGGAYSLRKAAMLIRPLKL

FIGURE 7A

GCAGCTGGTTACTGCATTTCTCCATGTGGCAGACAGAGCAAAGCCACAACGCTTTCTCTGCT  
GGATTAAAGACGGCCACAGACCAGAACTTCCACTATACTACTTAAAATTACATAGGTGGCT  
TGTCAAATTCAATTGATTAGTATTGTAAAAGGAAAAAGAAGTTCCTTCTTACAGCTTGGATT  
CAACGGTCCAAAACAAAATGCAGCTGCCATTAAAGTCTCAGATGAACAAACTTCTACACTG  
ATTTTAAAATCAAGAATAAGGGCAGCAAGTTTCTGGATTCACTGAATCAACAGACACAAAA  
AGCTGGCAATATAGCAACTATGAAGAGAAAAGCTACTAATAAAATTAACCCAACGCATAGAA  
GACTTTTTTTTTCTTCTAAAAACAATAAGTAAAGACTTAAATTTAAACACATCATTTTAC  
AACCTCATTTCAAAATGAAGACTTTTACCTGGACCCTAGGTGTGCTATTCTTCCTACTAGTG  
GACACTGGACATTGCAGAGGTGGACAATTCAAATTAATAAAATAAACCCAGAGAAGATACCC  
TCGTGCCACAGATGGTAAAGAGGAAGCAAAGAAATGTGCATACACATTCCTGGTACCTGAAC  
AAAGAATAACAGGGCCAATCTGTGTCAACACCAAGGGGCAAGATGCAAGTACCATTAAAGAC  
ATGATCACCAGGATGGACCTTGAAAACCTGAAGGATGTGCTCTCCAGGCAGAAGCGGGAGAT  
AGATGTTCTGCAACTGGTGGTGGATGTAGATGGAAACATTGTGAATGAGGTAAAGCTGCTGA  
GAAAGGAAAGCCGTAACATGAACTCTCGTGTTACTCAACTCTATATGCAATTATTACATGAG  
ATTATCCGTAAGAGGGATAATTCACCTGAACTTTCCCAACTGGAAAACAAAATCCTCAATGT  
CACCACAGAAATGTTGAAGATGGCAACAAGATACAGGGAACTAGAGGTGAAATACGCTTCCT  
TGACTGATCTTGTCAATAACCAATCTGTGATGATCACTTTGTTGGAAGAACAGTGCTTGAGG  
ATATTTTCCCGACAAGACACCCATGTGTCTCCCCACTTGTCCAGGTGGTGCCACAACATAT  
TCCTAACAGCCAACAGTATACTCCTGGTCTGCTGGGAGGTAACGAGATTAGAGGGGATCCAG  
GTTATCCCAGAGATTTAATGCCACCACCTGATCTGGCAACTTCTCCACCAAAGCCCTTTC  
AAGATAACCACCGGTAACCTTTTCATCAATGAAGGACCATTCAAAGACTGTCAGCAAGCAAAAGA  
AGCTGGGCATTCCGGTCAGTGGGATTTATATGATTAAACCTGAAAACAGCAATGGACCAATGC  
AGTTATGGTGTGAAAACAGTTTGGACCCTGGGGGTTGGACTGTTATTAGAAAAGAACAGAC  
GGCTCTGTCAACTTCTTCAGAAATTTGGGAAAATTATAAGAAAGGGTTTGGAAACATTGACGG  
AGAATACTGGCTTGACTGGAAAATATCTATATGCTTAGCAATCAAGATAATTACAAGTTAT  
TGATTGAATTAGAAGACTGGAGTGATAAAAAAGTCTATGCAGAATACAGCAGCTTTCGTCTG  
GAACCTGAAAGTGAATTCTATAGACTGCGCCTGGGAACTTACCAGGGAAATGCAGGGGATTC  
TATGATGTGGCATAATGGTAAACAATTCACCACACTGGACAGAGATAAAGATATGTATGCAG  
GAACTGCGCCCACTTTCATAAAGGAGGCTGGTGGTACAATGCCTGTGCACATTCTAACCTA  
AATGGAGTATGGTACAGAGGAGGCCATTACAGAAGCAAGCACCAAGATGGAATTTTCTGGGC  
CGAATACAGAGGCGGGTCATACTCCTTAAGAGCAGTTCAGATGATGATCAAGCCTATTGACT  
GAAGAGAGACACTCGCCAATTTAAATGACACAGAACTTTGTACTTTTCAGCTCTTAAAAATG

FIGURE 7B

TAAATGTTACATGTATATTACTTGGCACAATTTATTTCTACACAGAAAGTTTTTAAAATGAA  
TTTTACCGTAACTATAAAAGGGAACCTATAAATGTAGTTTCATCTGTCTCAATTACTGCAG  
AAAATTATGTGTATCCACAACCTAGTTATTTTAAAAATTATGTTGACTAAATACAAAGTTTG  
TTTTCTAAAATGTAAATATTTGCCACAATGTAAAGCAAATCTTAGCTATATTTTAAATCATA  
AATAACATGTTCAAGATACTTAACAATTTATTTAAAATCTAAGATTGCTCTAACGTCTAGTG  
AAAAAATATTTTTTAAATTTTCAGCCAAATAATGCATTTTATTTTATAAAAATACAGACAGA  
AAATTAGGGAGAACTTCTAGTTTTTGCCAATAGAAAATGTTCTTCCATTGAATAAAAGTTAT  
TTCAAATTGAATTTGTGCCTTTCACACGTAATGATTAAATCTGAATTCTTAATAATATATCC  
TATGCTGATTTTCCCAAACATGACCCATAGTATTAAATACATATCATTTTTTAAAAATAAAA  
AAAAACCCAAAAATAATGCATGCATAATTTAAATGGTCAATTTATAAAGACAAATCTATGAA  
TGAATTTTTTCAGTGTTATCTTCATATGATATGCTGAACACCAAATCTCCAGAAATGCATTT  
TATGTAGTTCTAAAATCAGCAAAATATTGGTATTACAAAAATGCAGAATATTTAGTGTGCTA  
CAGATCTGAATTATAGTTCTAATTTATTATTACTTTTTTTCTAATTTACTGATCTTACTACT  
ACAAAGAAAAAAAAACCCAACCCATCTGCAATTCAAATCAGAAAGTTTGGACAGCTTTACAA  
GTATTAGTGCATGCTCAGAACAGGTGGGACTAAAACAACTCAAGGAACTGTTGGCTGTTTT  
CCCGATACTGAGAATTCAACAGCTCCAGAGCAGAAGCCACAGGGGCATAGCTTAGTCCAAAC  
TGCTAATTTCAATTTACAGTGTATGTAACGCTTAGTCTCACAGTGTCTTTAACTCATCTTTG  
CAATCAACAACTTTACTAGTGACTTTCTGGAACAATTTCTTTTCAGGAATACATATTCCTG  
CTTAGAGGTGACCTTGCCTTAATATATTTGTGAAGTTAAAATTTTAAAGATAGCTCATGAAA  
CTTTTGCTTAAGCAAAAAGAAAACCTCGAATTGAAATGTGTGAGGCAAACTATGCATGGGAA  
TAGCTTAATGTGAAGATAATCATTTGGACAACCTCAAATCCATCAACATGACCAATGTTTTTC  
ATCTGCCACATCTCAAAATAAACTTCTGGTGAAACAAATTAACAAAAATATCCAAACCTCA  
AAAAAA

## FIGURE 8

Signal peptide: Amino acids 1-23  
N-glycosylation sites: Amino acids 160-164;188-192  
cAMP- and cGMP-dependent protein kinase phosphorylation site:  
Amino acids 120-124  
Tyrosine kinase phosphorylation sites:  
Amino acids 173-180;387-396  
N-myristoylation sites: Amino acids 70-76;110-116;  
232-238;343-349;400-406;  
467-473;475-481  
Fibrinogen beta and gamma chains C-terminal domain signature:  
Amino acids 440-453

MKTFTWTLGVLFLLVDTGHCRGGQFKIKKINQRRYPRATDGKEEAKKCAYTFLVPEQRITG  
PICVNTKGQDASTIKDMI TRMDLENLKDVL SRQKREIDVLQLVVDVDGNIVNEVKLLRKESR  
NMNSRVTQLYMQLLHEIIRKRDNSLELSQLENKILNVTTEMLKMATRYRELEVKYASLTDLV  
NNQSVMITLLEEQLRIFSRQDTHVSPPLVQVVPQHIPNSQQYTPGLLGGNEIQRDPGYPRD  
LMPPPDLATSPKSPFKIPPVTFINEGPFKDCQQAKEAGHSVSGIYMIKPENSNGPMQLWCE  
NSLDPGGWTVIQKRTDGSVNFFRNWENYKKGFGNIDGEYWLGLENIYMLSNQDNYKLLIELE  
DWSDDKKVYAEYSSFRLEPESEFYRLRLGTYQGNAGDSMMWHNGKQFTTLDRDKDMYAGNCAH  
FHKGGWWYNACAHSNLNGVWYRGGHYRSKHQDGI FWA EYRGGSYSLRAVQMMIKPID

FIGURE 9A

GAAAGCTATAGGCTACCCATTTCAGCTCCCCTGTCAGAGACTCAAGCTTTGAGAAAGGCTAGC  
AAAGAGCAAGGAAAGAGAGAAAAACAACAAAGTGGCGAGGCCCTCAGAGTGAAAGCGTAAGGT  
TCAGTCAGCCTGCTGCAGCTTTGCAGACCTCAGCTGGGCATCTCCAGACTCCCCTGAAGGAA  
GAGCCTTCCTCACCCAAACCCACAAAAGATGCTGAAAAAGCCTCTCTCAGCTGTGACCTGGC  
TCTGCATTTTTCATCGTGGCCTTTGTCAGCCACCCAGCGTGGCTGCAGAAGCTCTCTAAGCAC  
AAGACACCAGCACAGCCACAGCTCAAAGCGGCCAACTGCTGTGAGGAGGTGAAGGAGCTCAA  
GGCCCAAGTTGCCAACCTTAGCAGCCTGCTGAGTGAAGTGAACAAGAAGCAGGAGAGGGACT  
GGGTGAGCGTGGTCATGCAGGTGATGGAGCTGGAGAGCAACAGCAAGCGCATGGAGTCGCGG  
CTCACAGATGCTGAGAGCAAGTACTCCGAGATGAACAACCAAATTGACATCATGCAGCTGCA  
GGCAGCACAGACGGTCACTCAGACCTCCGCAGATGCCATCTACGACTGCTCTTCCCTCTACC  
AGAAGAACTACCGCATCTCTGGAGTGTATAAGCTTCCTCCTGATGACTTCCTGGGCAGCCCT  
GAACTGGAGGTGTTCTGTGACATGGAGACTTCAGGCGGAGGCTGGACCATCATCCAGAGACG  
AAAAAGTGGCCTTGTCTCCTTCTACCGGGACTGGAAGCAGTACAAGCAGGGCTTTGGCAGCA  
TCCGTGGGGACTTCTGGCTGGGGAACGAACACATCCACCGGCTCTCCAGACAGCCAACCCGG  
CTGCGTGTAGAGATGGAGGACTGGGAGGGCAACCTGCGCTACGCTGAGTATAGCCACTTTGT  
TTTGGGCAATGAACTCAACAGCTATCGCCTCTTCCTGGGGAACTACACTGGCAATGTGGGGA  
ACGACGCCCTCCAGTATCATAACAACACAGCCTTCAGCACCAAGGACAAGGACAATGACAAC  
TGCTTGGACAAGTGTGCACAGCTCCGCAAAGGTGGCTACTGGTACAAGTGTGCACAGACTC  
CAACCTCAATGGAGTGTACTACCGCCTGGGTGAGCACAATAAGCACCTGGATGGCATCACCT  
GGTATGGCTGGCATGGATCTACCTACTCCCTCAAACGGGTGGAGATGAAAATCCGCCCAGAA  
GACTTCAAGCCTTTAAAAGGAGGCTGCCGTGGAGCACGGATACAGAACTGAGACACGTGGAG  
ACTGGATGAGGGCAGATGAGGACAGGAAGAGAGTGTTAGAAAGGGTAGGACTGAGAAACAGC  
CTATAATCTCCAAAGAAAGAATAAGTCTCCAAGGAGCACAAAAAATCATATGTACCAAGGA  
TGTTACAGTAAACAGGATGAACTATTTAAACCCACTGGGTCTGCCACATCCTTCTCAAGGT  
GGTAGACTGAGTGGGGTCTCTCTGCCCCAAGATCCCTGACATAGCAGTAGCTTGTCTTTTCCA  
CATGATTTGTCTGTGAAAGAAAATAATTTTGAGATCGTTTTATCTATTTTCTCTACGGCTTA  
GGCTATGTGAGGGCAAAACACAAATCCCTTTGCTAAAAAGAACCATATTATTTTGATTCTCA  
AAGGATAGGCCTTTGAGTGTAGAGAAAGGAGTGAAGGAGGCAGGTGGGAAATGGTATTTCT  
ATTTTTAAATCCAGTGAAATTATCTTGAGTCTACACATTATTTTTAAACACAAAAATTGTT  
CGGCTGGAAGTGAACCCAGGCTGGACTTGCGGGGAGGAACTCCAGGGCACTGCATCTGGCGA  
TCAGACTCTGAGCACTGCCCTGCTCGCCTTGGTCATGTACAGCACTGAAAGGAATGAAGCA

## FIGURE 9B

CCAGCAGGAGGTGGACAGAGTCTCTCATGGATGCCGGCACAAACTGCCTTAAATATTCAT  
AGTTAATACAGGTATATCTATTTTTATTTACTTTGTAAGAAACAAGCTCAAGGAGCTTCCTT  
TTAAATTTTGTCTGTAGGAAATGGTTGAAAACCTGAAGGTAGATGGTGTTATAGTTAATAATA  
AATGCTGTAAATAAGCATCTCACTTTGTAAAAATAAAATATTGTGGTTTTGTTTTAAACATT  
CAACGTTTCTTTTCCTTCTACAATAAACACTTTCAAAATGTG

## FIGURE 10

Signal peptide: Amino acids 1-26  
N-glycosylation sites: Amino acids 58-62;253-257;  
267-271  
Glycosaminoglycan attachment site: Amino acids 167-171  
cAMP- and cGMP-dependent protein kinase phosphorylation site:  
Amino acids 176-180  
N-myristoylation sites: Amino acids 168-174;196-202;  
241-247;252-258;256-262;  
327-333  
Cell attachment sequence: Amino acids 199-202

MLKKPLSAVTWLCIFIVAFVSHPAWLQKLSKHKTPAQPQLKAANCCEEVKELKAQVANLSSL  
LSELNKKQERDWVSVVMQVMELESNSKRME SRLTDAESKYSEMNNQIDIMQLQAAQTVTQTS  
ADAIYDCSSLYQKNYRISGVYKLPPDDFLGSPELEVFCDMETSGGGWTIIQRRKSGLVSFYR  
DWKQYKQGFGSIRGDFWLGNEHIHRLSRQPTRLRVEMEDWEGNLRVYAEYSHFVLGNELNSYR  
LFLGNYTGNVGN DALQYHNNTAFSTKDKDNDNCLDKCAQLRKGGYWYNCCTDSNLNGVYYRL  
GEHNKHLDGITWYGWHGSTYSLKR VEMKIRPEDFKP



FIGURE 11A

GCGGACGCGTGGGTGAAATTGAAAATCAAGATAAAAATGTTTCACAATTAAGCTCCTTCTTTT  
TATTGTTCTCTAGTTATTTCTCCAGAATTGATCAAGACAATTCATCATTTGATTCTCTAT  
CTCCAGAGCCAAAATCAAGATTTGCTATGTTAGACGATGTAAAAATTTTAGCCAATGGCCTC  
CTTCAGTTGGGACATGGTCTTAAAGACTTTGTCCATAAGACGAAGGGCCAAATTAATGACAT  
ATTTCAAAAATCAACATATTTGATCAGTCTTTTATGATCTATCGCTGCAAACCAGTGAAA  
TCAAAGAAGAAGAAAAGGAACTGAGAAGAACTACATATAAACTACAAGTCAAAAATGAAGAG  
GTAAAGAATATGTCACTTGAACCTCAACTCAAACTTGAAAGCCTCCTAGAAGAAAAAATTCT  
ACTTCAACAAAAAGTGAAATATTTAGAAGAGCAACTAACTAACTTAATTCAAAATCAACCTG  
AACTCCAGAACACCCAGAAGTAACTTCACTTAAAACTTTTGTAGAAAAACAAGATAATAGC  
ATCAAAGACCTTCTCCAGACCGTGGAAGACCAATATAACAATTAAACCAACAGCATAGTCA  
AATAAAAGAAATAGAAAATCAGCTCAGAAGGACTAGTATTCAAGAACCCACAGAAATTTCTC  
TATCTTCCAAGCCAAGAGCACCAAGAACTACTCCCTTTCTTCAGTTGAATGAAATAAGAAAT  
GTAAACATGATGGCATTCTTGCTGAATGTACCACCATTTATAACAGAGGTGAACATACAAG  
TGGCATGTATGCCATCAGACCCAGCAACTCTCAAGTTTTTCATGTCTACTGTGATGTTATAT  
CAGGTAGTCCATGGACATTAATTCAACATCGAATAGATGGATCACAAAATTCATGAAACG  
TGGGAGAACTACAAATATGGTTTTTGGGAGGCTTGATGGAGAATTTTGGTTGGGCCTAGAGAA  
GATATACTCCATAGTGAAGCAATCTAATTATGTTTTACGAATTGAGTTGGAAGACTGGAAAGAC  
AACAAACATTATATTGAATATTTCTTTTTACTTGGGAAATCACGAAACCAACTATACGCTACA  
TCTAGTTGCGATTACTGGCAATGTCCCCAATGCAATCCCGGAAAACAAAGATTTGGTGTTTT  
CTACTTGGGATCACAAAGCAAAAGGACACTTCAACTGTCCAGAGGGTTATTTCAGGAGGCTGG  
TGGTGGCATGATGAGTGTGGAGAAAACAACCTAAATGGTAAATATAACAAACCAAGAGCAAA  
ATCTAAGCCAGAGAGGAGAAGAGGATTATCTTGGAAGTCTCAAAATGGAAGGTTATACTCTA  
TAAAATCAACCAAAATGTTGATCCATCCAACAGATTCAGAAAGCTTTGAATGAACTGAGGCA  
ATTTAAAGGCATATTTAACCATTAACTCATTCCAAGTTAATGTGGTCTAATAATCTGGTATA  
AATCCTTAAGAGAAAAGCTTGAGAAATAGATTTTTTTTATCTTAAAGTCACTGTCTATTTAAG  
ATTAAACATACAATCACATAACCTTAAAGAATACCGTTTACATTTCTCAATCAAAATTTCTTA  
TAATACTATTTGTTTTAAATTTTGTGATGTGGGAATCAATTTTAGATGGTCACAATCTAGAT  
TATAATCAATAGGTGAACTTATTAATAAATTTTCTAAATAAAAAATTTAGAGACTTTTATT  
TAAAAGGCATCATATGAGCTAATATCAAACTTTCCAGTTTAAAAAACTAGTACTCTTGT  
TAAAATCTAACTTGACTAAATACAGAGGACTGGTAATTGTACAGTTCTTAAATGTTGTAG  
TATTAATTTCAAACTAAAAATCGTCAGCACAGAGTATGTGTAAAAATCTGTAATACAAATT

FIGURE 11B

TTTAAACTGATGCTTCATTTTGCTACAAAATAATTTGGAGTAAATGTTTGATATGATTTATT  
TATGAAACCTAATGAAGCAGAATTAAATACTGTATTAAAATAAGTTCGCTGTCTTT

## FIGURE 12

Signal peptide: Amino acids 1-16  
N-glycosylation sites: Amino acids 23-27;115-119;  
296-300;357-361  
cAMP- and cGMP-dependent protein kinase phosphorylation sites:  
Amino acids 100-104;204-208  
Tyrosine kinase phosphorylation site:  
Amino acids 342-348  
N-myristoylation sites: Amino acids 279-285;352-358;  
367-373  
Leucine zipper patterns: Amino acids 120-142;127-149

MFTIKLLLFIVPLVISSRIDQDNSSFDSLSPKSRFAMLDDVKILANGLLQLGHGLKDFVH  
KTKGQINDIFQKLNIFDQSFYDLSLQTSEIKEEEKELRRTTYKLQVKNEEVKNMSLELNSKL  
ESLLEEKILLQQKVYLEEQLTNLIQNQPETPEHPEVTSKTFVEKQDNSIKDLLQTVEDQY  
KQLNQQHSQIKEIENQLRRTSIQEPTEISLSSKPRAPRTTPFLQLNEIRNVKHDGIPAECTT  
IYNRGEHTSGMYAIRPSNSQVFHVYCDVISGSPWTLIQHRIDGSQNFNETWENYKYGFGRLD  
GEFWLGLEKIYSIVKQSNYVLRIELEDWKDNKHYYEYSFYLGNHETNYTLHLVAITGNVPNA  
IPENKDLVFSTWDHKAKGHFNCPEGYSGGWWHDECGENNLNGKYNKPRAKSKPERRRGLSW  
KSQNGRLYSIKSTKMLIHPTDSESFE

FIGURE 13

GCCGAGCTGAGCGGATCCTCACATGACTGTGATCCGATTCTTTCCAGCGGCTTCTGCAACCA  
AGCGGGTCTTACCCCCGGTCCTCCGCGTCTCCAGTCCTCGCACCTGGAACCCCAACGTCCCC  
GAGAGTCCCCGAATCCCCGCTCCCAGGCTACCTAAGAGGATGAGCGGTGCTCCGACGGCCGG  
GGCAGCCCTGATGCTCTGCGCCGCCACCGCCGTGCTACTGAGCGCTCAGGGCGGACCCGTGC  
AGTCCAAGTCGCCGCGCTTTGCGTCTTGGGACGAGATGAATGTCCTGGCGCACGGACTCCTG  
CAGCTCGGCCAGGGGCTGCGCGAACACGCGGAGCGCACCCGCAGTCAGCTGAGCGCGCTGGA  
GCGGCGCCTGAGCGCGTGCGGGTCCGCCTGTCAGGGAACCGAGGGGTCCACCGACCTCCCGT  
TAGCCCCCTGAGAGCCGGGTGGACCCTGAGGTCCTTCACAGCCTGCAGACACAACCTCAAGGCT  
CAGAACAGCAGGATCCAGCAACTCTTCCACAAGGTGGCCCAGCAGCAGCGGCACCTGGAGAA  
GCAGCACCTGCGAATTCAGCATCTGCAAAGCCAGTTTGGCCTCCTGGACCACAAGCACCTAG  
ACCATGAGGTGGCCAAGCCTGCCCCAAGAAAGAGGCTGCCCCGAGATGGCCCAGCCAGTTGAC  
CCGGCTCACAATGTCAGCCGCTGCACCGGCTGCCCAGGGATTGCCAGGAGCTGTTCCAGGT  
TGGGGAGAGGCAGAGTGGACTATTTGAAATCCAGCCTCAGGGGTCTCCGCCATTTTTTGGTGA  
ACTGCAAGATGACCTCAGATGGAGGCTGGACAGTAATTCAGAGGCGCCACGATGGCTCAGTG  
GACTTCAACCGGCCCTGGGAAGCCTACAAGGCGGGGTTTGGGGATCCCCACGGCGAGTTCTG  
GCTGGGTCTGGAGAAGGTGCATAGCATCACGGGGGACCGCAACAGCCGCCTGGCCGTGCAGC  
TGCGGGACTGGGATGGCAACGCCGAGTTGCTGCAGTTCTCCGTGCACCTGGGTGGCGAGGAC  
ACGGCCTATAGCCTGCAGCTCACTGCACCCGTGGCCGGCCAGCTGGGCGCCACCACCGTCCC  
ACCCAGCGGCCTCTCCGTACCCTTCTCCACTTGGGACCAGGATCACGACCTCCGCAGGGACA  
AGAACTGCGCCAAGAGCCTCTCTGGAGGCTGGTGGTTTGGCACCTGCAGCCATTCCAACCTC  
AACGGCCAGTACTTCCGCTCCATCCCACAGCAGCGGCAGAAGCTTAAGAAGGGAATCTTCTG  
GAAGACCTGGCGGGGGCCGCTACTACCCGCTGCAGGCCACCACCATGTTGATCCAGCCCATGG  
CAGCAGAGGCAGCCTCCTAGCGTCCTGGCTGGGCCTGGTCCCAGGCCACGAAAGACGGTGA  
CTCTTGGCTCTGCCCCAGGATGTGGCCGTTCCCTGCCTGGGCAGGGGCTCCAAGGAGGGGCC  
ATCTGGAAACTTGTGGACAGAGAAGAAGACCACGACTGGAGAAGCCCCCTTTCTGAGTGCAG  
GGGGGCTGCATGCGTTGCCTCCTGAGATCGAGGCTGCAGGATATGCTCAGACTCTAGAGGCG  
TGGACCAAGGGGCATGGAGCTTCACTCCTTGCTGGCCAGGGAGTTGGGGACTCAGAGGGACC  
ACTTGGGGCCAGCCAGACTGGCCTCAATGGCGGACTCAGTCACATTGACTGACGGGGACCAG  
GGCTTGTGTGGGTCGAGAGCGCCCTCATGGTGCTGGTGCTGTTGTGTGTAGGTCCCCTGGGG  
ACACAAGCAGGCGCCAATGGTATCTGGGCGGAGCTCACAGAGTTCTTGAATAAAAGCAACC  
TCAGAACAC

## FIGURE 14

Signal peptide:	Amino acids 1-13
Transmembrane domain:	Amino acids 53-70
N-glycosylation site:	Amino acids 224-228
cAMP- and cGMP-dependent protein kinase phosphorylation sites:	Amino acids 46-50;118-122
N-myristoylation sites:	Amino acids 50-56;129-135; 341-347;357-363
Fibrinogen beta and gamma chains C-terminal domain signature:	Amino acids 396-409

MTVIRFFPAASATKRVLPVLRVSSPRTWNPNVPESPRIPAPRLPKRMSGAPTAGAALMLCA  
ATAVLLSAQGGPVQSKSPRFASWDEMNVLAHGLLQLGQGLREHAERTRSQLSALERRLSACG  
SACQGTEGSTDLPLAPESRVDPEVLHSLQTQLKAQNSRIQQLFHKVAQQQRHLEKQHLRIQH  
LQSQFGLLDHKHLDHEVAKPARRKRLPEMAQPVDPAHNVSRHLHRLPRDCQELFQVGERQSG  
FEIQPQGSPPFLVNCKMTSDGGWTVIQRRHDGSVDFNRPWEAYKAGFGDPHGEFWLGLEKVH  
SITGDRNSRLAVQLRDWDGNAELLQFSVHLGGEDTAYSLQLTAPVAGQLGATTVPPSGLSVP  
FSTWDQDHDLRDKNCAKSLSGGWFGTCSHNLNGQYFRSIPQQRQKLKKGIFWKTWRGRY  
YPLQATTMLIQPMAAEAAS

19/79

FIGURE 15

CGGACGCGTG GGGCCCCTGGTGGGCCCAGCAAGATGGATCTACTGTGGATCCTGCCCTCCCTG  
TGGCTTCTCCTGCTTGGGGGGCCTGCCTGCCTGAAGACCCAGGAACACCCAGCTGCCCAGG  
ACCCAGGGAACTGGAAGCCAGCAAAGTTGTCCTCCTGCCCAGTTGTCCCGGAGCTCCAGGAA  
GTCCTGGGGAGAAGGGAGCCCCAGGTCCTCAAGGGCCACCTGGACCACCAGGCAAGATGGGC  
CCCAAGGGTGAGCCAGGCCCCAGAACTGCCGGGAGCTGTTGAGCCAGGGCGCCACCTTGAG  
CGGCTGGTACCATCTGTGCCTACCTGAGGGCAGGGCCCTCCCAGTCTTTTGTGACATGGACA  
CCGAGGGGGGCGGCTGGCTGGTGTTCAGAGGCGCCAGGATGGTTCTGTGGATTTCTTCCCG  
TCTTGGTCCTCCTACAGAGCAGGTTTTGGGAACCAAGAGTCTGAATTCTGGCTGGGAAATGA  
GAATTTGCACCAGCTTACTCTCCAGGGTAACTGGGAGCTGCGGGTAGAGCTGGAAGACTTTA  
ATGGTAACCGTACTTTTCGCCCACTATGCGACCTTCCGCCTCCTCGGTGAGGTAGACCACTAC  
CAGCTGGCACTGGGCAAGTTCTCAGAGGGCACTGCAGGGGATTCCCTGAGCCTCCACAGTGG  
GAGGCCCTTTACCACCTATGACGCTGACCACGATTCAAGCAACAGCAACTGTGCAGTGATTG  
TCCACGGTGCCCTGGTGGTATGCATCCTGTTACCGATCAAATCTCAATGGTCGCTATGCAGTG  
TCTGAGGCTGCCGCCACAAATATGGCATTGACTGGGCCTCAGGCCGTGGTGTGGGCCACCC  
CTACCGCAGGGTTCGGATGATGCTTCGATAGGGCACTCTGGCAGCCAGTGCCCTTATCTCTC  
CTGTACAGCTTCCGGATCGTCAGCCACCTTGCCTTTGCCAACCACCTCTGCTTGCCTGTCCA  
CATTTAAAAATAAAATCATTTTAGCCCTTTCA

## FIGURE 16

Signal peptide: Amino acids 1-16  
N-glycosylation site: Amino acids 178-182  
Glycosaminoglycan attachment site: Amino acids 272-276  
Tyrosine kinase phosphorylation site:  
Amino acids 188-197  
N-myristoylation sites: Amino acids 16-22;89-95;  
144-150;267-273  
Fibrinogen beta and gamma chains C-terminal domain signature:  
Amino acids 242-255

MDLLWILPSLWLLLLGGPACLKTQEHPSCPGPRELEASKVLLPSCPGAPGSPGEKGAPGPQ  
GPPGPPGKMGPKEPGPRNCRELLSQGATLSGWYHLCLPEGRALPVFCDMDTEGGGWLVFQR  
RQDGSVDFFRSWSSYRAGFGNQSEFWLGNNLHQLTQGNWELRVELEDFNGNRTFAHYAT  
FRLLGESVDHYQLALGKFSEGTAGDSLHSGRPFTTYDADHDSSNSNCAVIVHGAWWYASCY  
RSNLNGRYAVSEAAAHKYGIDWASGRGVGHPYRRVRMMLR

FIGURE 17

TATTTACCATATCAGATTCACATTTCAGTCCTCAGCAAAATGAAGGGCTCCATTTTCACTCTG  
TTTTTATTCTCTGTCCTATTTGCCATCTCAGAAGTGCGGAGCAAGGAGTCTGTGAGACTCTG  
TGGGCTAGAATACATACGGACAGTCATCTATATCTGTGCTAGCTCCAGGTGGAGAAGGCATC  
TGGAGGGGATCCCTCAAGCTCAGCAAGCTGAGACAGGAACTCCTTCCAGCTCCACATAAA  
CGTGAGTTTTCTGAGGAAAATCCAGCGCAAAACCTTCCGAAGGTGGATGCCTCAGGGGAAGA  
CCGTCTTTGGGGTGGACAGATGCCCCACTGAAGAGCTTTGGAAGTCAAAGAAGCATTTCAGTGA  
TGTCAAGACAAGATTTACAACTTTGTGTTGCACTGATGGCTGTTCCATGACTGATTTGAGT  
GCTCTTTGCTTAAGACAAGAGCAAATACCCAATGGGTGGCAGAGCTTTATCACATGTTTAATT  
ACAGTGTTTTACTGCCTGGTAGAACACTAATATTGTGTTATTAAAATGATGGCTTTTGGGTA  
GGCAAACTTCTTTTCTAAAAGGTATAGCTGAGCGGTTGAAACCACAGTGATCTCTATTTTC  
TCCCTTTGCCAAGGTTAATGAACTGTTCTTTTCAAATTCTACTAATGCTTTGAAATTTCAA  
TGCTGCGCAAAATTGCAATAAAAATGCTATAAA



## FIGURE 18

Signal peptide: Amino acids 1-18  
cAMP- and cGMP-dependent protein kinase phosphorylation site:  
Amino acids 107-111  
N-myristoylation sites: Amino acids 3-9;52-58;96-102;  
125-131  
Insulin family signature: Amino acids 121-136

MKGSIFTLFLFSVLFAISEVRSKESVRLCGLEYIRTVIYICASSRWRRHLEGIPQAQQAETG  
NSFQLPHKREFSEENPAQNLPKVDASGEDRLWGGQMPTEELWKSKKHSVMSRQDLQTLCTD  
GCSMTDLSALC

FIGURE 19

CGGACGCGTG GGGGAAACCCCTTCCGAGAAAACAGCAACAAGCTGAGCTGCTGTGACAGAGGG  
GAACAAGATGGCGGCGCCGAAGGGGAGCCTCTGGGTGAGGACCCAACTGGGGCTCCCGCCGC  
TGCTGCTGCTGACCATGGCCTTGGCCGGAGGTTGCGGGACCGCTTCGGCTGAAGCATTTGAC  
TCGGTCTTGGGTGATACGGCGTCTTGCCACCGGGCCTGTCAGTTGACCTACCCCTTGACAC  
CTACCCTAAGGAAGAGGAGTTGTACGCATGTCAGAGAGGTTGCAGGCTGTTTTCAATTTGTC  
AGTTTGTGGATGATGGAATTGACTTAAATCGAACTAAATTGGAATGTGAATCTGCATGTACA  
GAAGCATATTCCCAATCTGATGAGCAATATGCTTGCCATCTTGGTTGCCAGAATCAGCTGCC  
ATTGCTGAACTGAGACAAGAACAATTATGTCCCTGATGCCAAAAATGCACCTACTCTTTT  
CTCTAACTCTGGTGAGGTCATTCTGGAGTGACATGATGGACTCCGCACAGAGCTTCATAACC  
TCTTCATGGACTTTTTATCTTCAAGCCGATGACGGAAAAATAGTTATATTCCAGTCTAAGCC  
AGAAATCCAGTACGCACCACATTTGGAGCAGGAGCCTACAAATTTGAGAGAATCATCTCTAA  
GCAAAATGTCCTATCTGCAAATGAGAAATTCACAAGCGCACAGGAATTTTCTGAAGATGGA  
GAAAGTGATGGCTTTTTAAGATGCCTCTCTCTTAACTCTGGGTGGATTTTAACTACAACCTCT  
TGTCTCTCGGTGATGGTATTGCTTTGGATTTGTTGTGCAACTGTTGCTACAGCTGTGGAGC  
AGTATGTTCCCTCTGAGAAGCTGAGTATCTATGGTGACTTGGAGTTTATGAATGAACAAAAG  
CTAAACAGATATCCAGCTTCTTCTCTTGTGGTTGTTAGATCTAAACTGAAGATCATGAAGA  
AGCAGGGCCTCTACCTACAAAAGTGAATCTTGCTCATTCTGAAATTTAAGCATTTTTCTTTT  
AAAAGACAAGTGTAATAGACATCTAAATTCCTCCTCATAGAGCTTTTAAATGGTTTCA  
TTGGATATAGGCCTTAAGAAATCACTATAAAATGCAAATAAAGTTACTCAAATCTGTG

## FIGURE 20

Signal peptide:	Amino acids 1-31
Transmembrane domain:	Amino acids 241-260
N-myristoylation sites:	Amino acids 28-34;29-35; 31-37;86-92
N-glycosylation site:	Amino acids 90-94

MAAPKGSLWVRTQLGLPPLLLLTMALAGGSGTASAEAFDSVLGDTASCHRAQCLTYPLHTYP  
KEEELYACQRCRLFSICQFVDDGIDLNRKLECESACTEAYSQSDEQYACHLGCQNQLPFA  
ELRQEQQLMSLMPKMHLFPLTLVRSFWSDMMDSAQSFITSSWTFYLQADDGKIVIFQSKPEI  
QYAPHLEQEPTNLRESSLSKMSYLQMRNSQAHRNFLEDGESDGFLRCLSLNSGWILTTTLVL  
SVMVLLWICCATVATAVEQYVPSEKLSIYGDLEFMNEQKLNRYPASSLVVVRSKTEDHEEAG  
PLPTKVNLAHSEI

FIGURE 21A

CGGACGCGTGGGCGGACGCGTGGGCGGACGCGTGGGCGGACGCGTGGGCTGGTTCAGGTCCA  
GGTTTTGCTTTGATCCTTTTCAAAAAGTGGAGACACAGAAGAGGGCTCTAGGAAAAAGTTTT  
GGATGGGATTATGTGGAACTACCTTGCATTCTCTGCTGCCAGAGCAGGCTCGGCGCTTCC  
ACCCAGTGCAGCCTTCCCCTGGCGGTGGTGAAAGAGACTCGGGAGTCGCTGCTTCCAAAGT  
GCCCCCGGTGAGTGAGCTCTCACCCAGTCAGCCAAATGAGCCTCTTCGGGCTTCTCCTGCT  
GACATCTGCCCTGGCCGGCCAGAGACAGGGGACTCAGGCGGAATCCAACCTGAGTAGTAAAT  
TCCAGTTTTCCAGCAACAAGGAACAGAACGGAGTACAAGATCCTCAGCATGAGAGAATTATT  
ACTGTGTCTACTAATGGAAGTATTCACAGCCCAAGGTTTCCTCATACTTATCCAAGAAATAC  
GGTCTTGGTATGGAGATTAGTAGCAGTAGAGGAAAATGTATGGATACAACTTACGTTTGATG  
AAAGATTTGGGCTTGAAGACCCAGAAGATGACATATGCAAGTATGATTTTGTAGAAGTTGAG  
GAACCCAGTGATGGAACCTATATTAGGGCGCTGGTGTGGTTCTGGTACTGTACCAGGAAAACA  
GATTTCTAAAGGAAATCAAATTAGGATAAGATTTGTATCTGATGAATATTTTCCTTCTGAAC  
CAGGGTTCTGCATCCACTACAACATTGTCATGCCACAATTCACAGAAGCTGTGAGTCCTTCA  
GTGCTACCCCCTTCAGCTTTGCCACTGGACCTGCTTAATAATGCTATAACTGCCTTTAGTAC  
CTTGAAGACCTTATTCGATATCTTGAACCAGAGAGATGGCAGTTGGACTTAGAAGATCTAT  
ATAGGCCAACTTGGCAACTTCTTGGCAAGGCTTTTGTGTTTTTGAAGAAAATCCAGAGTGGTG  
GATCTGAACCTTCTAACAGAGGAGGTAAGATTATACAGCTGCACACCTCGTAACTTCTCAGT  
GTCCATAAGGGAAGAACTAAAGAGAACCGATACCATTTTCTGGCCAGGTTGTCTCCTGGTTA  
AACGCTGTGGTGGGAACTGTGCCTGTTGTCTCCACAATTGCAATGAATGTCAATGTGTCCCA  
AGCAAAGTTACTAAAAAATACCACGAGGTCCTTCAGTTGAGACCAAAGACCGGTGTCAGGGG  
ATTGCACAAATCACTACCGACGTGGCCCTGGAGCACCATGAGGAGTGTGACTGTGTGTGCAGA  
GGGAGCACAGGAGGATAGCCGCATCACCACCAGCAGCTCTTGCCAGAGCTGTGCAGTGCAG  
TGGCTGATTCTATTAGAGAACGTATGCGTTATCTCCATCCTTAATCTCAGTTGTTTGCTTCA  
AGGACCTTTCATCTTCAGGATTTACAGTGCATTCTGAAAGAGGAGACATCAAACAGAATTAG  
GAGTTGTGCAACAGCTCTTTTGAGAGGAGGCCTAAAGGACAGGAGAAAAGGTCTTCAATCGT  
GGAAAGAAAATTAAATGTTGTATTAAATAGATCACCAGCTAGTTTCAGAGTTACCATGTACG  
TATTCCACTAGCTGGGTTCTGTATTTAGTTCTTTCGATACGGCTTAGGGTAATGTCAGTAC  
AGGAAAAAACTGTGCAAGTGAGCACCTGATTCCGTTGCCTTGCTTAACCTCTAAAGCTCCAT  
GTCCTGGGCCTAAAATCGTATAAAATCTGGATTTTTTTTTTTTTTTTGTCTCATATTCACAT  
ATGTAAACCAGAACATTCATGTACTACAAACCTGGTTTTTTTAAAAAGGAAGTATGTTGCTAT  
GAATTAACTTGTGTCTGCTGATAGGACAGACTGGATTTTTCATATTTCTTATTAAATTT

FIGURE 21B

CTGCCATTTAGAGAAGAGAACTACATTCATGGTTTGGAAGAGATAAACCTGAAAAGAAGAG  
TGGCCTTATCTTCACTTTATCGATAAGTCAGTTTATTTGTTTCATTGTGTACATTTTTATAT  
TCTCCTTTTGACATTATAACTGTTGGCTTTTCTAATCTTGTTAAATATATCTATTTTTACCA  
AAGGTATTTAATATTCCTTTTTTATGACAACTTAGATCAACTATTTTTAGCTTGGTAAATTTT  
TCTAAACACAATTGTTATAGCCAGAGGAACAAAGATGATATAAAATATTGTTGCTCTGACAA  
AAATACATGTATTTCACTCTCGTATGGTGCTAGAGTTAGATTAATCTGCATTTTAAAAAACT  
GAATTGGAATAGAATTGGTAAGTTGCAAAGACTTTTTGAAAATAATTAAATTATCATATCTT  
CCATTCTGTATTATTGGAGATGAAAATAAAAAGCAACTTATGAAAGTAGACATTCAGATCCAG  
CCATTACTAACCTATTCCTTTTTTGGGGAAATCTGAGCCTAGCTCAGAAAAACATAAAGCAC  
CTTGAAAAGACTTGGCAGCTTCCTGATAAAGCGTGCTGTGCTGTGCAGTAGGAACACATCC  
TATTTATTGTGATGTTGTGGTTTTATTATCTTAAACTCTGTTCCATACACTTGTATAAATACATG  
GATATTTTTATGTACAGAAGTATGTCTCTTAACCAGTTCACTTATTGTACTCTGGCAATTTA  
AAAGAAAATCAGTAAAATATTTTGCTTGTAATAATGCTTAATATNGTGCCTAGGTTATGTGGT  
GACTATTTGAATCAAAAATGTATTGAATCATCAAATAAAGAATGTGGCTATTTTGGGGAGA  
AAATTAAAAAAGGTTTAGGGATAACAGGGTAATGCGGCC

## FIGURE 22

Signal peptide:	Amino acids 1-14
N-glycosylation sites:	Amino acids 25-29;55-59; 254-258
N-myristoylation sites:	Amino acids 15-21;117-123; 127-133;281-287;282-288; 319-325
Amidation site:	Amino acids 229-233

MSLFGLLLLTSALAGQRQGTQAESNLSSKFQFSSNKEQNGVQDPQHERIITVSTNGSIHSPR  
FPHTYPRNTVLVWRLVAVEENVWQLTFDERFGLEDPEDDICKYDFVEVEEPSDGTILGRWC  
GSGTVPGKQISKGNQIRIRFVSDEYFPSEPGFCIHYNIVMPQFTEAVSPSVLPPSALPLDLL  
NNAITAFSTLEDLIRYLEPERWQLDLEDLYRPTWQLLGKAFVFGRKSRVVDLNNLLTEEVRLY  
SCTPRNFSVSIREELKRDTIFWPGCLLVKRCGGNCACCLHNCNECQCVPSKVTKKYHEVLQ  
LRPKTGVRGLHKSLTDVALEHHEECDVCVRGSGTGG

FIGURE 23

GGCCGGAGCAGCACGGCCGCAGGACCTGGAGCTCCGGCTGCGTCTTCCCGCAGCGCTACCCG  
CCATGCGCCTGCCGCGCCGGGCGCGCTGGGGCTCCTGCCGCTTCTGCTGCTGCTGCCGCCC  
GCGCCGGAGGCCGCCAAGAAGCCGACGCCCTGCCACCGGTGCCGGGGGCTGGTGGACAAGTT  
TAACCAGGGGATGGTGGACACCGCAAAGAAGAACTTTGGCGGCGGGAACACGGCTTGGGAGG  
AAAAGACGCTGTCCAAGTACGAGTCCAGCGAGATTCGCCTGCTGGAGATCCTGGAGGGGCTG  
TGCGAGAGCAGCGACTTCGAATGCAATCAGATGCTAGAGGCGCAGGAGGAGCACCTGGAGGC  
CTGGTGGCTGCAGCTGAAGAGCGAATATCCTGACTTATTTCGAGTGGTTTTGTGTGAAGACAC  
TGAAAGTGTGCTGCTCTCCAGGAACCTACGGTCCCGACTGTCTCGCATGCCAGGGCGGATCC  
CAGAGGCCCTGCAGCGGAATGGCCACTGCAGCGGAGATGGGAGCAGACAGGGCGACGGGTC  
CTGCCGGTGCCACATGGGGTACCAGGGCCCCGCTGTGCACTGACTGCATGGACGGCTACTTCA  
GCTCGCTCCGGAACGAGACCCACAGCATCTGCACAGCCTGTGACGAGTCTTGCAAGACGTGC  
TCGGGCCTGACCAACAGAGACTGCGGCGAGTGTGAAGTGGGCTGGGTGCTGGACGAGGGCGC  
CTGTGTGGATGTGGACGAGTGTGCGGCCGAGCCGCCTCCCTGCAGCGCTGCGCAGTTCTGTA  
AGAACGCCAACGGCTCCTACACGTGCGAAGAGTGTGACTCCAGCTGTGTGGGCTGCACAGGG  
GAAGGCCCAGGAAACTGTAAAGAGTGTATCTCTGGCTACGCGAGGGAGCACGGACAGTGTGC  
AGATGTGGACGAGTGTCTACTAGCAGAAAAACCTGTGTGAGGAAAAACGAAAACCTGCTACA  
ATACTCCAGGGAGCTACGTCTGTGTGTGTCTTGACGGCTTCGAAGAAACGGAAGATGCCTGT  
GTGCCCGCCGGCAGAGGCTGAAGCCACAGAAGGAGAAAGCCCGACACAGCTGCCCTCCCGCGA  
AGACCTGTAATGTGCCGGACTTACCCTTTAAATTATTCAGAAGGATGTCCCGTGGAAAATGT  
GGCCCTGAGGATGCCGTCTCCTGCAGTGGACAGCGGCGGGGAGAGGCTGCCTGCTCTCTAAC  
GGTTGATTCTCATTTGTCCCTTAAACAGCTGCATTTCTTGGTTGTTCTTAAACAGACTTGTA  
TATTTTGATACAGTCTTTGTAATAAAATTGACCATTGTAGGTAATCAGGAGGAAAAA

## FIGURE 24

Signal peptide:	Amino acids 1-24
N-glycosylation sites:	Amino acids 190-194;251-255
Glycosaminoglycan attachment sites:	Amino acids 149-153;155-159
cAMP- and cGMP-dependent protein kinase phosphorylation site:	Amino acids 26-30
Tyrosine kinase phosphorylation site:	Amino acids 303-310
N-myristoylation sites:	Amino acids 44-50;54-60; 55-61;81-87;150-156;158-164; 164-170;252-258;313-319
Aspartic acid and asparagine hydroxylation site:	Amino acids 308-320
EGF-like domain cysteine pattern signature:	Amino acids 166-178
Leucine zipper pattern:	Amino acids 94-116

MRLPRRAALGLLPLLLLLLPPAPEAAKKPTPCHRCRGLVDKFNQGMVDTAKKNFGGGNTAWEE  
 KTLISKYESSEIRLLEILEGLCESSDFECNQMLEAQEEHLEAWWLQLKSEYPDLFEWFVCVCTL  
 KVCCSPGTYGPDCLACQGGGSRPCSGNGHCSGDGSRQGDGSCRCHMGYQGGLCTDCMDGYFS  
 SLRNETHSICTACDESKTCSGLTNRDCGECEVGWVLDEGACVDVDECAAEPPPCSAAQFCK  
 NANGSYTCEECDSSCVGCTGEGPGNCKECISGYAREHGQCADVDECSLAEKTCVRKNENCYN  
 TPGSYVCVCPDGFEEEDACVPPAEAEATEGESPTQLPSREDL



FIGURE 25A

CCAGGCCGGGAGGCGACGCGCCCAGCCGTCTAAACGGGAACAGCCCTGGCTGAGGGAGCTGC  
AGCGCAGCAGAGTATCTGACGGCGCCAGGTTGCGTAGGTGCGGCACGAGGAGTTTTCCCGGC  
AGCGAGGAGGTCCTGAGCAGCATGGCCCCGGAGGAGCGCCTTCCCTGCCGCCGCGCTCTGGCT  
CTGGAGCATCCTCCTGTGCCTGCTGGCACTGCGGGCGGAGGCCGGGCCCGCAGGAGGAGA  
GCCTGTACCTATGGATCGATGCTCACCAGGCAAGAGTACTCATAGGATTTGAAGAAGATATC  
CTGATTGTTTCAGAGGGGAAAATGGCACCTTTTACACATGATTTTCAGAAAAGCGCAACAGAG  
AATGCCAGCTATTCCTGTCAATATCCATTCCATGAATTTTACCTGGCAAGCTGCAGGGCAGG  
CAGAATACTTCTATGAATTCCTGTCCTTGCGCTCCCTGGATAAAGGCATCATGGCAGATCCA  
ACCGTCAATGTCCCTCTGCTGGGAACAGTGCCTCACAAGGCATCAGTTGTTCAAGTTGGTTT  
CCCATGTCTTGGAACAGGATGGGGTGGCAGCATTTGAAGTGGATGTGATTGTTATGAATT  
CTGAAGGCAACACCATTCTCCAAACACCTCAAAATGCTATCTTCTTTAAACATGTCAACAA  
GCTGAGTGCCCGAGGCGGGTGCCGAAATGGAGGCTTTTGTAATGAAAGACGCATCTGCGAGTG  
TCCTGATGGGTTCCACGGACCTCACTGTGAGAAAGCCCTTTGTACCCACGATGTATGAATG  
GTGGACTTTGTGTGACTCCTGGTTTCTGCATCTGCCCACCTGGATTCTATGGAGTGAAGTGT  
GACAAAGCAAACCTGCTCAACCACCTGCTTTAATGGAGGGACCTGTTTCTACCCTGGAAAATG  
TATTTGCCCTCCAGGACTAGAGGGAGAGCAGTGTGAAATCAGCAAATGCCCAACCCCTGTC  
GAAATGGAGGTAAATGCATTGGTAAAAGCAAATGTAAGTGTTCAAAGGTTACCAGGGAGAC  
CTCTGTTCAAAGCCTGTCTGCGAGCCTGGCTGTGGTGCACATGGAACCTGCCATGAACCCAA  
CAAAATGCCAATGTCAAGAAGGTTGGCATGGAAGACACTGCAATAAAAGGTACGAAGCCAGCC  
TCATACATGCCCTGAGGCCAGCAGGCGCCAGCTCAGGCAGCACACGCCTTCACTTAAAAAG  
GCCGAGGAGCGGCGGGATCCACCTGAATCCAATTACATCTGGTGAACTCCGACATCTGAAAC  
GTTTTAAGTTACACCAAGTTCATAGCCTTTGTAAACCTTTCATGTGTTGAATGTTCAAATAA  
TGTTCAATTACACTTAAGAATACTGGCCTGAATTTTATTAGCTTCATTATAAATCACTGAGCT  
GATATTTACTCTTCCTTTTAAGTTTTCTAAGTACGTCTGTAGCATGATGGTATAGATTTTCT  
TGTTTCAGTGCTTTGGGACAGATTTTATATTATGTCAATTGATCAGGTAAAATTTTCAGTG  
TGTAAGTTGGCAGATATTTTCAAAATTACAATGCATTTATGGTGTCTGGGGGCAGGGGAACAT  
CAGAAAGGTTAAATTGGGCAAAATGCGTAAGTCACAAGAATTGGATGGTGCAGTTAATGT  
TGAAGTTACAGCATTTTCAGATTTTATTGTCAGATATTTAGATGTTTGTACATTTTAAAAA  
TTGCTCTTAATTTTAACTCTCAATACAATATATTTTGACCTTACCATTATTCCAGAGATT  
CAGTATTAAAAAATAAATTACACTGTGGTAGTGGCATTAAACAATATAATATATTCTA  
AACACAATGAAATAGGGAATATAATGTATGAACTTTTGCATTGGCTTGAAGCAATATAATA

FIGURE 25B

TATTGTAAACAAAACACAGCTCTTACCTAATAAACATTTTATACTGTTTGTATGTATAAAAT  
AAAGGTGCTGCTTTAGTTTTTTGGAAAAAAAAAAAAAAAAAAAAAAAAAAAA

## FIGURE 26

Signal peptide: Amino acids 1-28  
N-glycosylation sites: Amino acids 88-92;245-249  
Tyrosine kinase phosphorylation site:  
Amino acids 370-378  
N-myristoylation sites: Amino acids 184-190;185-191;  
189-195;315-321  
ATP/GTP-binding site motif A (P-loop):  
Amino acids 285-293  
EGF-like domain cysteine pattern signatures:  
Amino acids 198-210;230-242;  
262-274;294-306;326-338

MARRSAFPAAALWLWSILLCLLALRAEAGPPQEESLYLWIDAHQARVLIGFEEDILIVSEGK  
MAPFTHDFRKAQQORMPAIPVNIHSMNFTWQAAGQAEYFYEFSLRSLDKGIMADPTVNVPLL  
GTVPHKASVVQVGFPCLGKQDGVAAFEVDVIVMNSEGNLILQTPQNAIFFKTCQQAECPPGGC  
RNGGFCNERRICECPDGFHGPHEKALCTPRCMNGGLCVTPGFCICPPGFYGVNCDKANCST  
TCFNGGTCFYPGKCICPPGLEGEQCEISKCPQPCRNGGKCIGKSKCKCSKGYQGDLCSKPVC  
EPGCGAHGTCHEPNKCQCQEGWHGRHCNKRYEASLIHALRPAGAQLRQHPTSLKKAEEERRDP  
PESNYIW

FIGURE 27A

ACTTGGAGCAAGCGGCGGCGGAGACAGAGGCAGAGGCAGAAGCTGGGGCTCCGTCCTCG  
CCTCCCACGAGCGATCCCCGAGGAGAGCCGCGGCCCTCGGCGAGGCGAAGAGGCCGACGAGG  
AAGACCCGGGTGGCTGCGCCCCCTGCCTCGCTTCCCAGGCGCCGGCGGCTGCAGCCTTGCCCC  
TCTTGCTCGCCTTGAAAATGGAAAAGATGCTCGCAGGCTGCTTTCTGCTGATCCTCGGACAG  
ATCGTCCTCCTCCCTGCCGAGGCCAGGGAGCGGTACGTGGGAGGTCCATCTCTAGGGGCAG  
ACACGCTCGGACCCACCCGACAGCGGCCCTTCTGGAGAGTTCTGTGAGAACAAGCGGGCAG  
ACCTGGTTTTTCATCATTGACAGCTCTCGCAGTGTCAACACCCATGACTATGCAAAGGTCAAG  
GAGTTCATCGTGGACATCTTGCAATTCTTGACATTGGTCTCTGATGTCACCCGAGTGGGCCT  
GCTCCAATATGGCAGCACTGTCAAGAATGAGTTCTCCCTCAAGACCTTCAAGAGGAAGTCCG  
AGGTGGAGCGTGCTGTCAAGAGGATGCGGCATCTGTCCACGGGCACCATGACTGGGCTGGCCATC  
CAGTATGCCCTGAACATCGCATTCTCAGAAGCAGAGGGGGCCCGGCCCTGAGGGAGAATGT  
GCCACGGGTCATAATGATCGTGACAGATGGGAGACCTCAGGACTCCGTGGCCGAGGTGGCTG  
CTAAGGCACGGGACACGGGCATCCTAATCTTTGCCATTGGTGTGGGCCAGGTAGACTTCAAC  
ACCTTGAAGTCCATTGGGAGTGAGCCCCATGAGGACCATGTCTTCCTTGTGGCCAATTTGAG  
CCAGATTGAGACGCTGACCTCCGTGTTCCAGAAGAAGTTGTGCACGGCCCACATGTGCAGCA  
CCCTGGAGCATAACTGTGCCCACTTCTGCATCAACATCCCTGGCTCATACTGTGCAGGTGC  
AAACAAGGCTACATTCTCAACTCGGATCAGACGACTTGCAGAAATCCAGGATCTGTGTGCCAT  
GGAGGACCACAACCTGTGAGCAGCTCTGTGTGAATGTGCCGGGCTCCTTCGTCTGCCAGTGCT  
ACAGTGGCTACGCCCTGGCTGAGGATGGGAAGAGGTGTGTGGCTGTGGACTACTGTGCCTCA  
GAAAACACGGATGTGAACATGAGTGTGTAAATGCTGATGGCTCCTACCTTTGCCAGTGCCA  
TGAAGGATTTGCTCTTAACCCAGATGAAAAACGTGCACAAGGATCAACTACTGTGCACTGA  
ACAAACCGGGCTGTGAGCATGAGTGCCTAACATGGAGGAGAGCTACTACTGCCGCTGCCAC  
CGTGGCTACACTCTGGACCCCAATGGCAAAACCTGCAGCCGAGTGGACCACTGTGCACAGCA  
GGACCATGGCTGTGAGCAGCTGTGTCTGAACACGGAGGATTCCTTCGTCTGCCAGTGCTCAG  
AAGGCTTCCTCATCAACGAGGACCTCAAGACCTGCTCCCGGGTGGATTACTGCCTGCTGAGT  
GACCATGGTTGTGAATACTCCTGTGTCAACATGGACAGATCCTTTGCCTGTCAGTGTCTGA  
GGGACACGTGCTCCGCAGCGATGGGAAGACGTGTGCAAAATTGGACTCTTGTGCTCTGGGGG  
ACCACGGTTGTGAACATTTCGTGTGTAAGCAGTGAAGATTTCGTTTGTGTGCCAGTGCTTTGAA  
GGTTATATACTCCGTGAAGATGGAAAAACCTGCAGAAGGAAAGATGTCTGCCAAGCTATAGA  
CCATGGCTGTGAACACATTTGTGTGAACAGTGACGACTCATAACGTGCGAGTGCTTGGAGG  
GATTCCGGCTCGCTGAGGATGGGAAACGCTGCCGAAGGAAGGATGTCTGCAAATCAACCCAC  
CATGGCTGCGAACACATTTGTGTTAATAATGGGAATTCCTACATCTGCAAATGCTCAGAGGG  
ATTTGTTCTAGCTGAGGACGGAAGACGGTGCAAGAAATGCACTGAAGGCCCAATTGACCTGG

FIGURE 27B

TCTTTGTGATCGATGGATCCAAGAGTCTTGGAGAAGAGAATTTTGAGGTCGTGAAGCAGTTT  
GTCACTGGAATTATAGATTCCCTTGACAATTTCCCCCAAAGCCGCTCGAGTGGGGCTGCTCCA  
GTATTCCACACAGGTCCACACAGAGTTCACTCTGAGAACTTCAACTCAGCCAAAGACATGA  
AAAAAGCCGTGGCCACATGAAATACATGGGAAAGGGCTCTATGACTGGGCTGGCCCTGAAA  
CACATGTTTGAGAGAAGTTTTACCCAAGGAGAAGGGGCCAGGCCCTTCCACAAGGGTGCC  
CAGAGCAGCCATTGTGTTCCACCGACGGACGGGCTCAGGATGACGTCTCCGAGTGGGCCAGTAAA  
GCCAAGGCCAATGGTATCACTATGTATGCTGTTGGGGTAGGAAAAGCCATTGAGGAGGAACT  
ACAAGAGATTGCCTCTGAGCCCACAAACAAGCATCTCTTCTATGCCGAAGACTTCAGCACAA  
TGGATGAGATAAGTGAAAACTCAAGAAAGGCATCTGTGAAGCTCTAGAAGACTCCGATGGA  
AGACAGGACTCTCCAGCAGGGGAACTGCCAAAACGGTCCAACAGCCAACAGAATCTGAGCC  
AGTCACCATAAATATCCAAGACCTACTTTCCTGTTCTAATTTTGCAGTGCAACACAGATATC  
TGTTTGAAGAAGACAATCTTTTACGGTCTACACAAAAGCTTTCCCATTCACAAAACCTTCA  
GGAAGCCCTTTGGAAGAAAAACACGATCAATGCAAATGTGAAAACCTTATAATGTTCCAGAA  
CCTTGCAAACGAAGAAGTAAGAAAATTAACACAGCGCTTAGAAGAAATGACACAGAGAATGG  
AAGCCCTGGAAAATCGCCTGAGATACAGATGAAGATTAGAAATCGCGACACATTTGTAGTCA  
TTGTATCACGGATTACAATGAACGCAGTGCAGAGCCCCAAAGCTCAGGCTATTGTTAAATCA  
ATAATGTTGTGAAGTAAAACAATCAGTACTGAGAAACCTGGTTTGCCACAGAACAAAGACAA  
GAAGTATACACTAACTTGTATAAATTTATCTAGGAAAAAAATCCTTCAGAAATCTAAGATGA  
ATTTACCAGGTGAGAATGAATAAGCTATGCAAGGTATTTTGTAAATACTGTGGACACAACT  
TGCTTCTGCCTCATCCTGCCTTAGTGTGCAATCTCATTTGACTATACGATAAAGTTTGCACA  
GTCTTACTTCTGTAGAACACTGGCCATAGGAAATGCTGTTTTTTTGTACTGGACTTTACCTT  
GATATATGTATATGGATGTATGCATAAAATCATAGGACATATGTACTTGTGGAACAAGTTGG  
ATTTTTTATACAATATTAAAATTCACCACTTCAG

FIGURE 28

Signal peptide:	Amino acids 1-23
N-glycosylation site:	Amino acids 221-225
cAMP- and cGMP-dependent protein kinase phosphorylation sites:	Amino acids 115-119;606-610; 892-896
N-myristoylation sites:	Amino acids 133-139;258-291; 299-305;340-346;453-459; 494-500;639-645;690-696; 752-758;792-798
Amidation sites:	Amino acids 314-318;560-564; 601-605
Aspartic acid and asparagine hydroxylation sites:	Amino acids 253-265;294-306; 335-347;376-388;417-429; 458-470;540-552;581-593

MEKMLAGCFLLLILGQIVLLPAEARERSRGRSISRGRHARTHPTALLESSCENKRADLVFII  
DSSRSVNTHDYAKVKEFIVDILQFLDIGPDVTRVGLLQYGSTVKNEFSLKTFKRKSEVERAV  
KMRHLSTGTMGLAIQYALNIAFSEAEGARPLRENVPRVIMIVTDGRPQDSVAEVAAKARD  
TGILIFAIGVGQVDFNTLKSIGSEPHEDHVFLVANFSQIETLTSVFQKKLCTAHMCSTLEHN  
CAHFCINIPGSYVCRCKQGYILNSDQTTCRIQDLCAMEDHNCEQLCVNVPGSFVCQCYSGYA  
LAEDGKRCVAVDYCASENHGCEHECVNADGSYLCQCHEGFALNPDEKTCINTRINYCALNKPGE  
EHECVNMEESYCRCHRGYTLDPNGKTC SRVDHCAQQDHGCEQLCLNTEDSFVCQCSEGFLI  
NEDLKTC SRVDYCLLSDHGCEYSCVNMDRSFACQCPEGHVLRSDGKTC AKLDSCALGDHGCE  
HSCVSSSEDSFVCQCFEGYILREDGKTCRRKDVCQAIDHGCEHICVNSDDSYTCECLEGFRLA  
EDGKRCRRKDVCKSTHHGCEHICVNNGNSYICKCSEGFVLAEDGRRCKKCTEGPIDLVFVID  
GSKSLGEENFEVVKQFVTGIIDSLTISPKAARVGLLQYSTQVHTEFTLRNFNSAKDMKKAVA  
HMKYMGKGSMTGLALKHMFERSFTQGEGARPLSTRVPRAAIVFTDGRAQDDVSEWASKAKAN  
GITMYAVGVGKAIEEELQEIASPTNKLHLYAEDFSTMDIESEKLKKGICEALEDSDGRQDS  
PAGELPKTVQQPTESEPVTINIQDLLSCSNFAVQHRYLFEEDNLLRSTQKLSHSTKPSGSPL  
EEKHDQCKCENLIMFQNLANEEVRKLTQRLEEMTQRMEALENRLRYR

FIGURE 29

GCCCGGGACTGGCGCAAGGTGCCCAAGCAAGGAAAGAAATAATGAAGAGACACATGTGTTAG  
CTGCAGCCTTTTGAACACGCAAGAAGGAAATCAATAGTGTGGACAGGGCTGGAACCTTTAC  
CACGCTTGTTGGAGTAGATGAGGAATGGGCTCGTGATTATGCTGACATTCCAGCATGAATCT  
GGTAGACCTGTGGTTAACCCGTTCCCTCTCCATGTGTCTCCTCCTACAAAGTTTTGTTCTTA  
TGATACTGTGCTTTCATTCTGCCAGTATGTGTCCCAAGGGCTGTCTTTGTTCTTCCTCTGGG  
GGTTTAAATGTCACCTGTAGCAATGCAAATCTCAAGGAAATACCTAGAGATCTTCCTCCTGA  
AACAGTCTTACTGTATCTGGACTCCAATCAGATCACATCTATTCCCAATGAAATTTTAAAGG  
ACCTCCATCAACTGAGAGTTCTCAACCTGTCCAAAATGGCATTGAGTTTATCGATGAGCAT  
GCCTTCAAAGGAGTAGCTGAAACCTTGCACTCTGGACTTGTCCGACAATCGGATTCAAAG  
TGTGCACAAAAATGCCTTCAATAACCTGAAGGCCAGGGCCAGAATTGCCAACAACCCCTGGC  
ACTGCGACTGTACTCTACAGCAAGTTCTGAGGAGCATGGCGTCCAATCATGAGACAGCCCAC  
AACGTGATCTGTAAAACGTCCGTGTTGGATGAACATGCTGGCAGACCATTCTCAATGCTGC  
CAACGACGCTGACCTTTGTAACCTCCCTAAAAAACTACCGATTATGCCATGCTGGTCACCA  
TGTTTGGCTGGTTCATATGGTGATCTCATATGTGGTATATTATGTGAGGCAAAATCAGGAG  
GATGCCCCGAGACACCTCGAATACTTGAAATCCCTGCCAAGCAGGCAGAAGAAAGCAGATGA  
ACCTGATGATATTAGCACTGTGGTATAGTGTGCCAACTGACTGTCATTGAGAAAGAAAGAAA  
GTAGTTTGCGATTGCAGTAGAAATAAGTGGTTTACTTCTCCCATCCATTGTAAACATTTGAA  
ACTTTGTATTTTCAAGTTTTTTTGAATTATGCCACTGCTGAACTTTTAACAAACACTACAACA  
TAAATAATTTGAGTTTAGGTGATCCACCCCTTAATTGTACCCCGATGGTATATTTCTGAGT  
AAGCTACTATCTGAACATTAGTTAGATCCATCTCACTATTTAATAATGAAATTTATTTTTTT  
AATTTAAAAGCAAATAAAAGCTTAACCTTTGAACCATGGGAAAAAAAAAAAAAAAAAAAAACA

## FIGURE 30

Signal peptide:	Amino acids 1-33
Transmembrane domain:	Amino acids 206-225
N-glycosylation sites:	Amino acids 47-51;94-98
cAMP- and cGMP-dependent protein kinase phosphorylation site:	Amino acids 199-203
N-myristoylation sites:	Amino acids 37-43;45-51; 110-116

MNLVDLWLTRSLMCLLLQSFLMILCFHSASMCPKGCLCSSSGGLNVTCSNANLKEIPRDL  
PPETVLLYLDSNQITSIPNEIFKDLHQLRVLNLSKNGIEFIDEHAFKGVAETLQTLDSLSDNR  
IQSVHKNAFNNLKARARIANNPWHCDCTLQQVLRSMASNHETAHNVICKTSVLDEHAGRPFL  
NAANDADLCNLPPKTTDYAMLVTMFGWFTMVISYVVYYVRQNQEDARRHLEYLKSLSRQKK  
ADEPDDISTVV



FIGURE 31

CAGCGCGTGGCCGGCGCCGCTGTGGGGACAGCATGAGCGGCGGTTGGATGGCGCAGGTTGGA  
GCGTGGCGAACAGGGGCTCTGGGCCTGGCGCTGCTGCTGCTCGGCCTCGGACTAGGCCT  
GGAGGCCGCCGCGAGCCCGCTTTCCACCCCGACCTCTGCCCAGGCCGCGAGCCCCAGCTCAG  
GCTCGTGCCACCCACCAAGTTCCAGTGCCGCACCAAGTGGCTTATGCGTGCCCTCACCTGG  
CGCTGCGACAGGGACTTGACTGCAGCGATGGCAGCGATGAGGAGGAGTGCAGGATTGAGCC  
ATGTACCCAGAAAGGGCAATGCCCACCGCCCCCTGGCCTCCCCTGCCCCTGCACCGGCGTCA  
GTGACTGCTCTGGGGGAAGTACAAGAACTGCGCAACTGCAGCCGCCTGGCCTGCCTAGCA  
GGCGAGCTCCGTTGCACGCTGAGCGATGACTGCATTCCACTCACGTGGCGCTGCGACGGCCA  
CCCAGACTGTCCCGACTCCAGCGACGAGCTCGGCTGTGGAACCAATGAGATCCTCCCGGAAG  
GGGATGCCACAACCATGGGGCCCCCTGTGACCCTGGAGAGTGTACCTCTCTCAGGAATGCC  
ACAACCATGGGGCCCCCTGTGACCCTGGAGAGTGTCCCCTCTGTGCGGAATGCCACATCCTC  
CTCTGCCGGAGACCAGTCTGGAAGCCCAACTGCCTATGGGGTTATTGCAGCTGCTGCGGTGC  
TCAGTGCAAGCCTGGTCACCGCCACCCTCCTCCTTTTGTCTGGCTCCGAGCCCAGGAGCGC  
CTCCGCCCCACTGGGGTTACTGGTGGCCATGAAGGAGTCCCTGCTGCTGTGAGAACAGAAGAC  
CTCGCTGCCCCTGAGGACAAGCACTTGCCACCACCGTCACTCAGCCCTGGGCGTAGCCGGACA  
GGAGGAGAGCAGTGATGCGGATGGGTACCCGGGCACACCAGCCCTCAGAGACCTGAGTTCTT  
CTGGCCACGTGGAACCTCGAACCCGAGCTCCTGCAGAAGTGGCCCTGGAGATTGAGGGTCCC  
TGGACACTCCCTATGGAGATCCGGGGAGCTAGGATGGGGAACCTGCCACAGCCAGAAGT  
GGGCTGGCCCCAGGCAGCTCCAGGGGGTAGAACGGCCCTGTGCTTAAGACACTCCCTGCTG  
CCCCGTCTGAGGGTGGCGATTAAAGTTGCTTC

## FIGURE 32

Signal peptide:	Amino acids 1-30
Transmembrane domain:	Amino acids 231-248
N-glycosylation sites:	Amino acids 126-130;195-199; 213-217
N-myristoylation sites:	Amino acids 3-9;10-16;26-32; 30-36;112-118;166-172; 212-218;224-230;230-236; 263-269
Prokaryotic membrane lipoprotein lipid attachment site:	Amino acids 44-55
Leucine zipper pattern:	Amino acids 17-39

MSGGWMAQVGAWRTGALGLALLLLLGLGLGLEAAASPLSTPTSAQAAGPSSGSCPPTKFQCR  
TSGLCVPLTWRCRDLDCSDGSDEEEECRIEPTQKGQCPPPPGLPCPCTGVSDCSGGTDKKL  
RNC SRLACLAGELRCTLSDDCIPLTWRC DGHPCPDSSDELGCGTNEILPEGDATTMGPPVT  
LESVTSLRNATTMGPPVTLESVPSVGNATSSSAGDQSGSPTAYGVIAAAVLSASLVTATLL  
LLSWLRAQERLRPLGLLVAMKESLLLSEQKTSLP

FIGURE 33A

CGCCACCACTGCGGCCACCGCCAATGAAACGCCTCCCGCTCCTAGTGGTTTTTTTCCACTTTG  
TTGAATTGTTCTTACTCAAAATTGCACCAAGACACCTTGTCTCCCAAATGCAAAATGTGA  
AATACGCAATGGAATTGAAGCCTGCTATTGCAACATGGGATTTTCAGGAAATGGTGTACAA  
TTTGTGAAGATGATAATGAATGTGGAAATTTAACTCAGTCCTGTGGCGAAAATGCTAATTGC  
ACTAACACAGAAGGAAGTTATTATTGTATGTGTGTACCTGGCTTCAGATCCAGCAGTAACCA  
AGACAGGTTTATCACTAATGATGGAACCGTCTGTATAGAAAATGTGAATGCAAACTGCCATT  
TAGATAATGTCTGTATAGCTGCAAATATTAATAAACTTTAACAAAAATCAGATCCATAAAA  
GAACCTGTGGCTTTGCTACAAGAAGTCTATAGAAATTTCTGTGACAGATCTTTCACCAACAGA  
TATAATTACATATATAGAAATATTAGCTGAATCATCTTCATTACTAGGTTACAAGAACAACA  
CTATCTCAGCCAAGGACACCCTTTCTAACTCAACTCTTACTGAATTTGTAAAAACCGTGAAT  
AATTTTGTTCAAAGGGATACATTTGTAGTTTGGGACAAGTTATCTGTGAATCATAGGAGAAC  
ACATCTTACAAAACCTCATGCACACTGTTGAACAAGCTACTTTAAGGATATCCCAGAGCTTCC  
AAAAGACCACAGAGTTTGATACAAATTCAACGGATATAGCTCTCAAAGTTTTCTTTTTTGAT  
TCATATAACATGAAACATATTCATCCTCATATGAATATGGATGGAGACTACATAAATATATT  
TCCAAAGAGAAAAGCTGCATATGATTCAAATGGCAATGTTGCAGTTGCATTTTTTATATTATA  
AGAGTATTGGTCCTTTGCTTTCATCATCTGACAACCTTCTTATTGAAACCTCAAATTTATGAT  
AATTCTGAAGAGGAGGAAAGAGTCATATCTTCAGTAATTTAGTCTCAATGAGCTCAAACCC  
ACCCACATTATATGAACCTGAAAAATAACATTTACATTAAGTCATCGAAAGGTCACAGATA  
GGTATAGGAGTCTATGTGCATTTTGGAAATTACTCACCTGATACCATGAATGGCAGCTGGTCT  
TCAGAGGGCTGTGAGCTGACATACTCAAATGAGACCCACACCTCATGCCGCTGTAATCACCT  
GACACATTTTGCAATTTTGATGTCCTCTGGTCCTTCCATTGGTATTAAAGATTATAATATTC  
TTACAAGGATCACTCAACTAGGAATAATTATTTCACTGATTTGTCTTGCCATATGCATTTTT  
ACCTTCTGGTCTTTCAGTGAAATTCAAAGCACCAGGACAACAATTCACAAAAATCTTTGCTG  
TAGCCTATTTCTTGCTGAACTTGTTTTCTTGTTGGGATCAATACAAATACTAATAAGCTCT  
TCTGTTCAATCATTGCCGACTGCTACACTACTTCTTTTAGCTGCTTTTGCATGGATGTGC  
ATTGAAGGCATACATCTCTATCTCATTGTTGTGGGTGTCATCTACAACAAGGGATTTTTGCA  
CAAGAATTTTTATATCTTTGGCTATCTAAGCCCAGCCGTGGTAGTTGGATTTTCGGCAGCAC  
TAGGATACAGATATTATGGCACAACCAAAGTATGTTGGCTTAGCACCGAAAACAACTTTATT  
TGGAGTTTTATAGGACCAGCATGCCTAATCATTCTTGTTAATCTCTGGCTTTTGGAGTCAT  
CATATACAAAGTTTTTCGTCACACTGCAGGGTTGAAACCAGAAGTTAGTTGCTTTGAGAACAA  
TAAGGTCTTGTCGAAGAGGAGCCCTCGCTCTTCTGTTCTTCTCGGCACCACCTGGATCTTT

FIGURE 33B

GGGGTTCTCCATGTTGTGCACGCATCAGTGGTTACAGCTTACCTCTTCACAGTCAGCAATGC  
TTTCCAGGGGATGTTTCATTTTTTTTATTCTGTGTGTTTTATCTAGAAAGATTCAAGAAGAAT  
ATTACAGATTGTTCAAAAATGTCCCCTGTTGTTTGGATGTTTAAGGTAACATAGAGAATG  
GTGGATAATTACAACCTGCACAAAATAAAAATTCCAAGCTGTGGATGACCAATGTATAAAAA  
TGACTCATCAAATTATCCAATTATTAATACTACTAGACAAAAGTATTTTAAATCAGTTTTTCT  
GTTTATGCTATAGGAACTGTAGATAATAAGGTAAAATTATGTATCATATAGATATACTATGT  
TTTTCTATGTGAAATAGTTCTGTCAAAAATAGTATTGCAGATATTTGGAAAGTAATTGGTTT  
CTCAGGAGTGATATCACTGCACCCAAGGAAAGATTTTCTTTCTAACACGAGAAGTATATGAA  
TGTCTGAAGGAAACCACTGGCTTGATATTTCTGTGACTCGTGTTGCCTTTGAAACTAGTCC  
CCTACCACCTCGGTAATGAGCTCCATTACAGAAAGTGGAACATAAGAGAATGAAGGGGCAGA  
ATATCAAACAGTGAAAAGGGAATGATAAGATGTATTTTGAATGAACTGTTTTTCTGTAGAC  
TAGCTGAGAAATTGTTGACATAAAATAAAGAATTGAAGAAACACATTTTACCATTTTGTGAA  
TTGTTCTGAACTTAAATGTCCACTAAAACAACCTTAGACTTCTGTTTGCTAAATCTGTTTCTT  
TTTCTAATATTCTAAAAAAAAAAAAAAAAAGGTTTACCTCCACAAATTGAAAAAAAAAAAAAAAA  
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

FIGURE 34

Signal peptide: Amino acids 1-19

Transmembrane domain: Amino acids 431-450;494-515;  
573-594;619-636;646-664

N-glycosylation sites: Amino acids 15-19;21-25;  
64-68;74-78;127-131;177-181;  
188-192;249-253;381-385;  
395-399

Glycosaminoglycan attachment site: Amino acids 49-53

cAMP- and cGMP-dependent protein kinase phosphorylation site:  
Amino acids 360-364

Tyrosine kinase phosphorylation sites:  
Amino acids 36-44;670-677

N-myristoylation sites: Amino acids 38-44;50-56;  
52-58;80-86;382-388;388-394;  
434-440;480-486;521-527

Aspartic acid and asparagine hydroxylation site:  
Amino acids 75-87

MKRLPLLVVFSTLLNCSYTONCTKTPCLPNAKCEIRNGIEACYCNMGFSGNGVTICEDDNEC  
GNLTQSCGENANCTNTEGSYYCMCVPGFRSSSNQDRFITNDGTVCIENVNANCHLDNVICIAA  
NINKTLTKIRSIKEPVALLQEVYRNSVTDLSPTDIITYIEILAESSLLGYKNNTISAKDTL  
SNSTLTEFVKTVNNFVQRDTFVWVDKLSVNHRRTHLTKLMHTVEQATLRISQSFQKTTEFDT  
NSTDIALKVFFFDSDYNMKHIHPHMNMDGDYINIFPKRKAAYDSNGNVAVAFLLYKSGIPLLS  
SSDNFLLKPQNYDNSEEEERVISVISVSMSSNPPTLYELEKITFTLSHRKVTDYRSLCAF  
WNYSPTMNGSWSSEGCETYSNETHTSCRCNHLTHFAILMSSGPSIGIKDYNILTRITQLG  
IISLICLAICIFTFWFFSEIQSTRTTIHKNLCCSLFLAELVFLVGINTNTNKLFCSTIAGL  
LHYFFLAFAWMCIEGIHLYLIVVGVIYNKGFLHKNFYIFGYLSPAVVVGFSAAALGYRYYGT  
TKVCWLSTENNFIWSFIGPACLIILVNLLAFGVIIYKVRHTAGLKPEVSCFENIRSCARGA  
LALLFLLGTTWIFGVLHVHASVVTAYLFTVSNAFQGMFIFLFLCVLSRKIQEEYYRLFKNV  
PCCFGCLR

FIGURE 35

CCCAGAAGTTCAAGGGCCCCCGGCCTCCTGCGCTCCTGCCGCCGGGACCCTCGACCTCCTCA  
GAGCAGCCGGCTGCCGCCCCGGGAAGATGGCGAGGAGGAGCCGCCACCGCCTCCTCCTGCTG  
CTGCTGCGCTACCTGGTGGTCGCCCTGGGCTATCATAAGGCCTATGGGTTTTCTGCCCCAAA  
AGACCAACAAGTAGTCACAGCAGTAGAGTACCAAGAGGCTATTTTAGCCTGCAAAACCCCAA  
AGAAGACTGTTTCCTCCAGATTAGAGTGGAAGAACTGGGTCGGAGTGTCTCCTTTGTCTAC  
TATCAACAGACTCTTCAAGGTGATTTTAAAAATCGAGCTGAGATGATAGATTTCAATATCCG  
GATCAAAAATGTGACAAGAAGTGATGCGGGGAAATATCGTTGTGAAGTTAGTGCCCCATCTG  
AGCAAGGCCAAAACCTGGAAGAGGATACAGTCACTCTGGAAGTATTAGTGGCTCCAGCAGTT  
CCATCATGTGAAGTACCCTCTTCTGCTCTGAGTGGAAGTGTGGTAGAGCTACGATGTCAAGA  
CAAAGAAGGGAATCCAGCTCCTGAATACACATGGTTTAAGGATGGCATCCGTTTGCTAGAAA  
ATCCCAGACTTGGCTCCCAAAGCACCAACAGCTCATAACAATGAATACAAAACTGGAACT  
CTGCAATTTAATACTGTTTCCAAACTGGACACTGGAGAATATTCCTGTGAAGCCCGCAATTC  
TGTTGGATATCGCAGGTGTCCTGGGAAACGAATGCAAGTAGATGATCTCAACATAAGTGGCA  
TCATAGCAGCCGTAGTAGTTGTGGCCTTAGTGATTTCCGTTTGTGGCCTTGGTGTATGCTAT  
GCTCAGAGGAAAGGCTACTTTTCAAAGAAACCTCCTTCCAGAAGAGTAATTCTTCATCTAA  
AGCCACGACAATGAGTGAAAATGTGCAGTGGCTCACGCCTGTAATCCCAGCACTTTGGAAGG  
CCGCGGCGGGCGGATCACGAGGTCAGGAGTTCTAGACCAGTCTGGCCAATATGGTGAAACCC  
CATCTCTACTAAAATACAAAAATTAGCTGGGCATGGTGGCATGTGCCTGCAGTTCCAGCTGC  
TTGGGAGACAGGAGAATCACTTGAACCCGGGAGGCGGAGGTTGCAGTGAGCTGAGATCACGC  
CACTGCAGTCCAGCCTGGGTAAACAGAGCAAGATTCCATCTCAAAAAATAAAATAAATAAATA  
AATAAATACTGGTTTTTACCTGTAGAATTCTTACAATAAATATAGCTTGATATTC

## FIGURE 36

Signal peptide:	Amino acids 1-20
Transmembrane domain:	Amino acids 235-254
N-glycosylation sites:	Amino acids 98-102;187-191; 236-240;277-281
N-myristoylation sites:	Amino acids 182-188;239-245; 255-261;257-263;305-311
Amidation site:	Amino acids 226-230

MARRSRHRLLLLLLRYLVVALGYHKAYGFSAPKDQQVVTAVEYQEAILACKTPKKTVSSRLE  
WKKLGRSVSFVYYQQTLQGDFKNRAEMIDFNIRIKNVTRSDAGKYRCEVSAPSEQQNLEED  
TVTLEVLVAPAVPSCEVPSSALSGTVVELRCQDKEGNPAPEYTWFKDGIRLLENPRLGSQST  
NSSYTMNTKTGTLQFNTVSKLDTGEYSCEARNVSGYRRCPGKRMQVDDLNIISGIIAAVVVVA  
LVISVCGLGVCYAQRKGYFSKETSFQKSNSSSKATTMSENVQWLTPVIPALWCAAAGGSRGQEF

FIGURE 37

GGAGCCGCCCTGGGTGTCAGCGGCTCGGCTCCCGCGCACGCTCCGGCCGTCGCGCAGCCTCG  
GCACCTGCAGGTCCGTGCGTCCCGCGGCTGGCGCCCCTGACTCCGTCCCGGCCAGGGAGGGC  
CATGATTTCCCTCCCGGGGCCCCCTGGTGACCAACTTGCTGCGGTTTTTGTTCCTGGGGCTGA  
GTGCCCTCGCGCCCCCCTCGCGGGCCCAGCTGCAACTGCACTTGCCCCGCAACCGGTTGCAG  
GCGGTGGAGGGAGGGGAAGTGGTGCTTCCAGCGTGGTACACCTTGACGGGGAGGTGTCTTC  
ATCCCAGCCATGGGAGGTGCCCTTTGTGATGTGGTTCTTCAAACAGAAAGAAAAGGAGGATC  
AGGTGTTGTCTACATCAATGGGGTCACAACAAGCAAACCTGGAGTATCCTTGGTCTACTCC  
ATGCCCTCCCGGAACCTGTCCCTGCGGCTGGAGGGTCTCCAGGAGAAAGACTCTGGCCCCCTA  
CAGCTGCTCCGTGAATGTGCAAGACAAACAAGGCAAATCTAGGGGCCACAGCATCAAAACCT  
TAGAACTCAATGTACTGGTTCCCTCCAGCTCCTCCATCCTGCCGTCTCCAGGGTGTGCCCCAT  
GTGGGGGCAAACGTGACCCTGAGCTGCCAGTCTCCAAGGAGTAAGCCCGCTGTCCAATACCA  
GTGGGATCGGCAGCTTCCATCCTTCCAGACTTTCTTTGCACCAGCATTAGATGTATCCGTG  
GGTCTTTAAGCCTCACCAACCTTTCGTCTTCCATGGCTGGAGTCTATGTCTGCAAGGGCCAC  
AATGAGGTGGGCACTGCCCAATGTAATGTGACGCTGGAAGTGAGCACAGGGCCTGGAGCTGC  
AGTGGTTGCTGGAGCTGTTGTGGGTACCCTGGTTGGACTGGGGTTGCTGGCTGGGCTGGTCC  
TCTTGTACCACCGCCGGGCAAGGCCCTGGAGGAGCCAGCCAATGATATCAAGGAGGATGCC  
ATTGCTCCCCGGAACCTGCCCTGGCCCAAGAGCTCAGACACAATCTCCAAGAATGGGACCCT  
TTCCTCTGTACCTCCGCACGAGCCCTCCGGCCACCCCATGGCCCTCCCAGGCCTGGTGCAT  
TGACCCCCACGCCCAGTCTCTCCAGCCAGGCCCTGCCCTCACCAAGACTGCCACGACAGAT  
GGGGCCCAACCTCAACCAATATCCCCCATCCCTGGTGGGGTTTCTTCCTCTGGCTTGAGCCG  
CATGGGTGCTGTGCCTGTGATGGTGCCTGCCAGAGTCAAGCTGGCTCTCTGGTATTGATGAC  
CCCACCACTCATTGGCTAAAGGATTTGGGGTCTCTCCTTCCTATAAGGGTCACCTCTAGCAC  
AGAGGCCTGAGTCATGGGAAAGAGTCACACTCCTGACCCTTAGTACTCTGCCCCACCTCTC  
TTTACTGTGGGAAAACCATCTCAGTAAGACCTAAGTGTCCAGGAGACAGAAGGAGAAGAGGA  
AGTGGATCTGGAATTGGGAGGAGCCTCCACCCACCCCTGACTCCTCCTTATGAAGCCAGCTG  
CTGAAATTAGCTACTACCAAGAGTGAGGGGCAGAGACTTCCAGTCACTGAGTCTCCCAGGC  
CCCCTTGATCTGTACCCACCCCTATCTAACACCACCCCTTGGCTCCCACTCCAGCTCCCTGT  
ATTGATATAACCTGTCAGGCTGGCTTGGTTAGGTTTTACTGGGGCAGAGGATAGGGAATCTC  
TTATTAAACTAACATGAAATATGTGTTGTTTTCATTTGCAAATTTAAATAAAGATACATAA  
TGTTTGTATGAAAAA



FIGURE 38

Signal peptide:	Amino acids 1-29
Transmembrane domain:	Amino acids 244-265
N-glycosylation sites:	Amino acids 108-112;169-173; 213-217;236-240;307-311
N-myristoylation sites:	Amino acids 90-96;167-173; 220-226;231-237;252-258; 256-262;262-268;308-314; 363-369;364-370
Prokaryotic membrane lipoprotein lipid attachment site:	Amino acids 164-175

MISLPGPLVTNLLRFLFLGLSALAPPSRAQLQLHLPANRLQAVEGGEVVLPAWYTLHGEVSS  
SQPWEVPFVMWFFKQKEKEDQVLSYINGVTTSKPGVSLVYSMPSRNLSLRLEGLQEKDSGPY  
SCSVNVQDKQGKSRGHSIKTLELNLVPPAPPSCRLQGVPHVGANVTLSQCSPRSKPAVQYQ  
WDRQLPSFQTFFAPALDVIRGSLSLTNLSSSMAGVYVCKAHNEVGTAQCNVTLEVSTGPGAA  
VVAGAVVGTLVGLGLLAGLVLLYHRRGKALEEPANDIKEDAIAPRTLPPWKSSDTISKNGTL  
SSVTSARALRPPHGPPRPGALTPTPSLSSQALPSPRLPTTDGAHPQPISPIPGGVSSSGLSR  
MGAVPVMVPAQSQAGSLV

FIGURE 39

CCCACGCGTCCGCACCTCGGCCCCGGGCTCCGAAGCGGCTCGGGGGCGCCCTTTCGGTCAAC  
ATCGTAGTCCACCCCCTCCCCATCCCCAGCCCCCGGGATTTCAGGCTCGCCAGCGCCAGCC  
AGGGAGCCGGCCGGGAAGCGCGATGGGGGCCCCAGCCGCCTCGCTCCTGCTCCTGCTCCTGC  
TGTTTCGCTGCTGCTGGGCGCCCGGGCGGGGCCAACCTCTCCAGGACGACAGCCAGCCCTGG  
ACATCTGATGAAACAGTGGTGGCTGGTGGCACCGTGGTGCTCAAGTGCCAAGTGAAAGATCA  
CGAGGACTCATCCCTGCAATGGTCTAACCTGCTCAGCAGACTCTCTACTTTGGGGAGAAGA  
GAGCCCTTCGAGATAATCGAATTTCAGCTGGTTACCTCTACGCCCCACGAGCTCAGCATCAGC  
ATCAGCAATGTGGCCCTGGCAGACGAGGGCGAGTACACCTGCTCAATCTTCACTATGCCTGT  
GCGAACTGCCAAGTCCCTCGTCACTGTGCTAGGAATTCCACAGAAGCCCATCATCACTGGTT  
ATAAATCTTCATTACGGGAAAAAGACACAGCCACCCTAAACTGTCAGTCTTCTGGGAGCAAG  
CCTGCAGCCCGGCTCACCTGGAGAAAGGGTGACCAAGAACTCCACGGAGAACCAACCCGCAT  
ACAGGAAGATCCCAATGGTAAAACCTTCACTGTCAGCAGCTCGGTGACATTCCAGGTTACCC  
GGGAGGATGATGGGGCGAGCATCGTGTGCTCTGTGAACCATGAATCTCTAAAGGGAGCTGAC  
AGATCCACCTCTCAACGCATTGAAGTTTATACACACCAACTGCGATGATTAGGCCAGACCC  
TCCCCATCCTCGTGAGGGCCAGAAGCTGTTGCTACACTGTGAGGGTCGCGGCAATCCAGTCC  
CCCAGCAGTACCTATGGGAGAAGGAGGGCAGTGTGCCACCCCTGAAGATGACCCAGGAGAGT  
GCCCTGATCTTCCCTTTCTCAACAAGAGTGACAGTGGCACCTACGGCTGCACAGCCACCAG  
CAACATGGGCAGCTACAAGGCCTACTACACCCTCAATGTTAATGACCCAGTCCGGTGCCCT  
CCTCCTCCAGCACCTACCACGCCATCATCGGTGGGATCGTGGCTTTCATTGTCTTCCTGCTG  
CTCATCATGCTCATCTTCCTTGGCCACTACTTGATCCGGCACAAAGGAACCTACCTGACACA  
TGAGGCAAAAGGCTCCGACGATGCTCCAGACGCGGACACGGCCATCATCAATGCAGAAGGCG  
GGCAGTCAGGAGGGGACGACAAGAAGGAATATTTTCATCTAGAGGCGCCTGCCCACTTCCTGC  
GGGGGGCAGGGGCCCTGTGGGGACTGCTGGGGCCGTACCAACCCGGACTTGTACAGAGCAA  
CCGCAGGGCCGCCCTCCCGCTTGCTCCCCAGCCCCACCCACCCCTGTACAGAATGTCTGC  
TTTGGGTGCGGTTTTGTACTCGGTTTGGAATGGGGAGGGAGGAGGGCGGGGGAGGGGAGGG  
TTGCCCTCAGCCCTTTCGTGGCTTCTCTGCATTTGGGTTATTATTATTTTTGTAAACAATCC  
CAAATCAAATCTGTCTCCAGGCTGGAGAGGCAGGAGCCCTGGGGTGAGAAAAGCAAAAAACA  
AACAAAAACA

FIGURE 40

Signal peptide:	Amino acids 1-20
Transmembrane domain:	Amino acids 331-352
N-glycosylation sites:	Amino acids 25-29;290-294
N-myristoylation sites:	Amino acids 2-8;23-29; 156-162;218-224;295-301; 298-304;306-312;334-340; 360-366;385-391;386-392
Prokaryotic membrane lipoprotein lipid attachment site:	Amino acids 7-18

MGAPAASLLLLLLLLFACCWAPGGANLSQDDSQPWTSDETVVAGGTVVLKCQVKDHEDSSLQW  
SNPAQQTLYFGEKRALRDNRIQLVTSTPHELISISNVALADEGEYTCISFTMPVRTAKSLV  
TVLGIPQKPIITGYKSSLREKDTATLNCQSSGSKPAARLTWRKGDQELHGEPTRIQEDPNGK  
TFTVSSSVTFQVTREDDGASIVCSVNHESLKGADRSTSQRIEVLYTPTAMIRPDPPHPREGQ  
KLLHCEGRGNPVPQQYLWEKEGSVPPLKMTQESALIFPFLNKSDSGTYGCTATSNMGSYKA  
YYTLNVNDPSPVPSSSSTYHAIIGGIVAFIVFLLLIIMLIFLGHYLIRHKGTYLTHEAKGSDD  
APDADTAIINAEGGQSGGDDKKEYFI

FIGURE 41

GCTGGGGACATGAGAGGCACACCGAAGACCCACCTCCTGGCCTTCTCCCTCCTCTGCCTCCT  
CTCAAAGGTGCGTACCCAGCTGTGCCCCGACACCATGTACCTGCCCCCTGGCCACCTCCCCGAT  
GCCCCGCTGGGAGTACCCCTGGTGCTGGATGGCTGTGGCTGCTGCCGGGTATGTGCACGGCGG  
CTGGGGGAGCCCTGCGACCAACTCCACGTCTGCGACGCCAGCCAGGGCCTGGTCTGCCAGCC  
CGGGGCAGGACCCGGTGGCCGGGGGGCCCTGTGCCTCTTGCCAGAGGACGACAGCAGCTGTG  
AGGTGAACGGCCGCCTGTATCGGGAAGGGGAGACCTTCCAGCCCCACTGCAGCATCCGCTGC  
CGCTGCGAGGACGGCGGCTTCACCTGCGTGCCGCTGTGCAGCGAGGATGTGCGGCTGCCAG  
CTGGGACTGCCCCCAGGAGGGTTCGAGGTCCTGGGCAAGTGCTGCCCTGAGTGGGTGT  
GCGGCCAAGGAGGGGGACTGGGGACCCAGCCCCCTTCCAGCCCAAGGACCCAGTTTTCTGGC  
CTTGTCTCTTCCCTGCCCCCTGGTGTCCCCTGCCCAGAATGGAGCACGGCCTGGGGACCCTG  
CTCGACCACCTGTGGGCTGGGCATGGCCACCCGGGTGTCCAACCAGAACCCTTCTGCCGAC  
TGGAGACCCAGCGCCGCCTGTGCCTGTCCAGGCCCTGCCACCCTCCAGGGGTGCGAGTCCA  
CAAAACAGTGCCTTCTTAGAGCCGGGCTGGGAATGGGGACACGGTGTCCACCATCCCCAGCTG  
GTGGCCCTGTGCCTGGGCCCTGGGCTGATGGAAGATGGTCCGTGCCCAGGCCCTTGGCTGCA  
GGCAACACTTTAGCTTGGGTCCACCATGCAGAACACCAATATTAACACGCTGCCTGGTCTGT  
CTGGATCCCCAGGTATGGCAGAGGTGCAAGACCTAGTCCCCTTTCTCTAACTCACTGCCTA  
GGAGGCTGGCCAAGGTGTCCAGGGTCCTCTAGCCCACTCCCTGCCTACACACACAGCCTATA  
TCAAACATGCACACGGGCGAGCTTTCTCTCCGACTTCCCCTGGGCAAGAGATGGGACAAGCA  
GTCCCTTAATATTGAGGCTGCAGCAGGTGCTGGGCTGGACTGGCCATTTTTCTGGGGGTAGG  
ATGAAGAGAAGGCACACAGAGATTCTGGATCTCCTGCTGCCTTTTCTGGAGTTTGTAAAATT  
GTTCTGAATACAAGCCTATGCGTGA

## FIGURE 42

Signal peptide:	Amino acids 1-23
N-myristoylation sites:	Amino acids 3-9;49-55;81-87; 85-91;126-132;164-170; 166-172;167-173;183-189; 209-215
Insulin-like growth factor binding proteins signature:	Amino acids 49-65
von Willebrand C1 domain:	Amino acids 107-124
Thrombospondin 1:	Amino acids 201-216
IGF binding protein:	Amino acids 49-58

MRGTPKTHLLAFSLLCLLSKVRTQLCPTPCTCPWPPPRCPLGVPLVLDGCGCCRVCCARRLGE  
PCDQLHVCDASQGLVCQPGAGPGGRGALCLLAEDDSSCEVNGRLYREGETFQPHCSIRCRCE  
DGGFTCVPLCSEDVRLPSWDCPHPRRVEVLGKCCPEWVCGQGGGLGTQPLPAQGPQFSGLVLS  
SLPPGVPCPEWSTAWGPCSTTCGLGMATRVSNQNRFCRLETQRRRLCLSRPCPPSRGRSPQNSAF

FIGURE 43

GGAGAGCGGAGCGAAGCTGGATAACAGGGGACCGATGATGTGGCGACCATCAGTTCTGCTGC  
TTCTGTTGCTACTGAGGCACGGGGCCAGGGGAAGCCATCCCCAGACGCAGGCCCTCATGGC  
CAGGGGAGGGTGCACCAGGCGGGCCCCCTGAGCGACGCTCCCCATGATGACGCCACGGGAA  
CTTCCAGTACGACCATGAGGCTTTCCTGGGACGGGAAGTGGCCAAGGAATTCGACCAACTCA  
CCCCAGAGGAAAGCCAGGCCCGTCTGGGGCGGATCGTGGACCGCATGGACCGCGCGGGGGAC  
GGCGACGGCTGGGTGTGCTGGCCGAGCTTCGCGCGTGGATCGCGCACACGCAGCAGCGGCA  
CATACGGGACTCGGTGAGCGCGGCCTGGGACACGTACGACACGGACCGCGACGGGCGTGTGG  
GTTGGGAGGAGCTGCGCAACGCCACCTATGGCCACTACGCGCCCGGTGAAGAATTTTCATGAC  
GTGGAGGATGCAGAGACCTACAAAAGATGCTGGCTCGGGACGAGCGGCGTTTCCGGGTGGC  
CGACCAGGATGGGGACTCGATGGCCACTCGAGAGGAGCTGACAGCCTTCCTGCACCCCGAGG  
AGTTCCCTCACATGCGGGACATCGTGATTGCTGAAACCCTGGAGGACCTGGACAGAAACAAA  
GATGGCTATGTCCAGGTGGAGGAGTACATCGCGGATCTGTACTCAGCCGAGCCTGGGGAGGA  
GGAGCCGGCGTGGGTGCAGACGGAGAGGCAGCAGTTCCGGGACTTCCGGGATCTGAACAAGG  
ATGGGCACCTGGATGGGAGTGAGGTGGGCCACTGGGTGCTGCCCCCTGCCCAGGACCAGCCC  
CTGGTGAAGCCAACCACCTGCTGCACGAGAGCGACACGGACAAGGATGGGCGGCTGAGCAA  
AGCGGAAATCCTGGGTAATTGGAACATGTTTGTGGGCAGTCAGGCCACCAACTATGGCGAGG  
ACCTGACCCGGCACACGATGAGCTGTGAGCACCGCGCACCTGCCACAGCCTCAGAGGCCCG  
CACAATGACCGGAGGAGGGCCGCTGTGGTCTGGCCCCCTCCCTGTCCAGGCCCGCAGGAG  
GCAGATGCAGTCCCAGGCATCCTCCTGCCCCTGGGCTCTCAGGGACCCCTGGGTGGCTTC  
TGTCCCTGTCACACCCCAACCCAGGGAGGGGCTGTATAGTCCCAGAGGATAAGCAATAC  
CTATTTCTGACTGAGTCTCCCAGCCCAGACCCAGGGACCCTTGCCCCAAGCTCAGCTCTAA  
GAACCGCCCCAACCCCTCCAGCTCCAAATCTGAGCCTCCACCACATAGACTGAAACTCCCT  
GGCCCCAGCCCTCTCCTGCCTGGCCTGGCCTGGGACACCTCCTCTCTGCCAGGAGGCAATAA  
AAGCCAGCGCCGGGACCTTGAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA  
AAAAAAAAAAAAAA

## FIGURE 44

Signal peptide:	Amino acids 1-20
N-glycosylation site:	Amino acids 140-144
N-myristoylation sites:	Amino acids 263-269;311-317
Endoplasmic reticulum targeting sequence:	Amino acids 325-330

MMWRPSVLLLLLLLLRHGAQGKPSPDAGPHGQGRVHQAAPLSDA PHDDAHGNFQYDHEAFLGR  
EVAKEFDQLTPEESQARLGRIVDRMDRAGDGDGWVSLAELRAWIAHTQQRHIRDSVSAAWDT  
YDTRDGRVGVWEELRNATYGHYAPGEEFHDVEDAETYKKMLARDERRFRVADQDGDSMATRE  
ELTAFLHPEEFPHMRDIVIAETLEDLDRNKDGYVQVEEYIADLYSAEPGEEEPAWVQTERQQ  
FRDFRDLNKDGHLDGSEVGHVLP PAQDQPLVEANHLLHESDTDKDGRLSKAEILGNWNMFV  
GSQATNYGEDLTRHHDEL

FIGURE 45

GTCTGTTCCAGGAGTCCTTCGGCGGCTGTTGTGTCAGTGGCCTGATCGCGATGGGGACAAA  
GGCGCAAGTCGAGAGGAACTGTTGTGCCTCTTCATATTGGCGATCCTGTTGTGCTCCCTGG  
CATTGGGCAGTGTTACAGTGCACCTCTTCTGAACCTGAAGTCAGAATTCCTGAGAATAATCCT  
GTGAAGTTGTCCTGTGCCTACTCGGGCTTTTCTTCTCCCCGTGTGGAGTGGAAGTTTGACCA  
AGGAGACACCACCAGACTCGTTTGTCTATAATAACAAGATCACAGCTTCCTATGAGGACCGGG  
TGACCTTCTTGCCAACTGGTATCACCTTCAAGTCCGTGACACGGGAAGACACTGGGACATAC  
ACTTGTATGGTCTCTGAGGAAGGCGGCAACAGCTATGGGGAGGTCAAGGTCAAGCTCATCGT  
GCTTGTGCCTCCATCCAAGCCTACAGTTAACATCCCCCTCCTCTGCCACCATTGGGAACCGGG  
CAGTGCTGACATGCTCAGAACAAAGATGGTTCCCCACCTTCTGAATACACCTGGTTCAAAGAT  
GGGATAGTGATGCCTACGAATCCCAAAAGCACCCGTGCCTTCAGCAACTCTTCCTATGTCTCT  
GAATCCCACAACAGGAGAGCTGGTCTTTGATCCCCTGTCAGCCTCTGATACTGGAGAATACA  
GCTGTGAGGCACGGAATGGGTATGGGACACCCATGACTTCAAATGCTGTGCGCATGGAAGCT  
GTGGAGCGGAATGTGGGGGTGTCGTGGCAGCCGTCCTTGTAACCCTGATTCTCCTGGGAAT  
CTTGGTTTTTTGGCATCTGGTTTGCCTATAGCCGAGGCCACTTTGACAGAACAAAGAAAGGGA  
CTTCGAGTAAGAAGGTGATTTACAGCCAGCCTAGTGCCCGAAGTGAAGGAGAATTCAAACAG  
ACCTCGTCATTCTGGTGTGAGCCTGGTTCGGCTCACCGCCTATCATCTGCATTTGCCTTACT  
CAGGTGCTACCGGACTCTGGCCCCCTGATGTCTGTAGTTTACAGGATGCCTTATTTGTCTTC  
TACACCCACAGGGCCCCCTACTTCTTCGGATGTGTTTTTAATAATGTCAGCTATGTGCCCC  
ATCCTCCTTCATGCCCTCCCTCCCTTTCTACCACTGCTGAGTGGCCTGGAACCTGTTTAAA  
GTGTTTATTCCCCATTTCTTTGAGGGATCAGGAAGGAATCCTGGGTATGCCATTGACTTCCC  
TTCTAAGTAGACAGCAAAAATGGCGGGGGTTCGAGGAATCTGCACTCAACTGCCCACCTGGC  
TGGCAGGGATCTTTGAATAGGTATCTTGAGCTTGGTTCTGGGCTCTTTCCTTGTGTACTGAC  
GACCAGGGCCAGCTGTTCTAGAGCGGGAATTAGAGGCTAGAGCGGCTGAAATGGTTGTTTGG  
TGATGACACTGGGGTCCTTCCATCTCTGGGGCCCACTCTCTTCTGTCTTCCCATGGGAAGTG  
CCACTGGGATCCCTCTGCCCTGTCCCTCCTGAATACAAGCTGACTGACATTGACTGTGTCTGT  
GGAAAATGGGAGCTCTTGTTGTGGAGAGCATAGTAAATTTTCAGAGAACTTGAAGCCAAAAG  
GATTTAAAACCGCTGCTCTAAAGAAAAGAAAAGCTGGAGGCTGGGCGCAGTGGCTCACGCCTG  
TAATCCCAGAGGCTGAGGCAGGCGGATCACCTGAGGTGCGGAGTTTCGGGATCAGCCTGACCA  
ACATGGAGAAACCTACTGGAAATACAAAGTTAGCCAGGCATGGTGGTGCATGCCTGTAGTC  
CCAGCTGCTCAGGAGCCTGGCAACAAGAGCAAACTCCAGCTCAAAAAAAAAAAAAAAAAA



## FIGURE 46

Signal peptide:	Amino acids 1-27
Transmembrane domain:	Amino acids 235-256
N-glycosylation site:	Amino acids 185-189
cAMP- and cGMP-dependent protein kinase phosphorylation site:	Amino acids 270-274
N-myristoylation sites:	Amino acids 105-111;116-122; 158-164;219-225;237-243; 256-262

MGTKAQVERKLLCLFILAILLCSLALGSVTVHSSEPEVRIPENNPVKLSCAYSGFSSPRVEW  
KFDQGD TTRLVCYNNKITASYEDRVTF LPTGITFKSVTREDTGTYTCMVSEEGGNSYGEVKV  
KLIVLVPPSKPTVNIPSSATIGNRAVLTCSEQDGSPPEYTWFKDGI VMPTNPKSTRAFSNS  
SYVLNPTTGELVFDPLSASDTGEYSCEARNGYGT PMT SNAVRMEAVERNVGVIVA AVLVTLI  
LLGILVFGIWFAYS RGHFDRTKKGTSSKKVIYSQPSARSEGEFKQTSSFLV

FIGURE 47

CCCAGGCTCTAGTGCAGGAGGAGAAGGAGGAGGAGCAGGAGGTGGAGATTCCCAGTTAAAAG  
GCTCCAGAATCGTGTACCAGGCAGAGAACTGAAGTACTGGGGCCTCCTCCACTGGGTCCGAA  
TCAGTAGGTGACCCCGCCCCTGGATTCTGGAAGACCTCACCATGGGACGCCCCGACCTCGT  
GCGGCCAAGACGTGGATGTTCTTGCTCTTGCTGGGGGAGCCTGGGCAGGACACTCCAGGGC  
ACAGGAGGACAAGGTGCTGGGGGGTCATGAGTGCCAACCCCATTCGCAGCCTTGGCAGGCGG  
CCTTGTTCCAGGGCCAGCAACTACTCTGTGGCGGTGTCCTTGTAGGTGGCAACTGGGTCTTT  
ACAGCTGCCCCACTGTAAAAAACCGAAATACACAGTACGCCTGGGAGACCACAGCCTACAGAA  
TAAAGATGGCCCAGAGCAAGAAATACCTGTGGTTCAGTCCATCCCACACCCCTGCTACAACA  
GCAGCGATGTGGAGGACCACAACCATGATCTGATGCTTCTTCAACTGCGTGACCAGGCATCC  
CTGGGGTCCAAAGTGAAGCCCATCAGCCTGGCAGATCATTGCACCCAGCCTGGCCAGAAGTG  
CACCGTCTCAGGCTGGGGCACTGTCAACAGTCCCCGAGAGAATTTTCCTGACACTCTCAACT  
GTGCAGAAGTAAAAATCTTTCCCCAGAAGAAGTGTGAGGATGCTTACCCGGGGCAGATCACA  
GATGGCATGGTCTGTGCAGGCAGCAGCAAAGGGGCTGACACGTGCCAGGGCGATTCTGGAGG  
CCCCCTGGTGTGTGATGGTGCCTCCAGGGCATCACATCCTGGGGCTCAGACCCCTGTGGGA  
GGTCCGACAAACCTGGCGTCTATACCAACATCTGCCGCTACCTGGACTGGATCAAGAAGATC  
ATAGGCAGCAAGGGCTTGATTCTAGGATAAGCACTAGATCTCCCTTAATAAACTCACAACTCT  
CTGGTTC

## FIGURE 48

Signal peptide: Amino acids 1-23  
Transmembrane domain: Amino acids 51-71  
Glycosaminoglycan attachment site: Amino acids 155-159  
Tyrosine kinase phosphorylation site:  
Amino acids 182-189  
N-myristoylation sites: Amino acids 19-25;20-26;  
35-41;60-66;149-155;195-201;  
200-206;204-210;220-226;  
229-235;240-246  
N-glycosylation site: Amino acid 110-114  
Serine proteases, trypsin  
family and histidine active site: Amino acids 69-75  
Consensus sequence: Amino acids 207-217  
Kringle domain protein motif: Amino acids 205-217

MGRPRPRAAKTWMFLLLLGGAWAGHSRAQEDKVLGGHECQPHSQPWQAALFQGQQLLCGGVL  
VGGNWVLTAAHCKKPKYTVRLGDHSLQNKDGPEQEIPVVQSI PHPCYNSSDVEDHNHDLMLL  
QLRDQASLGSKVKPISLADHCTQPGQKCTVSGWGTVTSPRENFPDTLNCAEVKIFPQKKCED  
AYPGQITDGMVCAGSSKGADTCQGDSSGGPLVCDGALQGITSWGSDPCGRSDKPGVYTNICRY  
LDWIKKIIGSKG

FIGURE 49

GACGGCTGGCCACCATGCACGGCTCCTGCAGTTTCCTGATGCTTCTGCTGCCGCTACTGCTA  
CTGCTGGTGGCCACCACAGGCCCCGTTGGAGCCCTCACAGATGAGGAGAAACGTTTGATGGT  
GGAGCTGCACAACCTCTACCGGGCCCAGGTATCCCCGACGGCCTCAGACATGCTGCACATGA  
GATGGGACGAGGAGCTGGCCGCCTTCGCCAAGGCCTACGCACGGCAGTGCGTGTGGGGCCAC  
AACAAGGAGCGCGGGCGCCGCGGCGAGAATCTGTTCCGCATCACAGACGAGGGCATGGACGT  
GCCGCTGGCCATGGAGGAGTGGCACCACGAGCGTGAGCACTACAACCTCAGCGCCGCCACCT  
GCAGCCCAGGCCAGATGTGCGGCCACTACACGCAGGTGGTATGGGCCAAGACAGAGAGGATC  
GGCTGTGGTTCCCACTTCTGTGAGAAGCTCCAGGGTGTTGAGGAGACCAACATCGAATTACT  
GGTGTGCAACTATGAGCCTCCGGGGAACGTGAAGGGGAAACGGCCCTACCAGGAGGGGACTC  
CGTGCTCCCAATGTCCCTCTGGCTACCACTGCAAGAACTCCCTCTGTGAACCCATCGGAAGC  
CCGGAAGATGCTCAGGATTTGCCTTACCTGGTAACTGAGGCCCCATCCTTCCGGGCGACTGA  
AGCATCAGACTCTAGGAAAATGGGTACTCCTTCTTCCCTAGCAACGGGGATTCCGGCTTTCT  
TGTTAACAGAGGTCTCAGGCTCCCTGGCAACCAAGGCTCTGCCTGCTGTGGAAACCCAGGCC  
CCAACCTTCCCTAGCAACGAAAGACCCGCCCTCCATGGCAACAGAGGCTCCACCTTGCCTAAC  
AACTGAGGTCCCTTCCATTTTGGCAGCTCACAGCCTGCCCTCCTTGGATGAGGAGCCAGTTA  
CCTTCCCCAAATCGACCCATGTTTCTATCCCAAATCAGCAGACAAAGTGACAGACAAAACA  
AAAGTGCCCTCTAGGAGCCCAGAGAACTCTCTGGACCCCAAGATGTCCCTGACAGGGGCAAG  
GGAACCTCTACCCCATGCCCAGGAGGAGGCTGAGGCTGAGGCTGAGTTGCCTCCTTCCAGTG  
AGGTCTTGGCCTCAGTTTTTCCAGCCCAGGACAAGCCAGGTGAGCTGCAGGCCACACTGGAC  
CACACGGGGCACACCTCCTCCAAGTCCCTGCCCAATTTCCCAATACCTCTGCCACCGCTAA  
TGCCACGGGTGGGCGTGCCCTGGCTCTGCAGTCGTCTTGCCAGGTGCAGAGGGCCCTGACA  
AGCCTAGCGTTGTGTCAGGGCTGAACTCGGGCCCTGGTCATGTGTGGGGCCCTCTCCTGGGA  
CTACTGCTCCTGCCTCCTCTGGTGTGGCTGGAATCTTCTTGAATGGGATACCACTCAAAGGG  
TGAAGAGGTGAGCTGTCCTCCTGTCATCTTCCCCACCCTGTCCCCAGCCCCTAAACAAGATA  
CTTCTTGGTTAAGGCCCTCCGGAAGGGAAAGGCTACGGGGCATGTGCCTCATCACACCATCC  
ATCCTGGAGGCACAAGGCCTGGCTGGCTGCGAGCTCAGGAGGCCGCCTGAGGACTGCACACC  
GGGCCCACACCTCTCCTGCCCTCCCTCCTGAGTCCTGGGGGTGGGAGGATTTGAGGGAGCT  
CACTGCCTACCTGGCCTGGGGCTGTCTGCCCACACAGCATGTGCGCTCTCCCTGAGTGCCCTG  
TGTAAGCTGGGGATGGGGATTCTAGGGGCAGATGAAGGACAAGCCCCACTGGAGTGGGGTTT  
TTTGAGTGGGGGAGGCAGGGACGAGGGAAGGAAAGTAACTCCTGACTCTCCAATAAAAACCT  
GTCCAACCTGTGAAA

## FIGURE 50

Signal peptide: Amino acids 1-22

N-glycosylation sites: Amino acids 114-118;403-407;  
409-413

Glycosaminoglycan attachment site: Amino acids 439-443

N-myristoylation sites: Amino acids 123-129;143-149;  
152-158;169-175;180-186;  
231-237;250-256

Amidation sites: Amino acids 82-86;172-176

Peroxidase proximal heme-ligand signature:  
Amino acids 287-298

Extracellular protein SCP/Tpx-1/Ag5/PR-1/Sc7 signature 1:  
Amino acids 127-138

Extracellular protein SCP/Tpx-1/Ag5/PR-1/Sc7 signature 2:  
Amino acids 160-172

MHGSCSFLMLLLPLLLLLLVATTGPVGALTDEEKRLMVELHNLRYAQVSPTASDMLHMRWDEE  
LAAFAKAYARQCVWGHNKERGRGENLFAITDEGMDVPLAMEEWHHEREHYNLSAATCSPGQ  
MCGHYTQVWAKTERIGCGSHFCEKLQGVETNIELLVCNYEPPGNVKGKRPYQEGTPCSQC  
PSGYHCKNSLCEPIGSPEDAQDLPYLVTEAPSF RATEASDSRKMGT PSSLATGIPAFLVTEV  
SGSLATKALPAVETQAPTSLATKDPPSMATEAPPCVTTEVPSILAAHSLPSLDEEPVTFPKS  
THVPIPKSADKVTDKTKVPSRSPENSLDPKMSLTGARELLPHAQEEAEAEELPPSSEVLAS  
VFPAQDKPGELQATLDHTGHTSSKSLPNFPNTSATANATGGRALALQSSLPGAEGPDKPSVV  
SGLNSGPGHVWGPLLGLLLLPLVLGIF

FIGURE 51A

GGGGAGAGGAATTGACCATGTAAAAGGAGACTTTTTTTTTTGGTGGTGGTGGCTGTTGGGTG  
CCTTGCAAAAATGAAGGATGCAGGACGCAGCTTCTCCTGGAACCGAACGCAATGGATAAAC  
TGATTGTGCAAGAGAGAAGGAAGAACGAAGCTTTTTCTTGTGAGCCCTGGATCTTAACACAA  
ATGTGTATATGTGCACACAGGGAGCATTCAAGAATGAAATAAACAGAGTTAGACCCGCGGG  
GGTTGGTGTGTTCTGACATAAATAAATAATCTTAAAGCAGCTGTTCCCCCTCCCCACCCCCAA  
AAAAAAGGATGATTGGAAATGAAGAACCGAGGATTCACAAAGAAAAAAGTATGTTCATTTTT  
CTCTATAAAGGAGAAAAGTGAGCCAAGGAGATATTTTTGGAATGAAAAGTTTGGGGCTTTTTT  
AGTAAAGTAAAGAACTGGTGTGGTGGTGTTCCTTTCTTTTGAATTTCCCACAAGAGGAG  
AGGAAATTAATAATACATCTGCAAAGAAATTTAGAGAGAAAAGTTGACCGCGGCAGATTG  
AGGCATTGATTGGGGGAGAGAAACCAGCAGAGCACAGTTGGATTTGTGCCTATGTTGACTAA  
AATTGACGGATAATTGCAGTTGGATTTTTCTTCATCAACCTCCTTTTTTTTAAATTTTTATT  
CCTTTTGGTATCAAGATCATGCGTTTTCTCTTGTTCCTTAACCACCTGGATTTCCATCTGGAT  
GTTGCTGTGATCAGTCTGAAATACAACGTGTTTGAATTCAGAAAGGACCAACACCAGATAAAT  
TATGAATGTTGAACAAGATGACCTTACATCCACAGCAGATAATGATAGGTCTAGGTTTAAAC  
AGGGCCCTATTTGACCCCCTGCTTGTGGTGTGCTGGCTCTTCAACTTCTTGTGGTGGCTGG  
TCTGGTGCGGGCTCAGACCTGCCCTTCTGTGTGCTCCTGCAGCAACCAGTTCAGCAAGGTGA  
TTTGTGTTTCGAAAAACCTGCGTGAGGTTCCGGATGGCATCTCCACCAACACACGGCTGCTG  
AACCTCCATGAGAACCAAATCCAGATCATCAAAGTGAACAGCTTCAAGCACTTGAGGCACCTT  
GGAAATCCTACAGTTGAGTAGGAACCATATCAGAACCATTGAAATTGGGGCTTTCAATGGTC  
TGGCGAACCTCAACACTCTGGAACCTTTTGACAATCGTCTTACTACCATCCCGAATGGAGCT  
TTTGTATACTTGTCTAAACTGAAGGAGCTCTGGTTGCGAAACAACCCCATTGAAAGCATCCC  
TTCTTATGCTTTTAAACAGAATTCCTTCTTTGCGCCGACTAGACTTAGGGGAATTGAAAAGAC  
TTTCATACATCTCAGAAGGTGCCTTTGAAGGTCTGTCCAACCTTGAGGTATTTGAACCTTGCC  
ATGTGCAACCTTCGGGAAATCCCTAACCTCACACCGCTCATAAACTAGATGAGCTGGATCT  
TTCTGGGAATCATTTATCTGCCATCAGGCCTGGCTCTTCCAGGGTTTGATGCACCTTCAAA  
AACTGTGGATGATACAGTCCCAGATTCAAGTGATTGAACGGAATGCCTTTGACAACCTTCAG  
TCACTAGTGGAGATCAACCTGGCACACAATAATCTAACATTACTGCCTCATGACCTCTTCAC  
TCCCTTGATCATCTAGAGCGGATACATTTACATCACAACCCTTGGAACGTAACTGTGACA  
TACTGTGGCTCAGCTGGTGGATAAAAGACATGGCCCCCTCGAACACAGCTTGTGTGCCCGG  
TGTAACACTCCTCCCAATCTAAAGGGGAGGTACATTGGAGAGCTCGACCAGAATTACTTCAC  
ATGCTATGCTCCGGTGATTGTGGAGCCCCCTGCAGACCTCAATGTCACTGAAGGCATGGCAG

FIGURE 51B

CTGAGCTGAAATGTCGGGCCTCCACATCCCTGACATCTGTATCTTGGATTACTCCAAATGGA  
ACAGTCATGACACATGGGGCGTACAAAGTGCGGATAGCTGTGCTCAGTGATGGTACGTTAAA  
TTTCACAAATGTAAGTGTGCAAGATACAGGCATGTACACATGTATGGTGAGTAATTCCGTTG  
GGAATACTACTGCTTCAGCCACCCTGAATGTTACTGCAGCAACCACTACTCCTTTCTCTTAC  
TTTTCAACCGTCACAGTAGAGACTATGGAACCGTCTCAGGATGAGGCACGGACCACAGATAA  
CAATGTGGGTCCCCTCCAGTGGTTCGACTGGGAGACCACCAATGTGACCACCTCTCTCACAC  
CACAGAGCACAAGGTCGACAGAGAAAACCTTCACCATCCCAGTGACTGATATAAACAGTGGG  
ATCCCAGGAATTGATGAGGTCATGAAGACTACCAAATCATCATTGGGTGTTTTGTGGCCAT  
CACACTCATGGCTGCAGTGATGCTGGTCATTTTCTACAAGATGAGGAAGCAGCACCATCGGC  
AAAACCATCACGCCCCAACAAAGGACTGTTGAAATTATTAATGTGGATGATGAGATTACGGGA  
GACACACCCATGGAAAGCCACCTGCCCATGCCTGCTATCGAGCATGAGCACCTAAATCACTA  
TAACTCATACAAATCTCCCTTCAACCACACAACAACAGTTAACACAATAAATTCAATACACA  
GTTTCAGTGCATGAACCGTTATTGATCCGAATGAACTCTAAAGACAATGTACAAGAGACTCAA  
ATCTAAACATTTACAGAGTTACAAAAACAAACAATCAAAAAAAGACAGTTTATTAAAA  
ATGACACAAATGACTGGGCTAAATCTACTGTTTCAAAAAAGTGTCTTTACAAAAAACAAAA  
AAGAAAAGAAATTTATTATTAAAAATTCTATTGTGATCTAAAGCAGACAAAAA

## FIGURE 52

Signal peptide:	Amino acids 1-44
Transmembrane domain:	Amino acids 525-545
N-glycosylation sites:	Amino acids 278-282;364-368; 390-394;412-416;415-419; 434-438;442-446;488-492; 606-610
cAMP- and cGMP-dependent protein kinase phosphorylation site:	Amino acids 183-187
N-myristoylation sites:	Amino acids 40-46;73-79; 118-124;191-197;228-234; 237-243;391-397;422-428; 433-439;531-537

MLNKMTLHPQQIMIGPRFNRALFDPLLVLALLQLLVVAGLVRAQTCPSVCSCSNQFSKVIC  
VRKNLREVPDGI STNTRLLNLHENQIQI I KVNSFKHLRHLEILQLSRNHIRTIEIGAFNGLA  
NLNTLELFDNRLTTIPNGAFVYLSKLELWLRNNPIESIPSYAFNRIPSLRRLDLGELKRLS  
YISEGAFEGLSNLRYNLAMCNLREIPNLTPLIKLELDLSGNHLSAIRPGSFQGLMHLQKL  
WMIQSQIQVIERNAFDNLQSLVEINLAHNNLTLLPHDLFTPLHHLERIHLHNPWNCNDIL  
WLSWWIKDAPSNTACCARCNTPPNLKGRYIGELDQNYFTCYAPVIVEPPADLNVTEGMAAE  
LKCRASLTSVSWITPNGTVMTHGAYKVRIAVLSDGTLNFTNVTVQDTGMYTCMVNSVGN  
TTASATLNVTAATTTTFFSYFSTVTVETMEPSQDEARTTDNNVGPTPVVDWETTNVTTSLTPQ  
STRSTEKTFTIPVTDINSGIPGIDEVMKTTKIIIGCFVAITLMAAVMLVIFYKMRKQHHRQN  
HHAPTRTVEIINVDDEITGDTPMESHLPMPAIEHEHLNHYSYKSPFNHTTTVNTINSIHSS  
VHEPLLIRMNSKDNVQETQI



FIGURE 53

CACGCACTTCACCTGGGTCGGGATTCTCAGGTCATGAACGGTCCCAGCCACCTCCGGGCAGG  
GCGGGTGAGGACGGGGACGGGGCGTGTCCAACCTGGCTGTGGGCTCTTGAAACCCGAGCATGG  
CACAGCACGGGGCGATGGGCGCGTTTCGGGCCCTGTGCGGCCTGGCGCTGCTGTGCGCGCTC  
AGCCTGGGTGAGCGCCCCACCGGGGGTCCCGGGTGCGGCCCTGGGCGCCTCCTGCTTGGGAC  
GGGAACGGACGCGCGCTGCTGCCGGGTTACACGACGCGCTGCTGCCGCGATTACCGGGCG  
AGGAGTGCTGTTCCGAGTGGGACTGCATGTGTGTCCAGCCTGAATTCCACTGCGGAGACCCT  
TGCTGCACGACCTGCCGGCACCACCCTTGTCCCCCAGGCCAGGGGGTACAGTCCCAGGGGAA  
ATTCAGTTTTGGCTTCCAGTGTATCGACTGTGCCTCGGGGACCTTCTCCGGGGGCCACGAAG  
GCCACTGCAAACCTTGGACAGACTGCACCCAGTTTCGGGTTTTCTCACTGTGTTCCCTGGGAAC  
AAGACCCACAACGCTGTGTGCGTCCCAGGGTCCCCGCCGGCAGAGCCGCTTGGGTGGCTGAC  
CGTCGTCCTCCTGGCCGTGGCCGCTGCGTCCTCCTCCTGACCTCGGCCAGCTTGGACTGC  
ACATCTGGCAGCTGAGGAGTCAGTGCATGTGGCCCCGAGAGACCCAGCTGCTGCTGGAGGTG  
CCGCCGTGACCGAAGACGCCAGAAGCTGCCAGTTCCCCGAGGAAGAGCGGGGCGAGCGATC  
GGCAGAGGAGAAGGGGCGGCTGGGAGACCTGTGGGTGTGAGCCTGGCCGTCTCCTCCGGGGCCA  
CCGACCGCAGCCAGCCCCCTCCCCAGGAGCTCCCCAGGCCGCAGGGGCTCTGCGTTCTGCTCT  
GGGCCGGGCCCTGCTCCCCTGGCAGCAGAAGTGGGTGCAGGAAGGTGGCAGTGACCAGCGCC  
CTGGACCATGCAGTTC

## FIGURE 54

Signal peptide:	Amino acids 1-25
Transmembrane domain:	Amino acids 162-180
N-glycosylation site:	Amino acids 146-150
N-myristoylation sites:	Amino acids 5-11;8-14;25-31; 30-36;33-39;118-124;122-128; 156-162
Prokaryotic membrane lipoprotein lipid attachment site:	Amino acids 166-177
Leucine zipper pattern:	Amino acids 171-193

MAQHGAMGAFRALCGLALLCALSLGQRPTGGPGCGPGRLLLGTGTDARCCRVHTTRCCRDYP  
GEECCSEWDCMCVQPEFHCGDPCCTTCRHHPCPPGQGVQSQGKFSFGFQCIDCASGTFSGGH  
EGHCKPWTDCQFGFLTVFPGNKTHNAVCPGSPPAEPLGWLTVVLLAVAACVLLLTSAQLG  
LHIWQLRSQCMWPRETQLLLEVPPSTEDARSCQFP EEER GERSAEEKGRLGDLWV

FIGURE 55

GCTGTGGGAACCTCTCCACGCGCACGAACTCAGCCAACGATTTCTGATAGATTTTGGGAGT  
TTGACCAGAGATGCAAGGGGTGAAGGAGCGCTTCTACCGTTAGGGAACCTCTGGGGACAGAG  
CGCCCCGGCCGCTGATGGCCGAGGCAGGGTGCACCCAGGACCCAGGACGGCGTCGGGAAC  
CATACCATGCCCCGGATCCCCAAGACCCTAAAGTTCTGTCGTCGTCATCGTCGCGGTCCTGCT  
GCCAGTCCTAGCTTACTCTGCCACCACTGCCCCGCGAGGAGGAAGTTCCCCAGCAGACAGTGG  
CCCCACAGCAACAGAGGCACAGCTTCAAGGGGGAGGAGTGTCCAGCAGGATCTCATAGATCA  
GAACATACTGGAGCCTGTAACCCGTGCACAGAGGGTGTGGATTACACCAACGCTTCCAACAA  
TGAACCTTCTTGCTTCCCATGTACAGTTTGTAAATCAGATCAAAAACATAAAAGTTCCTGCA  
CCATGACCAGAGACACAGTGTGTCAAGTGTAAAGAAGGCACCTTCCGGAATGAAAACCTCCCCA  
GAGATGTGCCGGAAGTGTAGCAGGTGCCCTAGTGGGGAAGTCCAAGTCAGTAATTGTACGTC  
CTGGGATGATATCCAGTGTGTTGAAGAATTTGGTGCCAATGCCACTGTGGAAACCCAGCTG  
CTGAAGAGACAATGAACACCAGCCCCGGGGACTCCTGCCCCAGCTGCTGAAGAGACAATGAAC  
ACCAGCCCAGGGACTCCTGCCCCAGCTGCTGAAGAGACAATGACCACCAGCCCCGGGGACTCC  
TGCCCCAGCTGCTGAAGAGACAATGACCACCAGCCCCGGGGACTCCTGCCCCAGCTGCTGAAG  
AGACAATGACCACCAGCCCCGGGGACTCCTGCCTCTTCTCATTACCTCTCATGCACCATCGTA  
GGGATCATAGTTCTAATTGTGCTTCTGATTGTGTTTGTTGAAAGACTTCACTGTGGAAGAA  
ATTCCTTCCTTACCTGAAAGGTTCAAGTAGGCGCTGGCTGAGGGCGGGGGCGCTGGACACT  
CTCTGCCCTGCCTCCCTCTGCTGTGTTCCACAGACAGAAACGCCTGC

## FIGURE 56

Signal peptide:	Amino acids 1-29
Transmembrane domain:	Amino acids 240-259
N-glycosylation site:	Amino acids 77-81;140-144; 156-160
cAMP- and cGMP-dependent protein kinase phosphorylation site:	Amino acids 126-130
N-myristoylation sites:	Amino acids 56-62;72-78; 114-120;154-160;233-239

MARIPKTLKFVVVIVAVLLPVLAYSATTARQEEVPQQTVPAPQQQRHSFKGEECPAGSHRSEH  
TGACNPCTEGVDYTNASNNEPSCFPCTVCKSDQKHKSSCTMTRDTVCQCKEGTFRNENSP  
CRKCSRCPSGEVQVSNCTSWDDIQCVVEFGANATVETPAAEETMNTSPGTPAPAAEETMNTS  
PGTPAPAAEETMTTSPGTPAPAAEETMTTSPGTPAPAAEETMTTSPGTPASSHYLSCTIVGI  
IVLIVLLIVFV

FIGURE 57

CCGCGGGAACGCTGTCCTGGCTGCCGCCACCCGAACAGCCTGTCCTGGTGCCCCGGCTCCCT  
GCCCCGCGCCCAGTCAATGACCCTGCGCCCCCTCACTCCTCCCGCTCCATCTGCTGCTGCTGCT  
GCTGCTCAGTGCGGCGGTGTGCCGGGCTGAGGCTGGGCTCGAAACCGAAAGTCCCGTCCGGA  
CCCTCCAAGTGGAGACCCTGGTGGAGCCCCCAGAACCATGTGCCGAGCCCGCTGCTTTTGGA  
GACACGCTTCACATACACTACACGGGAAGCTTGGTAGATGGACGTATTATTGACACCTCCCT  
GACCAGAGACCCTCTGGTTATAGAACTTGGCCAAAAGCAGGTGATTCCAGGTCTGGAGCAGA  
GTCTTCTCGACATGTGTGTGGGAGAGAAGCGAAGGGCAATCATTCTTCTCACTTGGCCTAT  
GGAAAACGGGGATTTCCACCATCTGTCCCAGCGGATGCAGTGGTGCAGTATGACGTGGAGCT  
GATTGCACTAATCCGAGCCAACTACTGGCTAAAGCTGGTGAAGGGCATTTCCTCTGGTAG  
GGATGGCCATGGTGCCAGCCCTCCTGGGCCTCATTGGGTATCACCTATACAGAAAGGCCAAT  
AGACCCAAAGTCTCCAAAAGAAGCTCAAGGAAGAGAAACGAAACAAGAGCAAAAAGAATA  
ATAAATAATAAATTTTAAAAAACTTAAAAAAAAAAAAAAAAAAAA

## FIGURE 58

Signal peptide:	Amino acids 1-25
Transmembrane domain:	Amino acids 155-174
N-glycosylation site:	Amino acids 196-200
N-myristoylation site:	Amino acids 95-100
Amidation site:	Amino acids 119-123
FKBP-type peptidyl-prolyl cis-trans isomerase signature 1:	Amino acids 62-78;87-123; 128-141

MTLRPSLLPLHLLLLLLLLLSAAVCRAEAGLETESPVRTLQVETLVEPPEPCAEPAAFGDTLHI  
HYTGSLVDGRIIDTSLTRDPLVIELGQKQVIPGLEQSLDMCVGEKRRRAIPSHLAYGKRGF  
PPSVPADAVVQYDVELIALIRANYWLKLVKGILPLVGMAMVPALLGLIGYHLYRKANRPKVS  
KKKLKEEKRNKSKKK

FIGURE 59

CAGTTCTGAAATCAATGGAGTTAATTTAGGGAATACAAACCAGCCATGGGGGTGGAGATTGC  
CTTTGCCTCAGTGATTCTCACCTGCCTCTCCCTTCTGGCAGCAGGAGTCTCCCAGGTTGTTC  
TTCTCCAGCCAGTTCCAACCTCAGGAGACAGGTCCCAAGGCCATGGGAGATCTCTCCTGTGGC  
TTTGCCGGCCACTCATGAGAGTGTTTTTGTGTAAAGTATTTTTTTAGAATACTGTTGACTTCT  
TCATGATTTAATAACCATCCTTTGCGAAGTTTTATGAGGCTTTAGGGGAATGTCAACCCTCA  
AATTTTTGTTATACTAGATGGCTTCCATTTACCCACCACTATTTTAAGGTCCCTTTATTTTT  
AGGTTCAAGGTTCAATTTGACTTGAGAAAGTGCCCTTCTGCAGCTTCATTGATTTTGTTTATC  
TTCATATTAATTGTAACGATTAAAAAGAATAAGAGCACGCAGACCTCTAGGAGAATATTT  
TATCCCTGGGTGCCCCCTGACACATTTATGTAGTGATCCCACAAATGTGATTGTTAATTTAAA  
TGTTATTCTAATATTAGTACATTCAGTTGTGATGTAATATGAATAACCAGAATCTATTTCTT  
AAAAGTTTTGAGTATATTTTTCAACTAGATATTTGTATAGAAAGACTGAATAGTGATG

FIGURE 60

Signal peptide: Amino acids 1-24  
N-myristoylation site: Amino acids 2-8  
Homology region to immuno-globulin light chain:  
Amino acids 5-33

MGVEIAFASVILTCLSLAAGVSQVLLQPVPVTQETGPKAMGDLSCGFAGHS



FIGURE 61

AAACTTGACGCCATGAAAGATCCCGGTCCTTCCTGCCGTGGTGCTCCTCTCCCTCCTGGTGCT  
CCTCTGCCCAGGGAGCCACCCTGGGTGGTCCTGAGGAAGAAAGCACCATTGAGAATTATG  
CGTCACGACCCGAGGCCTTTAACACCCCGTTCTGAACATCGACAAATTGCGATCTGCGTTT  
AAGGCTGATGAGTTCCTGAACTGGCACGCCCTCTTTGAGTCTATCAAAAGGAACTTCCTTT  
CCTCAACTGGGATGCCTTTCCTAAGCTGAAAGGACTGAGGAGCGCAACTCCTGATGCCCAGT  
GACCATGACCTCCACTGGAAGAGGGGGCTAGCGTGAGCGCTGATTCTCAACCTACCATAACT  
CTTTCCTGCCTCAGGAACTCCAATAAAACATTTTCATCCAAA

## FIGURE 62

Signal peptide:	Amino acids 1-22
N-myristoylation sites:	Amino acids 22-28;90-96
Homology region to peroxidase:	Amino acids 16-48

MKIPVLPVAVLLSLLVLHSAQGATLGGPEEESTIENYASRPEAFNTPFLNIDKLRSFAKDE  
FLNWHALFESIKRKLPFLNWDAFPKLGKGLRSATPDAQ

## FIGURE 63

GCATTTTTGTCTGTGCTCCCTGATCTTCAGGTCACCACCAATGAAGTTCTTAGCAGTCCTGGT  
ACTCTTGGGAGTTTCCATCTTTCTGGTCTCTGCCCAGAATCCGACAACAGCTGCTCCAGCTG  
ACACGTATCCAGCTACTGGTCCTGCTGATGATGAAGCCCCTGATGCTGAAACCACTGCTGCT  
GCAACCACTGCGACCACTGCTGCTCCTACCACTGCAACCACCGCTGCTTCTACCACTGCTCG  
TAAAGACATTCCAGTTTTACCCAAATGGGTTGGGGATCTCCCGAATGGTAGAGTGTGTCCCT  
GAGATGGAATCAGCTTGAGTCTTCTGCAATTGGTCACAACCTATTCATGCTTCCTGTGATTTC  
ATCCAACCTACTTACCTTGCCTACGATATCCCCTTTATCTCTAATCAGTTTATTTTCTTTCAA  
ATAAAAAATAACTATGAGCAACATAAAAAAAAAAAAAA

FIGURE 64

Signal peptide: Amino acids 1-19  
Protein Kinase C phosphorylation site: Amino acids 68-70

MKFLAVLVLLGVSI FLVSAQNPTTAAPADTYPATGPADDEAPDAETTAAATTATTAAPTTAT  
TAASTTARKDIPVLPKWVGDL PNGRVCP

FIGURE 65

TGGCCTCCCCAGCTTGCCAGGCACAAGGCTGAGCGGGAGGAAGCGAGAGGCATCTAAGCAGG  
CAGTGTTTTGCCTTCACCCCAAGTGACCAATGAGAGGTGCCACGCGAGTCTCAATCATGCTCC  
TCCTAGTAACTGTGTCTGACTGTGCTGTGATCACAGGGGCCTGTGAGCGGGATGTCCAGTGT  
GGGGCAGGCACCTGCTGTGCCATCAGCCTGTGGCTTCGAGGGCTGCGGATGTGCACCCCGCT  
GGGGCGGGAAGGCGAGGAGTGCCACCCCGGCAGCCACAAGGTCCCCTTCTTCAGGAAACGCA  
AGCACCACACCTGTCCTTGCTTGCCCAACCTGCTGTGCTCCAGGTTCCCGGACGGCAGGTAC  
CGCTGCTCCATGGACTTGAAGAACATCAATTTTTTAGGCGCTTGCTGCTCAGGATACCCA  
CCATCCTTTTCCTGAGCACAGCCTGGATTTTTATTTCTGCCATGAAACCCAGCTCCCATGAC  
TCTCCCAGTCCCTACACTGACTACCCTGATCTCTCTTGTCTAGTACGCACATATGCACACAG  
GCAGACATACCTCCCATCATGACATGGTCCCCAGGCTGGCCTGAGGATGTCACAGCTTGAGG  
CTGTGGTGTGAAAGGTGGCCAGCCTGGTTCTCTTCCCTGCTCAGGCTGCCAGAGAGGTGGTA  
AATGGCAGAAAGGACATTCCCCCTCCCCTCCCCAGGTGACCTGCTCTCTTTCTGGGCCCTG  
CCCCCTCTCCCCACATGTATCCCTCGGTCTGAATTAGACATTCTCTGGGCACAGGCTCTTGGGT  
GCATTGCTCAGAGTCCCAGGTCTGGCCTGACCCTCAGGCCCTTCACGTGAGGTCTGTGAGG  
ACCAATTTGTGGGTAGTTCATCTTCCCTCGATTGGTTAACTCCTTAGTTTCAGACCACAGAC  
TCAAGATTGGCTCTTCCCAGAGGGCAGCAGACAGTCACCCCAAGGCAGGTGTAGGGAGCCCA  
GGGAGGCCAATCAGCCCCCTGAAGACTCTGGTCCCAGTCAGCCTGTGGCTTGTGGCCTGTGA  
CCTGTGACCTTCTGCCAGAATTGTCATGCCTCTGAGGCCCCCTCTTACCACACTTTACCAGT  
TAACCACTGAAGCCCCCAATTCCCACAGCTTTTCCATTAAAATGCAAATGGTGGTGGTTCAA  
TCTAATCTGATATTGACATATTAGAAGGCAATTAGGGTGTTCCTTAAACAACCTCCTTTCCA  
AGGATCAGCCCTGAGAGCAGGTTGGTGACTTTGAGGAGGGCAGTCCTCTGTCCAGATTGGGG  
TGGGAGCAAGGGACAGGGAGCAGGGCAGGGGCTGAAAGGGGCACTGATTACAGACCAGGGAGG  
CAACTACACACCAACATGCTGGCTTTAGAATAAAAGCACCAACTGAAAAAA

## FIGURE 66

Signal peptide: Amino acids 1-19

Tyrosine kinase phosphorylation site:

Amino acids 88-95

N-myristoylation sites:

Amino acids 33-39;35-41;

46-52

MRGATRVSIMLLLVTVSDCAVITGACERDVQCGAGTCCAISLWLRGLRMCTPLGREGEECHP  
GSHKVPFFRKRKHHTCPCLPNLLCSRFPDGRYRCSMDLKNINF

FIGURE 67A

AAGCAACCAAAGCTGCAAGCTTTGGGAGTTGTTGCTGTCCCTGCCCTGCTCTGCTAGGGAGA  
GAACGCCAGAGGGAGGCGGCTGGCCCGGCGGCAGGCTCTCAGAACCGCTACCGGCGATGCTA  
CTGCTGTGGGTGTCGGTGGTTCGACGCTTGGCGCTGGCGGTACTGGCCCCGGAGCAGGGGA  
GCAGAGGCGGAGAGCAGCCAAAGCGCCCAATGTGGTGTGGTTCGTGAGCGACTCCTTCGATG  
GAAGGTAAACATTTTCATCCAGGAAGTCAGGTAGTGAACTTCCTTTTATCAACTTTATGAAG  
ACACGTGGGACTTCCTTTCTGAATGCCTACACAACTCTCCAATTTGTTGCCCATCACGCGC  
AGCAATGTGGAGTGGCCTCTTCACTCACTTAACAGAATCTTGGAATAATTTTAAGGTCTAG  
ATCCAAATTATACAACATGGATGGATGTCATGGAGAGGCATGGCTACCGAACACAGAAATTT  
GGGAACTGGACTATACTTCAGGACATCACTCCATTAGTAATCGTGTGGAAGCGTGGACAAG  
AGATGTTGCTTTCTTACTCAGACAAGAAGGCAGGCCCATGGTTAATCTTATCCGTAACAGGA  
CTAAAGTCAGAGTGATGGAAAGGGATTGGCAGAATACAGACAAAGCAGTAACTGGTTAAGA  
AAGGAAGCAATTAATTACACTGAACCATTTGTTATTTACTTGGGATTAAATTTACCACACCC  
TTACCCTTCACCATCTTCTGGAGAAAATTTGGATCTTCAACATTTACACATCTCTTTATT  
GGCTTGAAAAAGTGTCTCATGATGCCATCAAAATCCCAAAGTGGTCACCTTTGTCAGAAATG  
CACCTGTAGATTATTACTCTTCTTATACAAAAAAGTCACTGGAAGATTTACAAAAAAGA  
AATTAAGAATATTAGAGCATTTTATTATGCTATGTGTGCTGAGACAGATGCCATGCTTGGTG  
AAATTATTTTGGCCCTTCATCAATTAGATCTTCTTCAGAAAAGTATTGTCATATACTCCTCA  
GACCATGGAGAGCTGGCCATGGAACATCGACAGTTTTTATAAATGAGCATGTACGAGGCTAG  
TGCACATGTTCCGCTTTTGATGATGGGACCAGGAATTAAGCCGGCCTACAAGTATCAAATG  
TGGTTTCTCTTGTGGATATTTACCCTACCATGCTTGATATTGCTGGAATTCCTCTGCCTCAG  
AACCTGAGTGGATACTCTTTGTTGCCGTTATCATCAGAAACATTTAAGAATGAACATAAAGT  
CAAAAACCTGCATCCACCCTGGATTCTGAGTGAATTCATGGATGTAATGTGAATGCCTCCA  
CCTACATGCTTCGAACTAACCCTGGAAATATATAGCCTATTTCGGATGGTGCATCAATATTG  
CCTCAACTCTTTGATCTTTCCTCGGATCCAGATGAATTAACAAATGTTGCTGTAAAATTTCC  
AGAAATTACTTATTCTTTGGATCAGAAGCTTCATTCCATTATAAACTACCCTAAAGTTTCTG  
CTTCTGTCCACCAGTATAATAAAGAGCAGTTTATCAAGTGGAAACAAAGTATAGGACAGAAT  
TATTCAAACGTTATAGCAAATCTTAGGTGGCACCAAGACTGGCAGAAGGAACCAAGGAAGTA  
TGAAAATGCAATTGATCAGTGGCTTAAACCCATATGAATCCAAGAGCAGTTTGAACAAAAA  
GTTTAAAAATAGTGTTCTAGAGATACATATAAATATATTACAAGATCATAATTATGTATTTT  
AAATGAAACAGTTTTTAATAATTACCAAGTTTTGGCCGGGCACAGTGGCTCACACCTGTAATC  
CCAGGACTTTGGGAGGCTGAGGAAAGCAGATCACAAGGTCAAGAGATTGAGACCATCCTGGC

FIGURE 67B

CAACATGGTGAAACCCCTGTCTCTACTAAAAATACAAAAATTAGCTGGGCGCGGTGGTGCACA  
CCTATAGTCTCAGCTACTCAGAGGCTGAGGCAGGAGGATCGCTTGAACCCGGGAGGCAGCAG  
TTGCAGTGAGCTGAGATTGCGCCACTGTACTCCAGCCTGGCAACAGAGTGAGACTGTGTTCGC  
AAAAAATAAAAAATAAATAATAATTACCAATTTTTCATTATTTTGTAAGAATGTAGTG  
TATTTTAAGATAAAATGCCAATGATTATAAAATCACATATTTTCAAAAATGGTTATTATTTA  
GGCCTTTGTACAATTTCTAACAATTTAGTGGAAGTATCAAAAGGATTGAAGCAAATACTGTA  
ACAGTTATGTTCCTTTAAATAATAGAGAATATAAAATATTGTAATAATATGTATCATAAAAT  
AGTTGTATGTGAGCATTTGATGGTGAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA  
AAA



## FIGURE 68

Signal peptide: Amino acids 1-15  
Transmembrane domain: Amino acids 347-365  
N-glycosylation sites: Amino acids 108-112;166-170;  
193-197;262-266;375-379;  
413-417;498-502  
Tyrosine kinase phosphorylation sites:  
Amino acids 103-110;455-462  
N-myristoylation sites: Amino acids 104-110;344-350;  
348-354;409-415;496-502  
Homology block to sulfatase proteins:  
Amino acids 286-315;359-369;  
78-97

MLLLWVSVVAALALAVLAPGAGEQRRRAAKAPNVVLVSDSFDGRLTFHPGSQVVKLPFINF  
MKTRGTSFLNAYTNSPICCPSRAAMWSGLFTHLTESWNNFKGLDPNYTTWMDVMERHGYRTQ  
KFGKLDYTSGHHSISNRVEAWTRDVAFLLRQEGRPMVNLIRNRTKVRVMERDWQNTDKAVNW  
LRKEAINYTEPFVIYLG LNLPHYPSPSSGENFGSSTFHTSLYWLEKVS HDAIKIPKWSPLS  
EMHPVDYYSSYTKNCTGRFTKKEIKNIRAFYYAMCAETDAMLGEIILALHQDLLQKTIVIIY  
SSDHGELAMEHRQFYKMSMYEASAHVPLLMMGPGIKAGLQVSNVSLVDIYPTMLDIAGIPL  
PQNLSGYSLPLSSETFKNEHKVKNLHPPWILSEFHGCNVNASTYMLRTNHWKYIAYS DGAS  
ILPQLFDLSSDPDEL TNVAVKFPEITYSLDQKLHSIINYPKVSASVHQYNKEQFIKWKQSIG  
QNYSNVIANLRWHQDWQKEPRKYENAI DQWLKTHMNPRAV